



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>

NYPL RESEARCH LIBRARIES



3 3433 06641779 5









1

HANDY-BOOK OF ROCK NAMES.



HANDY-BOOK

OF

ROCK NAMES,

WITH

OF DESCRIPTIONS OF THE ROCKS.

BY
George
 G. HENRY KINAHAN,

DIPLOMATIST, TRIN. COLL. DUBLIN; MEMBER OF THE ROYAL IRISH
 ACADEMY; FELLOW OF THE ROYAL GEOLOGICAL SOCIETY, IRELAND;
 OF THE GEOLOGICAL SURVEY OF THE UNITED KINGDOM;
 ETC. ETC.



NEW
 EDITION
 1873

LONDON:
 ROBERT HARDWICKE, 192, PICCADILLY.
 1873. *hr*

GE



STON WOOD
CLUB
TREASURY

P R E F A C E.

A WANT expressed by many Geologists and often felt by the Author, is a classified **HANDY-BOOK OF ROCK NAMES**. It is hoped that, until a better appears, this Manual may be useful to the Public.

To write such a work satisfactorily is no easy task. In other branches of Natural Science, such as Botany, Zoology, and the like, some systematic nomenclature has been followed, but in Geology no one system has been used, or all system has been ignored. It is not uncommon to find eminent Petrologists calling the same rock by several names, or, what is more perplexing, using for a rock a name already given to a totally different one. At the present time Rock names are in such confusion that if an observer names a rock, without at the same time mentioning his authority, he may be supposed to be referring to a rock, or even rocks, totally different to that which he intended to illustrate. This irregularity has caused many names to be used, not only unscientifically but also absurdly, as the original name may refer to a mineral or character not possessed by the rock or rocks to which it is now applied. From the above it is evident that a compiler of such a book as this *Manual must run contrary to the opinions of*

prejudices of many geologists; therefore some will find fault even if the work could be well done.

I would wish to impress on my readers that this book has been written as a reference for surveyors and students while engaged in the field. Should they desire a perfect knowledge of rocks, they must study the various papers and works of the different eminent Petrologists. The work, necessarily, is far from perfect: still I hope it may be of some small use or help to learners of Geology.

During the compilation of this Manual, the writings of numerous Petrologists have been consulted and quoted; also works on general subjects from which information could be procured. Dana's suggestions as to the termination of Rock names have been adopted as far as practicable, while the older names for rocks are generally adopted, except when they are objectionable or better names have since been proposed. Such local names as were known are also given, as they may assist explorers in gleaning information about a country. To various fellow-labourers I am much indebted: to D. Forbes, F.R.S., &c., for information; also to W. King, Dep. Supt. Geol. Surv., India, and for his valuable assistance while arranging and classifying the rocks. I should also mention the names of the Rev. M. H. Close, M.R.I.A., &c.; Stackpoole Westropp, M.D., M.R.I.A., &c.; and H. Leonard, M.R.I.A.; besides others who have supplied me with lists of local names.

G. H. K.

CONTENTS.

PREFACE	Page iii
INTRODUCTORY REMARKS	1
PART I.—CLASSIFIED TABLE OF THE ROCKS	9
Order I. Ingenite Rocks	9
Class 1. Granitic Rocks	9
Class 2. Plutonic Rocks	10
Class 3. Volcanic Rocks	14
Class 4. Transition or Metamorphic Sedimentary Rocks	15
Order II. Derivate Rocks	19
Class 1. Subaqueous Rocks	19
Class 2. Subaërial Rocks	24
PART II.—Order I. INGENITE ROCKS	26
Class 1. Granitic Rocks	26
Group A. Intrusive Granite	27
Group B. Granite for the most part non-intrusive	32
Group C. Protogene...	36
Group D. Elvanyte	39
Class 2. Plutonic Rocks	43
Group E. Felstone	44
Group F. Whinstone	55
Group G. Rocks due to pseudomorphic action	63
Group H. Tuff	66
Class 3. Volcanic Rocks	69
Group I. Trachytic	70
Group J. Augitic	73
Group K. Tufa and Peperino	75
Class 4. Transition Rocks	76
Group L. Gneiss	77
Group M. Schist	80

Group N. Slate	Page	88
Group O. Calcareous Rocks		88
Group P. Pseudomorph Calcareous Rocks		90
PART III.—Order II. DERIVATE ROCKS		94
Class 1. Subaqueous Rocks		94
Group A. Rocks for the most part mechanically formed		94
Subgroup A. Arenaceous Rocks		94
Subgroup B. Argillous Rocks		94
Subgroup C. Fault-rock		100
Group B. Rocks for the most part chemically formed		106
Subgroup A. Haylyte		107
Subgroup B. Gypsum		107
Subgroup C. Anhydrite		108
Subgroup D. Dolomite		108
Subgroup E. Quartz		108
Subgroup F. Limonite Rock		109
Subgroup G. Hematite Rock		110
Subgroup H. Sphaerosiderite		110
Subgroup I. Minerals occurring as Rock Masses		111
Group C. Rocks partly mechanically, partly chemically, and partly organically formed		112
Subgroup A. Laminated Coal		112
Subgroup B. Limestone		115
Subgroup C. Dolomite		120
Class 2. Subaërial Rocks		122
Group D. Mechanically, chemically, and organically formed		122
Subgroup A. Coal		122
Subgroup B. Surface Deposit and Accumulations		126

HANDY-BOOK OF ROCK NAMES.

INTRODUCTORY REMARKS.

THE term Rock, in a geological sense, includes every solid substance that is an ingredient, or forms part, of the earth. Thus loose sand, clay, peat, and even vegetable mould, geologically speaking, are rocks. Jukes thus defines a rock:—"A mass of mineral matter consisting of many individual particles, either of one species of mineral, or of two or more species of minerals, or of fragments of such particles. These particles need not at all resemble each other in size, form, or composition; while, neither in its minute particles, nor in the external shape of the mass, need a rock have any regular symmetry of form." Rocks are most variable in condition and structure; soft or hard, loose or compact, friable or tenacious, coarse or fine, crystalline or homogeneous; or they may be scoriaceous, vesicular, hyaline, &c. &c.

Rocks may be chemically, mechanically, or organically formed, or two or more of these combined; they may be stratified or unstratified, igneous or aqueous, or partaking of the nature of both. Various classifications have been adopted by different writers on the subject, each taking dif-

1

HANDY-BOOK OF ROCK NAMES.



A HANDY-BOOK
OF
ROCK NAMES,
WITH
BRIEF DESCRIPTIONS OF THE ROCKS.

BY
George
G. HENRY KINAHAN,

ENGINEER DIPLOMATIST, TRIN. COLL. DUBLIN; MEMBER OF THE ROYAL IRISH
ACADEMY; FELLOW OF THE ROYAL GEOLOGICAL SOCIETY, IRELAND;
OF THE GEOLOGICAL SURVEY OF THE UNITED KINGDOM;
ETC. ETC.



NEW YORK
PUBLISHED BY
J. B. LIPPINCOTT & CO.
15 N. 2ND ST.

LONDON:
ROBERT HARDWICKE, 192, PICCADILLY.

1873. *hr*

E. 4

vate rocks as Dolomyte, Ophyte, Steatyte, &c., in which a secondary or pseudomorphic action has taken place—new minerals replacing the old constituents, and thereby changing the nature of the rock.

The second Order (DERIVATE ROCKS) may be divided into two classes; namely, I. SUBAQUEOUS, II. SUBAERIAL, which need not be dwelt upon now.

This Manual is divided into three parts. Part I. is a classified table of the rocks; Part II. is a description of the Ingenite rocks; and Part III. is a description of the Derivate rocks: while in the Index will be found an alphabetically arranged list of local, duplicate, and other rock names that do not appear *in the classified table of the rocks*. As before stated, Dana's termination of *yte* for rock names, on account of its convenience in distinction, is adopted; but at the same time the names ending in *ite* are also given for such as prefer them. The termination *oid* is only used to signify like, as *Granitoid*, granite-like; *ous* means a constituent that gives a character to the rock, as *Pyritous granite*, granite containing pyrite; while the termination *ic* points to quantity, as *felsitic*, containing a quantity of felsite (orthoclase intimately combined with quartz.—See quotation from Cotta, page 46).


In Parts II. and III. a general description of each class is given, and a general description of each group; while under each group will be found the subgroups, and under the latter the varieties, and, when necessary, the subvarieties. In these two parts names of classes are printed in large capitals, of groups in egyptian type, of subgroups *and particular varieties* in small capitals; while the

others, subvarieties and local names, are in italics. Synonymous names for groups, subgroups, and varieties are given when not highly objectionable, but the name that is considered most applicable appears first. Objectionable names, if mentioned, are printed in italics, and the objection to them pointed out.

All the groups are arranged under roman capitals; the subgroups under italic capitals; the varieties under small italic, and the subvarieties under small roman letters.

A student wishing to learn the description of a rock, must first look for the name in the Index, and if the name does not occur there, it will be found in the classified list in its proper group, class, and order; he must then refer for the description to either Parts II. or III.

NOTE.—While this Manual was in the press, Professor Joseph Le Conte's excellent paper on the "Features of the Earth's Surface" was published in "The American Journal of Science and Art," third series, vol. iv. As in it the formation of metamorphic granite is explained (p. 468), the student may be referred to it.



PART I.

CLASSIFIED TABLE OF THE ROCKS.

Order No. I.—*INGENITE ROCKS.*

(*Divided into Classes, Groups, Subgroups, and Varieties.*)

Class I.—GRANITIC ROCKS,

Group A.—INTRUSIVE GRANITE (Highly siliceous granite).

Subgroup A. Pyritous intrusive granite.

Var. *a.* Beresyte.

B. Albitic granite.

C. Schorlaceous granite.

D. Granityte.

E. Felsitic granite.

F. Greissen, or quartzitic granite.

G. Porphyritic intrusive granite.

H. Pegmatyte.

Var. *a.* Graphic granite.

b. Blumen granite.

c. Plumose granite.

Group B. — GRANITE FOR THE MOST PART NON-INTRUSIVE (Basic or oligoclase granite).

Subgroup A. Hornblendic granite.

Var. *a.* Titanitic granite.

b. Chloritic granite.

c. Talcose granite.

d. Epidotic granite.

B. Graphitic granite.

C. Hematitic granite.

D. Pyritous oligoclase granite.

E. Porphyritic oligoclase granite.

F. Gneissoid granite.

- Var. *a.* Nodular granite.
- b.* Lenticular granite.

- Subgroup *G.* Intrusive oligoclase granite.
- H.* Granitic felstone.

Group C.—PROTOGENE.

Group D.—ELVANYTE, or QUARTZ PORPHYRY.

Subgroup *A.* Felsitic and felspathic elvanyte.

- Var. *a.* Orthoclastic elvanyte.
- b.* Oligoclastic elvanyte.
- c.* Rappakivi elvanyte.

- B.* Quartzitic elvanyte.
- C.* Micaceous elvanyte.
- D.* Pyritous elvanyte.
- E.* Hornblendic elvanyte.

- Var. *a.* Micaceous hornblendic elvanyte.
- b.* Chloritic.
- c.* Felspathic.
- d.* Pyritous.

- F.* Chloritic elvanyte.
- G.* Granular elvanyte.
- H.* Compact elvanyte.
- I.* Gneissoid elvanyte.
- J.* Granitoid elvanyte.
- K.* Pyromeride.

Class II.—PLUTONIC ROCKS.

Group E.—FELSTONE, or FELSYTE.

Subgroup *A.* Petrosilex.

- Var. *a.* Felstone glass.
- b.* Ribaned petrosilex.

B. Quartzitic felstone.

- Var. *a.* Oligoclastic quartzitic felstone.
- b.* Micaceous.
- c.* Hornblendic.
- d.* Chloritic.
- e.* Pyritous.
- f.* Striped.
- g.* Variolitic.

Subgroup C. Minette.

Var. *a.* Fraidronyte.

D. Euryte.

Var. *a.* Orthoclastic euryte.*b.* Oligoclastic.*c.* Micaceous.*d.* Hornblendic.*e.* Chloritic.*f.* Quartzose.*g.* Euryte porphyry.*h.* Amygdaloidal euryte.*i.* Vesicular.*j.* Compact.*k.* Slab.*l.* Cleaved.*E.* Compact felstone.*F.* Granular felstone.*G.* Friable felstone.*H.* Rubbly felstone.*I.* Cleaved felstone.*J.* Foliated felstone.*K.* Granulyte.*L.* Felstone porphyry.Var. *a.* Amygdaloidal felstone.*b.* Vesicular felstone.

M. Gneissyte.

Var. *a.* Quartzose gneissyte.*b.* Felsitic.*c.* Micaceous.*d.* Hornblendic.*e.* Syenyte gneissyte.*f.* Chloritic.*g.* Talcose.*h.* Porphyritic.*i.* Schistoid.*N.* Schistoid felstone.

Group F.—WHINSTONE.

Subgroup A. Doleryte.

Var. *a.* White rock.*b.* Aphanyte-doleryte.

B. Melaphyre.

Var. *a.* Eukryte.*b.* Magnetic mel

- Var. *c.* Micaceous.
- d.* Porphyritic.
- e.* Amygdaloidal.
- f.* Vesicular.
- g.* Variolitic.

Subgroup *C.* Pyroxenite.*D.* Diabase.

- Var. *a.* Pyroxenic diabase.
- b.* Uralitic.
- c.* Labradoritic.
- d.* Oligoclastic.
- e.* Magnetic.
- f.* Kersantite.
- g.* Calcareous diabase.
- h.* Porphyritic.
- i.* Amygdaloidal.
- j.* Vesicular.
- k.* Variolitic.
- l.* Aphanite-diabase.

E. Gabbro.

- Var. *a.* Euphotide.
- b.* Noryte.
- c.* Hyperite.

F. Dioryte.

- Var. *a.* Amphibolyte.
- b.* Aphanite-dioryte.
- c.* Kersanton.
- d.* Napoleonyte.

G. Syenite.

- Var. *a.* Quartzose syenite.
- b.* Titanitic.

H. Hornblende rock.

- Var. *a.* Actinolite rock.
- b.* Tremolite rock.
- c.* Hyperite.
- d.* Dioryte.
- e.* Syenite.
- f.* Falso-syenite.
- g.* Mico-hornblende rock.

Group *G.*—ROCKS DUE TO PSEUDOMORPHIC ACTION OR WEATHERING (Ingenite in part, derivate in part).Group *A.* Ophyte.

- Var. *a.* Ophihornblende rock.

B. Steatite.

Var. *a.* Felspathic steatyte.

b. Pyroxenic.

c. Amphibolic.

Subgroup *C.* Eklogyte.

D. Epidosyte.

E. Kaolin.

F. Fuller's earth.

G. Meerschaum.

H. Magnesyte.

Group *H.*—**Tuff** (Ingenite in part, derivate in part).

Subgroup *A.* Felstone tuff.

Var. *a.* Quartzose.

b. Calcareous.

c. Pyritous.

d. Cupriferous.

e. Hematitic.

B. Whinstone, or Basic tuff.

Var. *a.* Hornblendic.

b. Pyroxenic.

c. Chloritic.

d. Calcareous.

e. Pyritous.

f. Cupriferous.

g. Hematitic.

C. Calcareous tuff.

D. Arenaceous.

E. Argillous.

F. Hematitic.

G. Pyritous.

H. Cupriferous.

I. Steatitic.

J. Porphyritic.

K. Nodular.

Var. *a.* Concretionary tuff.

b. Spheroidal.

L. Shaly tuff.

Var. *a.* Flaggy.

b. Slaty.

M. Conglomeritic tuff.

Var. *a.* Brecciated tuff.

b. Plutonic agglomerate.

Class III.—VOLCANIC ROCKS.

Group I.—TRACHYTIC GROUP.

Subgroup A. Rhyolyte.

- Var. *a.* Liparyte.
- b.* Perlyte.
- c.* Obsidian.
- d.* Pumice.
- e.* Phonolyte.

B. Trachyte.

- Var. *a.* Sanidine trachyte.
- b.* Drachenfels trachyte.
- c.* Domyte.
- d.* Andesyte.
- e.* Trachydoleryte.
- f.* Granular trachyte.
- g.* Compact trachyte.
- h.* Trachyte porphyry.
- i.* Trachyte lava.
- j.* Alumstone.

Group J.—AUGITIC GROUP.

Subgroup A. Augyte, or Doleryte.

- Var. *a.* Nephelite augyte.
- b.* Häüynophyre.
- c.* Allogovyte.
- d.* Common augyte.
- e.* Anamesyte.
- f.* Basalt.
- g.* Porphyritic basalt.
- h.* Amygdaloidal basalt.
- i.* Basalt lava.
- j.* Variolitic basalt.

B. Leucityte.

- Var. *a.* Compact leucityte.
- b.* Leucitophyre.
- c.* Amygdaloidal leucityte.
- d.* Leucityte lava.

Group K.—VOLCANIC TUFFA AND PEPERINO (Ingenite in part, derivate in part).

Subgroup A. Tufa.

- Var. *a.* Trachyte.
- b.* Pumiceous.
- c.* Phonolyte.
- d.* Pozzuolana.

B. Peperino.

- Var. *a.* Augyte.
- b.* Leucityte.
- c.* Palagonyte.

C. Volcanic conglomerate.

- Var. *a.* Volcanic breccia.
- b.* Volcanic agglomerate.

Class IV.—TRANSITION OR METAMORPHIC SEDIMENTARY ROCKS.

Group L.—GNEISS.**Subgroup A. Felspathic and felsitic gneiss.**

- Var. *a.* Orthoclase gneiss.
- b.* Oligoclase gneiss.
- c.* Alpynte.
- d.* Adularia gneiss.

B. Micaceous gneiss.

- Var. *a.* Two-mica gneiss.
- b.* White mica gneiss.
- c.* Black mica gneiss.

C. Quartzitic gneiss.**D. Basic gneiss.**

- Var. *a.* Hornblendic gneiss.
- b.* Chloritic gneiss.
- c.* Talcose gneiss.
- d.* Protogene gneiss.

E. Iolitic gneiss.**F. Hematitic gneiss.****G. Graphitic gneiss.****H. Common gneiss.**

- Var. *a.* Compact gneiss.
- b.* Slate gneiss.

- Var. *c.* Ribaned gneiss.
- d.* Fibrous gneiss.

Subgroup *I.* Oblique gneiss.

- Var. *a.* Curled gneiss.
- b.* Nodular gneiss.
- c.* Conglomeritic gneiss.

J. Porphyritic gneiss.

K. Granitoid gneiss.

Group *M.*—SCHIST.

Subgroup *A.* Argillyte.

- Var. *a.* Chistolithic schist.
- b.* Phyllitic schist.
- c.* Carbonaceous schist.
- d.* Alum schist.
- e.* Spilyte.
- f.* Dolomitic schist.
- g.* Folded argillyte.
- h.* Contorted argillyte.
- i.* Fibrous argillyte.

B. Quartzyte.

- Var. *a.* Itacolumyte.
- b.* Itabiryte.
- c.* Felsitic quartzyte.
- d.* Quartz rock.
- e.* Fibrous quartzyte.
- f.* Conglomeritic.

C. Felsyte schist.

- Var. *a.* Micaceous.
- b.* Quartzose.
- c.* Pyritous.
- d.* Ribaned.
- e.* Mealy.
- f.* Gneissoid.

D. Mica schist.

- Var. *a.* Two-mica schist.
- b.* Chloritic.
- c.* Talcose.
- d.* Garnetiferous.
- e.* Chistolithic.
- f.* Andalusitic.
- g.* Amphibolic.

- Var. *h.* Epidotic.
- i.* Quartzose.
- j.* Felsitic.
- k.* Calcareous.
- l.* Schorlaceous.
- m.* Pyritous.
- n.* Pyrrhotitic.
- o.* Hematitic.
- p.* Graphitic.
- q.* Fine.
- r.* Ribaned.
- s.* Fibrous.
- t.* Nummoid.
- u.* Wavy.
- v.* Frilled.
- w.* Curled.
- x.* Knotty.
- y.* Nodular.
- z.* Gneissoid.

Subgroup *E.* Basic schist.

- Var. *a.* Hornblende schist.
- b.* Actinolite.
- c.* Chlorite.
- d.* Chloritoid.
- e.* Potstone.
- f.* Talc schist.
- g.* Garnet.
- h.* Tourmaline.
- i.* Rhætitic.

F. Metallic schists.

- Var. *a.* Mico-iron schist.
- b.* Pyrite schist.
- c.* Pyrrhotite schist.

G. Burnt shale.

- Var. *a.* Burnt clay.
- b.* Rock slag.
- c.* Porcelanyte.

H. Minerals forming rock masses (*see* page 22).

- Var. *a.* Quartz.
- b.* Corundum.
- c.* Apatite.
- d.* Magnesite.

- Var. *e.* Orthoclase.
- f.* Pistacite.
- g.* Pycnite.
- h.* Lepidolite.
- i.* Lievrite.
- j.* Magnetite.
- k.* Anthracite.
- l.* Graphite.
- m.* Eisenkiesel.
- n.* Garnet-rock.

Group N.—SLATE. [See Order II., Class I., Group A, Subgroup *B*, Variety *d.*]

Group O.—CALCAREOUS ROCKS.

Subgroup *A.* Schistose limestone.

- Var. *a.* Cipollino.
- b.* Calciphyre.
- c.* Hemitrene.
- d.* Felsitic limestone.
- e.* Statuary marble.
- f.* Granular schistose limestone.

Group P.—PSEUDOMORPH CALCAREOUS ROCKS.

Subgroup *A.* Schistose dolomite.

- Var. *a.* Micaceous dolomite.
- b.* Pyritous.
- c.* Saccharoid.
- d.* Granular.

B. Ophyte.

- Var. *a.* Ophidolomite.
- b.* Ophimagnesyte.
- c.* Ophicalcyte.
- d.* Ophyte schist.

C. Steatyte.

- Var. *a.* Felsitic steatyte.
- b.* Hornblendic.
- c.* Pyroxenic.
- d.* Micaceous.
- e.* Steatyte schist.
- f.* Pyrallolyte.
- g.* Onkosin.

D. Magnesyte.

E. Smaragdite schist.

Order No. II.—*DERIVATE ROCKS.*

(*Divided into Classes, Groups, Subgroups, Varieties, and Subvarieties.*)

Class I.—SUBAQUEOUS ROCKS.

Group A.—ROCKS FOR THE MOST PART MECHANICALLY FORMED.

Subgroup A. Arenaceous rocks.

Var. *a.* Sand, gravel, and shingle.

b. Conglomerate.

Subvar. *a.* Calcareous.

b. Hematitic.

c. Cupriferous.

d. Pyritous.

e. Plumbeous.

f. Shale conglomerate.

g. Slate conglomerate.

c. Breccia.

Subvar. *a.* Calcareous.

b. Hematitic.

c. Cupriferous.

d. Pyritous.

e. Shale breccia.

d. Sandstone and grit.

Subvar. *a.* Calcareous.

b. Argillous.

c. Felspathic.

d. Micaceous.

e. Hematitic.

f. Cupriferous.

g. Pyritous.

h. Tuffose.

i. Pebbly.

j. Cleaved.

k. Oblique.

l. Concretionary.

m. Spheroidal.

n. Nodular.

o. Lenticular.

p. Friable.

Var. *e*. Flagstone.Subvar. *a*. Arenaceous.

b. Argillous.

c. Slabstone.

Subgroup *B*. Argillous or Argillaceous rocks.Var. *a*. Clay, loam, mud, and silt.Subvar. *a*. Kaolin.

b. Potter's clay.

c. Brick clay.

d. Fire clay.

e. Clunch.

f. Claystone.

g. Clayrock.

h. Lateryte.

i. Fuller's earth.

j. Bituminous clay.

k. Saliferous clay.

l. Variegated clay.

m. Hematitic clay.

n. Ferruginous clay.

o. Book clay.

b. Marl.Subvar. *a*. Calcareous.

b. Dolomitic.

c. Argillous.

d. Arenaceous.

e. Micaceous.

f. Oil-slate.

g. Glauconitic.

h. Gypseous.

i. Shell marl.

j. Mealy marl.

k. Marl-stone.

l. Tutenic marl.

m. Book marl.

c. Shale.Subvar. *a*. Calcareous.

b. Arenaceous.

c. Micaceous.

d. Bituminous.

e. Carbonaceous.

f. Hematitic.

g. Cupriferous.

Subvar. h. Flaggy.

- i. Conglomeritic.
- j. Nodular.
- k. Concretionary.
- l. Spheroidal.
- m. Lenticular.
- n. Mudstone.
- o. Tutenic shale.

Var. d. Slate.

Subvar. a. Roofing slate.

- b. Slab slate.
- c. Pencil slate.
- d. Novaculyte.
- e. Calcareous slate.
- f. Carbonaceous.
- g. Arenaceous.
- h. Tuffose.
- i. Ribaned.
- j. Conglomeritic.

Subgroup C. Fault rock.

Group B.—ROCKS FOR THE MOST PART CHEMICALLY FORMED.

Subgroup A. Halyte.

Var. a. Granular.

- b. Sparry.
- c. Fibrous.

B. Gypsum.

Var. a. Alabaster.

- b. Granular gypsum.
- c. Compact.
- d. Fibrous.
- e. Selenyte.
- f. Tripe-stone.

C. Anhydrite.

Var. a. Granular.

- b. Compact.
- c. Fibrous.

D. Dolomite.

E. Quartz.

Var. a. Chert.

- b. Flint.
- c. Menilite.
- d. Jasper.

Var. *e.* Agate.

f. BasanYTE.

Subgroup *F.* Limonite rock.

Var. *a.* Limonite shale.

b. Oolitic brown ore.

c. Reniform iron ore.

d. Pea-iron ore.

e. Bog-iron ore.

Subvar. *a.* Black.

b. Yellow.

c. Sparry.

G. Hematite rock.

Var. *a.* Red iron-mould.

b. Reddle.

c. Sparry red-ironstone.

d. Specular iron.

e. Topanhoacanga.

H. Sphaerosiderite.

Var. *a.* Black band.

I. Minerals occurring as rocks (*see* page 17).

Var. *a.* Asphalt.

b. Pyrite.

c. Cinnabar.

d. Boracite.

e. Trona.

f. Aragonite.

g. Manganese ores.

h. Rhodochrosite.

i. Bole.

j. Opal.

k. Fluorite.

l. Barite.

m. Ankerite.

n. Malachite.

o. Melanite.

p. Galmey.

q. Zincite.

r. Galenite.

s. Stibnite.

t. Arsenopyrite.

u. Sulphur.

v. Wavellite.

w. Chalcopyrite.

up C.—ROCKS PARTLY MECHANICALLY, PARTLY CHEMICALLY,
AND PARTLY ORGANICALLY FORMED.

ubgroup A. Laminated, or stratified coal.

Var. *a.* Cannel coal.

Subvar. *a.* Parrot coal.

b. Horn coal.

c. Torbanyte.

b. Splint coal.

c. Culm.

d. Bituminous shale.

e. Carbonaceous shale.

Subvar. *a.* Batt, or bass.

b. Dauks.

c. Kelve.

d. Pindy.

e. Slaty culm.

B. Limestone.

Var. *a.* Compact limestone.

b. Crystalline limestone.

c. Marble.

d. Oolyte.

Subvar. *a.* Pisolyte.

b. Dolomitic oolyte.

e. Chalk.

Subvar. *a.* Indurated chalk.

b. Chalk rock.

c. Pisolitic chalk.

d. Glauconitic chalk.

e. Ferruginous chalk.

f. Calcsinter.

Subvar. *a.* Travertine.

b. Stalactyte.

c. Stalagmyte.

d. Rockmeal.

g. Coral-reef limestone.

h. Brecciated limestone.

i. Limestone conglomerate.

j. Rubbly limestone.

k. Lithographyte.

l. Siliceous limestone.

m. Cherty.

n. Argillous.

o. Arenaceous.

- Var. *p.* Ferruginous.
- q.* Bituminous.
- r.* Stinkstone.
- s.* Dolomitic limestone.
- t.* Hydraulic limestone.

Subgroup *C.* Dolomite.

- Var. *a.* Granular.
- b.* Oolitic.
- c.* Compact.
- d.* Porous.
- e.* Cellular.
- f.* Concretionary.
- g.* Brecciated.
- h.* Rubbly.
- i.* Ferruginous.
- j.* Argillous.
- k.* Arenaceous.
- l.* Siliceous.
- m.* Cherty.
- n.* Tuffose.
- o.* Dolomitic sand.

D. Marine drift.

(See Class II., Subgroup *B*, Var. *c.*)

Class II.—SUBAERIAL ROCKS.

Group *D.*—MECHANICALLY, CHEMICALLY, OR ORGANICALLY FORMED.

Subgroup *A.* Coal.

Var. *a.* Peat.

- Subvar. *a.* White turf.
- b.* Brown turf.
- c.* Black or Stone turf.
- d.* Gas turf.

b. Lignite or Brown coal.

- Subvar. *a.* Woody lignite.
- b.* Compact lignite.
- c.* Jet.

c. Black coal or Steinkohle.

- Subvar. a. Caking coal.
- b. Non-caking coal.
- c. Cherry coal.
- Var. d. Anthracyte.
- Subvar. a. Native coke.
- Subgroup B. Surface deposits and accumulations.
- Var. a. Boulder-clay-drift.
- Subvar. a. Calcareous.
- b. Argillous.
- c. Arenaceous.
- b. Moraine drift.
- Subvar. a. Calcareous.
- b. Argillous.
- c. Arenaceous.
- d. Rocky.
- c. Sand, gravel, and shingle.
- Subvar. a. Esker drift.
- b. Post-drift gravel.
- c. Beaches.
- d. Æolian drift.
- e. Shell sand.
- f. Bergmehl.
- d. Guano.
- e. Soil.
- f. Meteoric drift.
- g. Ice.

PART II.

Order I.—INGENITE ROCKS.

Class I.—GRANITIC ROCKS.

THE typical rock of this class is a crystalline aggregate of quartz, feldspars, and micas, nearly universally known as granite or granyte [Celtic *gran*, Lat. *granum*, a grain.]* There are, however, rocks called granite in which part of the mica is replaced by other minerals, sometimes to such an extent that the mica is only microscopically visible.

Granite usually is a quaternary or quinary aggregate, containing, along with quartz, one or two feldspars and one or two micas; nevertheless, other minerals are often present, more especially pyrite and marcasite.† Some granites are undoubtedly intrusive, while others appear to be only in part intrusive, portions having been formed *in situ*; that is, having relations as to position with the associated rocks similar to those which now exist. Typical granites weather with a peculiar rough, rugged aspect.

* In Cornwall granite formerly was called *growan*, from *gronen*, a grain.

† Some of the granites that weather or disintegrate freely appear to have marcasite as an ingredient disseminated in *minute grains* throughout the mass.

A. Intrusive Granite; Highly Siliceous Granite; Leinster Granite; Oughterard Granite [intruded into the place it now occupies]. — An aggregate of quartz and orthoclase with black and white mica; pyrite and marcasite are often constituents; while Haughton seems to believe that most, if not all, intrusive granites also contain albite.

NOTE.—Haughton finds the intrusive granites of Cornwall, Devon, Leinster, and Ulster, to contain “quartz, orthoclase, margarodite, and lepidomelane.” He has also detected albite.

Varieties in Composition.

A. PYRITOUS INTRUSIVE, or HIGHLY SILICEOUS GRANITE.—When pyrite or marcasite is a component, galenite and chalcopyrite often occur as accessories.

a. Beresyte, Beresite.—Pyritous highly siliceous granite containing gold.

B. ALBITIC GRANITE (Dana).—Containing albite as well as orthoclase.

C. SCHORLACEOUS GRANITE. — With tourmaline (*schorl*) in addition to the mica.

In the highly siliceous or intrusive granite, as also in all granite, and some of the other ingenite rocks, are veins of segregation, which in the granites generally form two distinct varieties. One kind appears as irregular veins, nests, or patches, that have no deep-seated source, but die out every way, often in very short distances. The other variety makes regular, often dyke-like, veins, from half an inch or less in thickness, to about two or three yards in width. These latter appear to have segregated from the deep-seated fluid or semi-fluid portions of the mass, and to have been forced up into

the vacancies or the *shrinkage fissures* caused by the contraction consequent on the consolidation of the cooled portion or crust of the mass. The granite in such veins would necessarily be more siliceous than the granite mass through which they traverse, as basic rocks, although they melt at a lower temperature than the more siliceous, cool sooner and at a higher temperature, losing their heat much more rapidly. Consequently, it appears evident that the portions of a mass of granite which remain longest fluid ought to be more siliceous than the parts that cool first. The dyke-like *veins of segregation* are undoubtedly more siliceous than the rock which they traverse. Moreover, they are more compact, and finely crystalline. These latter peculiarities may, however, be due to the final rapid cooling when in the veins.

NOTE.—M'Farlane thus writes on the cooling of siliceous rocks:—"The scoria of iron-furnaces are usually very acid, containing as much as sixty per cent. of silex. They generally fuse at a temperature of 1,450° C. As they flow out of the breast of the furnace, they may be observed to do so very leisurely, to be sluggish and viscid, but nevertheless to continue fluid a long time; and even, in some cases, to flow out of the building in which they have been produced, before solidifying. On the other hand, slags from certain copper-furnaces, or from those used for puddling iron, are more or less basic, containing from thirty to forty-five per cent. of silica. As they flow out they are seen to be very fluid, and to run quickly, but they solidify much more rapidly than iron slag. Yet these basic slags fuse at 1,300° C., or about 150° less than the more acid slags. Those who have been accustomed to observe metallurgical processes will not find it difficult to conceive how a very siliceous slag might continue fluid at a temperature at which a more basic one might become solid."

They often are more or less felsytoid; that is, *have the aspect of a felstone or felsyte*; but they *always contain mica as a constituent*, although often

only visible under the lens or microscope. On account of their similitude in aspect to felsyte, also on account of their usual purplish reddish colour, Delesse and others erroneously call them "eurites," although they are quite dissimilar to the original *eurytes* of Daubuisson. Rose, on the other hand, describes a granite apparently identical with them, and calls it *granityte*, a name to which there seems to be no objection.

D. GRANITYTE ; GRANITITE (Rose).—A compact finely crystalline rock, usually purplish or reddish purple in colour. An aggregate of felspar (orthoclase), quartz, and mica. Black and white mica seem always to be present, but often in such minute flakes as to be only microscopically visible. Pyrite, and sometimes marcasite, are locally present.

Granityte* occurs as veins in granite from mere lines to two or more yards in thickness. It seems to be allied to *elvanyte* or *quartz-porphry*, hereafter described, as sometimes part of the quartz in the granityte seems to have crystallized out prior to the other constituents. Moreover, granityte nearly always weathers evenly like those rocks, and not with the rugged uneven surface, so characteristic of typical granite.

Allied to the intrusive granite are rocks that to the naked eye seem not mineralogically granite, yet petrologically they seem to be part of the intrusive granite, as they always occur associated

* If granityte, as suggested, fills cracks and fissures in the granite, it and *elvanyte* must necessarily have cooled under somewhat similar conditions. Jukes has described some of the granitytes of the counties of Wicklow and Dublin, Ireland, as *elvanytes*.

with it, into which they seem to merge, and it into them. They are as follows:—

E. FELSITIC GRANITE (King) ; FELSITE ROCK (Cotta).

—“ A rock of compact texture, about the hardness of felspar, with dull or smooth conchoidal or fissile fracture ; colour yellowish, reddish, grey, greyish, or bluish, weathering white.”—*Cotta*.

To the naked eye this rock does not appear to contain either mica or quartz ; however, with a lens, both of these substances can be detected, but usually they appear to be sparingly and partially developed.

F. GREISSEN ; QUARTZITIC GRANITE.—Apparently a crystalline granular aggregate of quartz and mica.

Greissen occurs associated with the intrusive granite, and merges into it. Sometimes, indeed, there is felspar developed, but so sparingly and at such wide intervals, that it seems to be more an accessory than an essential of the rock. In some places even the mica seems to be absent, or in such minute particles as only to be detected by a microscopical examination.

Structural Varieties.

G. PORPHYRITIC INTRUSIVE, OR HIGHLY SILICEOUS GRANITE.—In which crystals of felspar are largely and conspicuously developed.

H. PEGMATITE ; PEGMATITE [Gr. *pegma*, a hardened mass].—The second variety of the veins of segregation ; very coarsely and irregularly crystallized, the conspicuous constituents, according to Cotta, being “ orthoclase, quartz,

and silvery-white mica." Nevertheless, it usually also contains more or less dark-coloured mica, also pyrite or marcasite, and locally tourmaline.

Pegmatyte sometimes appears in large, well-marked veins, but more frequently it occurs in irregular veins, patches, and lentils, dying out in every direction. The dark-coloured mica, although the other constituents may be very coarsely crystalline, usually occurs in minute scales, but not always.

a. *Graphic granite* [Gr. *grapho*, I write].—A variety of Pegmatyte, in which the constituents are so arranged as to produce figures resembling written characters.

b. *Blumen granite*.—"The felspar assumes a form resembling flowering-stalks."—*Cotta*.

c. *Plumose granite*.—The mica assumes a plumose form, like Prince of Wales's feathers.
—*Jukes*.

The granites belonging to this type [intrusive granite] in general are not foliated; nevertheless Forbes mentions a "foliated granite" at Edisvand, Norway; of which he says: "No doubt could be entertained of its true eruptive origin." And my colleague, Mr. R. G. Symes, describes veins of a foliated granite which he observed N.E. of Castlebar, in the co. Mayo, Ireland. In south-west Mayo, associated with very slightly metamorphosed rock, are wide dykes of foliated rock, some being typical gneiss (leaves of quartz, felspar, and mica), others hornblendic gneiss, while some apparently are formed of leaves of quartz, orthoclase, and amphibole.

Not uncommon in the intrusive granite is a structure having an aspect somewhat like foliation; and yet it cannot be so classed, as the con

stituents of the rock have no tendency to occur in leaves. Perhaps it may possibly be some kind of close, irregular, rudely parallel-jointed structure, induced during the cooling of the rock, somewhat similar to the structure called by Scrope "shrinkage fissures." This structure of the intrusive granite is scarcely perceptible in a hard specimen, but in rock masses it is conspicuous.

Granites more or less due to metamorphism, having (for the most part) been altered into granite while in their present positions as regards the associated rocks (see note, page 7).

These rocks appear to have been formed from previously existing sedimentary and igneous rocks. Those due to the metamorphism of the sedimentary and of the basic-igneous rocks, are often very similar in aspect and composition, while some of the acid-igneous rocks have changed into granite somewhat similar to the *Intrusive Granite* (A).

B. Granite for the most part non-intrusive; Basic or Oligoclase Granite; Scandinavian Granite [in part metamorphic].—Usually a crystalline aggregate of quartz, black and white mica, orthoclase, and oligoclase. When typical, it is a quinary granite; however it is very variable in its composition, which in part seems due to its metamorphic origin. Many minerals occur locally as adjuncts; such as amphibole, titanite, ripidolite, pyrite, marcasite, &c., forming different varieties.

NOTE.—Houghton is of opinion that the normal constituents of this class of metamorphic granite in Ireland, Scotland, Norway, Sweden, and Finland, are *Quartz, Orthoclase, Oligoclase, Margarodite, and Lepidomelane*.

Varieties in Composition.

- A. **HORNBLENDIC GRANITE.**—An oligoclase granite, in which amphibole is a constituent.
- a. *Titanitic Granite ; Furbogh Granite.*
 - b. *Chloritic or Rhipidolitic Granite.*
 - c. *Talcose Granite.*
 - d. *Epidotic Granite.*

With amphibole titanite sometimes occurs, and this variety has been called *Furbogh Granite*, from a district in the county Galway, Ireland. In places rhipidolite, talc, or epidote may be present with, or in place of, the amphibole, making varieties. The last-named mineral is supposed to be due to secondary formation.

NOTE.—*Hornblendic granite* by some geologists has been called “*Syenitic granite*.” This name, however, seems not only unscientific, but also inapplicable, as the rock is not an aggregate of granite and syenite, but of granite and amphibole [*hornblende*], and it is to the latter mineral that the name is intended to direct attention. *Chloritic granite* and *Talcose granite* have erroneously been called “*Protogene granite*,” also a most incorrect name, as protogene (*quartz + felspar + talc*) is a constituent of neither. Furthermore, in general chloritic granite can have no relation to protogene, not containing any talc.

- B. **GRAPHITIC GRANITE.**—With graphite in addition to mica.
- C. **HEMATITIC OR FERRUGINOUS OLIGOCLASE GRANITE.**
—With micaceous iron-ore in addition to the mica.
- D. **PYRITOUS OLIGOCLASE GRANITE.** — Containing pyrite or marcasite, or even chalcopyrite, in addition to the other constituents.

Structural Varieties.

- E. PORPHYRITIC OLIGOCLEASE GRANITE; Galway Granite.*—Having one of the feldspars developed in crystals, larger and more conspicuous than those of the other constituents. Generally, it is the orthoclase that is porphyritically developed; however, sometimes it is the oligoclase. It is the typical granite of county Galway, Ireland.
- F. GNEISSOID GRANITE; FOLIATED GRANITE; Gneissic Granite (Cotta).*—An oligoclase granite, in which there is more or less distinct foliation. Sometimes one of the feldspars is porphyritically developed, when in general the mica does not curl round the large crystals of feldspar, but the latter lie promiscuously about.
- a. Nodular Granite.*—Containing round or semi-angular pieces or nodules of a dissimilar granite or of a granitic rock. The inliers often are a hornblendic granite or a rock, like minette, or like a gneissyte, but they vary considerably.
- b. Lenticular Granite.*—A more or less gneissoid granite, containing lenticular patches or lentils of granite or a rock, like a gneissyte or minette. The lentils sometimes occur closely together, at other times widely apart; but in all cases the foliation of the rock curls round them.
- G. INTRUSIVE OLIGOCLEASE GRANITE; Omev Granite.*—Very similar to the non-intrusive oligoclase granite; but such minerals as amphibole, titanite, ripidolite, &c., only occur locally, as accessories. It has been called "Omev granite," from an island on the west coast of Ireland.
- Besides the varieties mentioned above, others*

might be named, but those given seem to be the most important.

Veins of segregation occur in the oligoclase granite. Their composition appears very similar to the veins of segregation (*granityte*) that occur in the highly siliceous granite, being of a more or less siliceous nature. There are also irregular veins, nests, and lenticular patches, sometimes of considerable size, of a rock answering Cotta's description for the rock called *Pegmatyte*. The *Pegmatyte* found associated with the non-intrusive oligoclase granite is, at least in some instances, due to the metamorphism of irregular felspathic veins and lenticular patches that existed in the metamorphosed sedimentary and irruptive rocks prior to their being changed into granite.

Basic igneous rocks may be metamorphosed into rocks very similar to the *Basic* or *oligoclase granite*, or its varieties; and the *Acid* or highly siliceous igneous rocks may change into a rock like the finer varieties of the *Intrusive* or *highly siliceous granite* (A); while the basic felstones (*Euryte* and such rocks) may be altered into more or less hornblendic or chloritic, or even talcose granite.

H. GRANITIC FELSTONE.—A fine, more or less even-grained aggregate of quartz, felspar, and mica. In some varieties the felspar seems always to be orthoclase; the mica may be black or white, often both. Part of the mica may be replaced by amphibole or ripidolite. They weather neither like a true granite nor yet like a felstone, having a characteristic weathering of their own, partaking of the nature of both.

Granitic felstones probably are metamorphic *felstones*. They are often banded, the lines'

times being parallel, or nearly so, to the walls of the dykes, but at other times they are oblique. The bands are due to thin zones of different colour or texture, perhaps also of composition. In these bands, or between them, there is on rare occasions a structure like foliation.

All the metamorphosed igneous rocks are more or less granitoid, in accordance with the intensity of the metamorphism that has acted on them and the associated sedimentary rocks.

C. **Protogene** [Gr. *protogenes*, first formed; the rock, when first discovered, being supposed to be the oldest].—A granitoid rock; a crystalline aggregate of quartz, felspar, and talc.

Protogene, although not a true granite, seems allied to those of the *Oligoclase type* (B). The name is said to have been given originally to a metamorphic rock, but its original signification appears to have been lost, for Cotta thus describes it:—"A granite which contains talc or chlorite, or decomposed mica, instead of the usual mica." This is an erroneous definition, for if it contains either mica or ripidolite instead of talc, it cannot be Protogene.* However, Cotta would appear not to be the only geologist or petrologist who has found this rock a stumbling-block; as Haughton, after examining numerous specimens of the so-called Protogene from the Alps, states that most of them "contained not talc, but dull mica or ripidolite, or some kindred mineral." If a rock contains mica in addition to talc, or ripidolite in place of talc, it cannot be protogene, but is, in the one case, *Talcose Granite*, and in the other, *Chloritic* or *Ripidolitic Granite*.

* Some basic felstones, when metamorphosed, appear to be true protogene (quartz + felspar + talc).

The formation of granite is not fully understood; yet I cannot but believe that while some are undoubtedly intrusive, others have been formed by extreme metamorphic action. However, between the two extremes there is a rock (previously mentioned and described as *Intrusive oligoclase granite*) that partakes of the nature of both, having constituents very similar to the ordinary oligoclase granite, while it is intrusive, like the highly siliceous granite. This at first may appear an anomaly; nevertheless it seems capable of a simple explanation.

All authorities infer that granites were formed at a considerable depth beneath the earth's surface, under pressure, by extreme heat. They cooled and consolidated, more or less gradually, long previous to coming near, or being exposed at the surface of the earth.

Rocks, under the above conditions, would melt more or less rapidly in accordance with the intensity of the heat to which they were subjected, the amount of pressure, and constituents of which they are composed. Moreover, the more acid portions would remain longer fluid, during refrigeration, than the basic (*see page 28*).

If, while in a state of fusion, the heating action were to cease or begin to subside (either of which must at some time have happened), the rocks or strata affected would be in the following conditions: Part would be entirely fused or liquefied; part would be semifused; while the rocks adjacent to the latter would be only more or less altered. Of the latter class, the least altered rock would be represented by rocks belonging to the *Schist series*; while the more altered rocks, according to the intensity of the metamorphic action, would form the *Gneiss*

series and the *Gneissoid granite*. The second class, or the semifluid rocks, would form the *Oligoclase granite*, or the *Granite due to extreme metamorphism*; while the first class, or the completely fluid mass, would form the different intrusive or irruptive rocks.

Of the completely fluid mass, part ought to find its way to, or close to, the earth's surface, and then cool under little or no pressure, forming the *Plutonic* and *Volcanic rocks*; a portion would consolidate between the source, the caldron, if it may be so called, and the surface of the earth, under more or less pressure, and form *Elvanyte* or *Quartz-Porphry*, the granitic rocks next to be enumerated and described; while the residue would consolidate in or near the caldron.

The consolidation of the residue, that remained in or near the caldron or source, could scarcely have been simultaneous, as it is probable the outer or upper portion would consolidate sooner than the inner or lower part. Moreover the acid portion would consolidate slower than the basic part; consequently there ought to be two distinct rocks forming,—a more basic rock above, and a more acid or siliceous rock below. As the upper or more basic rock cooled and consolidated, it ought to shrink more or less, and thereby form cracks, fissures, and other vacancies, into which the underlying still fluid or semifluid matter would be squeezed; therefore at any time prior to their final consolidation, part of the *oligoclase granite* or part of the *highly siliceous granite* might be forced up into and occupy all vacancies in the supercrescent rocks, and thereby form *intrusive granite* of one or both types (*Intrusive oligoclase granite* and *Intrusive*

highly siliceous granite) ;—subsequently, when the overlying rocks were denuded away, tracts, veins, &c., of these granites would be exposed.

The above suggestions, if correct, give a solution for the formation of the three different kinds of granite,—Highly siliceous granite, Intrusive oligoclase granite, and the Non-intrusive oligoclase granite. The Non-intrusive oligoclase granite would be represented by the semifluid portion that first consolidated ; the Intrusive oligoclase granite by the more basic portion of the fluid mass that was forced out of its normal position ; while the Highly siliceous granite would be the part that took longest to cool and consolidate. That granite must cool and consolidate under laws somewhat similar to the above appears highly probable, because when two or more granites come from one source, the last formed is more siliceous than the rock through which it breaks up, while the veins of segregation in the former are more siliceous than the parent rock in which they appear.

D. Elvanyte (Jukes) ; **Quartz-Porphry** ; **Quartziferous Porphyry** ; *Granitic Porphyry* (Cotta), [Celtic, *el* or *oil*, rock, and *van* or *ban*, white, so called from the white crust of the rock when weathered]. —A granitic rock, having a granular or compact felsitic or feldspathic matrix, inclosing globules, blebs, and crystals of quartz, crystals of one or more feldspars, with more or less mica or amphibole, ripidolite, and the like. Pyrite and marcasite are often constituents. In many of the siliceous varieties, a considerable portion of the quartz crystallized out prior to the other minerals. The matrix may be siliceous (*felsitic*) or basic (*fel-*

spathic), and of a greenish, greyish, brownish, blackish, bluish, or even yellowish colour: all of the yellowish, as also some of the blues, seem due to weathering.

NOTE. — Elvanyte naturally includes not only the quartz-porphyrries, but also the syenytes (quartz+felspar+amphibole) of some British geologists.

The free silica or quartz in the Elvanyte is for the most part in irregular blebs or globules, from minute specks to the size of a pea, or even larger. The quartz internally has a glassy, often radiated structure, and externally on some, the crystal faces can be recognized, while others appear to be regular coated minute balls. The quartz is scattered through the mass, sometimes sparingly, but at other times so thickly as to give the rock a pisolitic aspect. Elvanytes, except some of the very granitic varieties, weather with the even flowing surface so characteristic of the Felstones, and not with the rough rugged aspect of granite. They often near the edge of large masses, at the walls of dykes, and in small branch dykes or veins merge into a rock in appearance identical with felstone: this compact rock is called by Cotta *the Base* or *Mother-rock*. The Mother-rock at dyke walls is rarely more than a few inches thick. Elvanytes being hypogene rocks, are never accompanied by *tuff*.

NOTE.—A true elvanyte is more crystalline at the centre of a dyke or mass than at the margins, while an elvanoid rock due to metamorphism is more crystalline at the margins than in the interior, the change in the latter taking place from the outside inwards.

Varieties in Composition.

A. FELSITIC and FELSPATHIC ELVANYTE.—Respectively a siliceous or a basic elvanyte, in which the quartz, mica, and other minerals are more or less developed.

a. Orthoclastic Elvanyte.—When the felspar is principally orthoclase.

b. Oligoclastic Elvanyte.—When the felspar crystals are principally oligoclase.

c. Rappakivi Elvanyte.—An elvanyte in which many of the orthoclase crystals (often pink or flesh-colour) are inclosed in an envelope of oligoclase (often light green). In the Finland variety the orthoclase is generally in rounded masses inclosed in orbicular envelopes of oligoclase.

B. QUARTZITIC or SILICEOUS ELVANYTE.—Having a large percentage of siliceous matter.

C. MICACEOUS ELVANYTE.—Having a large percentage of mica.

NOTE.—These rocks would be called “mica traps” by some authors. (See *Mica Trap*, page 48.)

D. PYRITOUS ELVANYTE.—Having a large percentage of either pyrite, marcasite, or chalcopyrite.

E. HORNBLENDIC ELVANYTE.—When typical, a granitoid rock, an aggregate of quartz, felspar, and amphibole, usually with some pyrite or marcasite; it however graduates into more or less compact varieties.

Varieties in composition are, — *a. Micaceous*; *b. Chloritic*; *c. Felspathic*; *d. Pyritous*; respec-

tively, as different minerals give a marked character; some structural varieties are gneissoid.

NOTE.—Some authors would call hornblendic elvanyte and its varieties "syenyte." Part of a mass may be a true syenyte (*amphibole*+*orthoclase*+*quartz*), but in general the rock contains two feldspars.

F. CHLORITIC OR RIPIDOLITIC ELVANYTE.—Having ripidolite replacing most or all of the mica or amphibole.

Structural Varieties.

G. GRANULAR ELVANYTE.—With a matrix somewhat resembling granite; but distinct crystals of one or more feldspars, with globules, blebs, and crystals of quartz, flakes of mica, or crystals of amphibole, are separately and prominently developed. In one variety ripidolite replaces the mica or amphibole.

H. COMPACT ELVANYTE, or PONTELLARYTE.—The matrix is more or less compact throughout, forming a paste, often dark-coloured. It contains separate crystals, globules, or flakes of one or more feldspars, amphibole, quartz, mica, or ripidolite.

I. GNEISSOID ELVANYTE.—Elvanyte with a structure that seems to be foliation, more or less well-defined. Some varieties are micaceous, others hornblendic; while some may be chloritic. Some gneissoid elvanytes may be metamorphosed igneous rock.

J. GRANITOID ELVANYTE.—For the most part an aggregate of quartz and feldspar, with mica, or ripidolite, or amphibole. Part of the matrix appears as a sort of paste, while some of the

quartz occurs in globules, blebs, or irregular crystals, having crystallized out prior to the other minerals.

Most *granitoid elvanytes* weather more like a felstone than like a granite; some, however, do not. At the walls of dykes or masses they often merge into a more or less compact rock—"The Mother-rock, or Base," of Cotta. Granitoid elvan-
yte seems to be the passage-rock between granite and elvanyte.

K. PYROMERIDE, or BALL ELVANYTE, *Ball Porphyry*.—

"This rock, in addition to the usual quartz crystals, contains balls of felsite (either small and numerous, or large and isolated. The small balls are frequently marked internally with radial streaks. The interiors of the larger ones are usually split after the manner of septaria, or they contain a geodic cavity). The clefts or cavities in the balls are wholly or partly filled with hornstone, chalcedony, agate, quartz, amethyst, calcite, fluorite, &c. &c."—Cotta.

Class II.—PLUTONIC ROCKS.

THIS name (from *Pluto*, the god of the infernal regions) has been given to the rocks that at one time were buried beneath the earth or sea, as distinct from the Volcanic, or those that are thrown up and consolidated on the present surface of the earth. The Plutonic rocks were irraptured or intruded, and consolidated at or near a former surface of the earth, or at the bottom of a sea or lake, under, comparatively speaking, little pressure; but subsequently they were covered by successive

deposits of "Derivate rocks." As they are not "hypogene rocks," they have their tuffs, agglomerates, and other mechanically-derived associates, with which they may be interstratified, as also with any class or variety of derivate or sedimentary rock.

In Nature there are no hard divisional lines between different kinds of rocks, more especially ingenite rocks; therefore the Plutonic rocks, on one hand merge into Granitic rocks, and on the other into Volcanic rocks. A boundary between the Granitic and Plutonic rocks can be defined, as the first are hypogene, while the latter are not; but the difference between the Plutonic and the Volcanic rocks is much more vague, we only knowing that the latter are intrusive rocks, erupted and consolidated at the *present* surface of the earth, while the former were formed in ages or geological periods *long past*, and subsequently were covered up by accumulations of strata that are now in part removed by the force of denudation, thereby bringing them again to the surface.

E. Felstone, Felsyte, or Felsite [Ger. *felspath*, rock spar].—A compact or granular, or splintery quartzitic felspathic rock—often porphyritic or quartzose; sometimes micaceous, or hornblendic, or ripidolitic; pyrite or marcasite, and sometimes chalcopyrite, may occur as constituents.

The Felstones, or Felsytes, include all the highly siliceous plutonic rocks, and are for the most part a felsitic compound. Various minerals, however, are locally ingredients, forming numerous subgroups and varieties. Felstones, or Felsytes, usually weather in flowing surfaces with even outlines. They are, however, affected by different structures. Some are *homogeneous*; in some there is a platy arrange-

ment, rudely parallel to the walls of the dykes ; while in others, sometimes, however, associated with the platy structure, is a spheroidal or concretionary arrangement. The platy structure may be due to the rock cooling in thin sections, or nearly vertical layers ; or perhaps, as suggested by Scrope, to the differential motion of the vesicular and non-vesicular portions of the cooling rock. The spheroidal or concretionary arrangement seems always to occur near the end of a dyke, and may be due to the cooling when the rock was in its last stage of motion : Phillips, in his description of "Vesuvius," figures and describes a similar structure in some of the dykes of lava belonging to that volcano.

In other felstones, or felsytes, there are lines cutting obliquely across ; while in some is a rudely columnar structure, except that the joint-planes are as often oblique as perpendicular to the walls of the dykes. Associated, in places, with the last-named is a very irregular platy development between the joint-lines usually perpendicular to them, but often more or less oblique.

The minerals most commonly appearing in Felstone or Felsyte, are—crystals of orthoclase, oligoclase, pyrite, marcasite, and amphibole, blebs or crystals of quartz, and flakes of black and white mica, and ripidolite. The quartz globules vary from minute particles to the size of a pea.

- A. PETROSILEX (Brongniart) [Lat. *petra*, stone, and *silex*, flint, on account of the hard matrix and flint-like appearance of the rock], *Felsyte*, or *Felsite*.—The type rock of the Felstones. A compact or granular, hard, siliceous felsitic rock ; a granular mixture of felspar and quartz, in which, although fine-grained, the minerals

constituents do not merge into one another, but are recognized by the eye, with or without the help of a magnifying-glass. Petrosilex is greyish, greenish, purplish, or bluish in colour; when yellowish or reddish, the colour seems due to weathering. Some are splintery, but usually they have from an uneven to a conchoidal fracture, and are translucent or sub-translucent. Some have a porcelanic, others a saccharoid aspect.

a. *Felstone glass*; *Pitchstone felsyte*.* — “The principal mass is homogeneous; of vitreous pitch-like appearance; conchoidal fracture; resinous lustre; translucent at the edges, and very variously coloured.”—*Cotta*.

b. *Ribbed petrosilex*; *Striped*, or *ribbed felsyte*, with bands of different colour, texture, and sometimes composition alternating. — In some varieties the rock splits into plates along the riban.

B. QUARTZITIC OR QUARTZOSE FELSTONE, or FELSYTE; QUARTZIFEROUS FELSTONE.—A compact felsitic mass, inclosing crystals or crystalline grains of quartz. Of quartzitic felstone there are different varieties; some due to composition, others to structure.

The compact matrix consists principally of felspar, and of it Cotta says: “Probably orthoclase; its proportion of silica is, however, too high even for orthoclase, and it is therefore probable that some quartz is intimately combined with the felspar.” The colour of the matrix is greyish, greenish, purplish, or bluish, with the surface

* *Pitchstone felsyte* must not be confounded with the volcanic rock.

weathering a dirty white; when the rock is yellowish or reddish, the colour seems due to secondary action. The texture of the matrix varies considerably: sometimes it is compact like horn, with a smooth conchoidal fracture; at other times it is granular, or saccharoid, or splintery. Usually it is compact, with a fracture from semiconchoidal to uneven, but sometimes it is vesicular or amygdaloidal.

Varieties in Composition.

- a. *Oligoclasic quartzitic felstone*; b. *Micaceous*;
 c. *Hornblendic*; d. *Chloritic*, or *Ripidolitic*;
 e. *Pyritous*, according to whichever mineral gives a character to the rock.

The micaceous varieties graduate into *Minette*, while other varieties graduate into *Felsitic elvanite*.

Structural Varieties.

- f. *Striped*, or *Ribbed quartzitic felstone*.—Thin layers of somewhat dissimilar texture; hence the fracture appears to be striped like a riban, and the rock splits more easily in the direction of those layers than across them.
- g. *Variolitic*, or *Spotted quartzitic felstone*, containing in the matrix worm-shaped spots, or blotches of different colour, texture, and usually of composition.—In the latter case the rock weathers into pock-marked, or ovate hollows.

C. **MINETTE, MICACEOUS FELSTONE, OR FELSYTE.**—A more or less felsitic or felspathic rock, containing much mica, and sometimes distinct crystals of orthoclase or oligoclase, or amphibole.

- a. *Fraïdronite*—*Fraïdronite*.—"A greenish felspathic principal mass combined with a greater

or less quantity of mica. Pyrite and quartz occur as accessories."—*Cotta*.

NOTE.—MICA TRAP.—If the grouping of some recent writers were followed, the sub-group just described, and all other highly micaceous rocks, should here be introduced and described as *Mica traps*. Under this name, however, have been included "micaceous elvanyte," "micaceous felstone or minette," "micaceous euryte," "micaceous diabase or kersantite," "micaceous dioryte or kersanton," "micaceous melaphyre," "micaceous doleryte," and many micaceous varieties of the above subgroups; rocks that are quite dissimilar, except that mica is largely developed in each. Mineralogically, such a group may exist, but geologically or petrologically it does not, for the different rocks which must be included under the name are varieties or subgroups not only of quite distinct groups, but also of distinct classes, for under it would be included granitic, plutonic, and volcanic rocks. Such a group will not be introduced into this Manual, but each micaceous rock will be found described in its natural group and place.

D. EURYTE, or EURITE [Gr. *eureo*, to flow well or easily].—A compact or granular felspathic rock, occasionally showing glistening specks of quartz in the mass, usually purplish or greenish in colour; may be vesicular or amygdaloidal; when heated before the blowpipe in thin pieces, it fuses with greater or less facility. It is often columnar.

In A.D. 1817 Daubuisson proposed the name *Eurite* for the basic felstones, on account of their fusibility. This name was afterwards, however, misapplied by Delesse and others, who called *Granityte* (veins of segregation in granite), by Daubuisson's name, apparently mistaking the rock, on account of Granityte being in colour and aspect somewhat similar to *Euryte*.*

* A *Granityte* to the eye may appear the same as a *Petro-silex* or a *Euryte*, but if carefully examined will always be found to

Subsequently Naumann has proposed for Daubuisson's Euryte the name "Porphyrite," and Cotta advocates its adoption. It however seems to be a most objectionable name, as many authors use the term synonymously with porphyry.* Moreover, the name refers to a structure which is not an essential feature of the subgroup, or, to quote Cotta,— "*The name of Porphyrite refers to a texture which is not an essential feature of these rocks, because the Porphyrites are not always in fact porphyritic.*"

NOTE.—Intermediate between the highly siliceous felstones, or *felsytes*, and the *whinstones*, are rocks that partake more or less of the nature of both. These intermediate rocks Jukes and other British geologists seem to class with the basic rocks under the general name of Greenstone; † a name derived from the prevailing colour of the rocks, as even the purplish varieties have a green tinge, or merge into a green. One reason for this classification seems to be, that all the rocks included, fuse with greater or less facility before the blowpipe. Within the last few years there appears to be a reaction among some of the British geologists, who seem inclined to go to the other extreme, and place not only the eurytes, but also all the micaceous dolerites, diorytes, and diabases in the Felstone group. This is evidently erroneous, as the latter rocks ought to be put with the basic rocks; or they and the eurytes should be placed in an intermediate group (called *Hybrid rocks* by Durocher) between the highly siliceous and the basic plutonic rock.

Varieties in Composition.

- a. *Orthoclastic euryte*; b. *Oligoclastic*; c. *Micaceous*; d. *Hornblendic*; e. *Chloritic*, or *Ripidolitic*; f. *Quartzose*.

contain minute flakes of mica and a granular structure, never having the compact or semivitreous appearance characteristic of a *Euryte*.

* Its advocate, Cotta, seems to use it in this sense.

† Jukes, in his Manual, mentions that there are "*intermediate rocks*," but he does not name them.

The orthoclase and oligoclase appear as crystals, changing the rock into a *Porphyry*; the micaceous variety by some would be called a *Mica-trap*, and in the quartzose variety the quartz appears in blebs, globules, and crystals.

Structural Varieties.

- g. Euryte porphyry*; *h. Amygdaloidal euryte*; *i. Vesicular*; *j. Compact*; *k. Slab or flaggy euryte*; and *l. Cleaved*; according to the structure that gives a character to the rock mass.

The compact varieties have a conchoidal or sub-conchoidal fracture, while some of the cleaved rocks make a coarse roofing-slate.

Structural Varieties of the Felstones.

E. COMPACT FELSTONE, or FELSYTE.—Typical felstone or felsyte (*see Petrosilex*).

F. GRANULAR FELSTONE, or FELSYTE. -

G. FRIABLE, or MEALY FELSTONE.—A more or less fissile, or friable felstone; usually of a pale greenish or greyish colour; tough, but splits easily; a mealy or scaly aspect; often has a soapy feel, showing a transition into *Steatyte*.

This rock is undistinguishable in aspect and composition from some of the fine mealy *Felstone tuff*. It may possibly be a true tuff, but as it occurs in dykes and intrusive masses, and often graduates into a compact felstone, it is here described.

H. RUBBLY or SHINGLY FELSTONE; THICKLY-JOINTED FELSTONE.—So traversed by three or more regular or irregular systems of joint-lines, that the rock breaks up into rubble or angular shingle, and is incapable of being quarried in large pieces.

I. CLEAVED or SLATY FELSTONE.—With cleavage-planes, sometimes so distinct that the rock splits into slates, but rarely into “Roofing slate.”

J. FOLIATED or SCHISTOID FELSTONE; SCHISTOSE FELSTONE.—A fissile felspathic rock, composed of leaves of felspar, with mica or amphibole, or ripidolite; sometimes with a little quartz.

NOTE.—Rocks very similar both in aspect and in composition are found among the metamorphic igneous rocks. Nevertheless it may be possible that some felstones are so constituted that on solidification their constituents arrange themselves in leaves. At Barna, co. Galway, Ireland, breaking up through the porphyritic oligoclase granite, are dykes belonging to this subgroup that apparently are newer than the time at which the general metamorphism of the associated rocks took place.

K. GRANULYTE, or GRANULITE.—“A fine-grained to compact fissile compound of felspar and quartz, usually with some mica.”—*Cotta*.

NOTE.—Granulyte, although allowed to remain provisionally among the Felstones, may properly belong to the metamorphic igneous rocks, and be a felstone into which a fissile structure has been introduced by metamorphism. Some of the metamorphic igneous rocks hereafter to be described seem to be identical with the granulite of *Cotta*.

L. FELSTONE or FELSYTE PORPHYRY; PORPHYRY; ORTHOPHYRE.—Felstone, or Felsyte, containing distinct and numerous crystals of one or more felspars.

- a. Amygdaloidal* (Gr. *amygdalon*, an almond);
- b. Vesicular.*

If the felsitic or felspathic matrix is full of cavities, the rock is called *vesicular*, while, if it contains crystalline secretions filling the original cavities, it is called *amygdaloidal*.

Felstones, or Felsytes, supposed to be metamorphosed.

Metamorphic action as yet has not been properly explained or understood. It cannot be denied that derivate rocks may be altered into schist and gneiss; yet many authorities seem to doubt that igneous rocks could be altered by the action that metamorphosed the others. That the latter have sometimes been altered, seems not only probable but also absolutely necessary, for the following reasons:—

There are derivate rocks of all ages, which have their associated igneous rocks. Such derivate rocks, except, however, those of the most recent age, are often metamorphosed; the metamorphism being clearly *normal* (to use Delesse's expression), *i.e.* not the result of the intrusion of the igneous rocks, although, as is often evident, effected after that intrusion. The agents which have prevailed to metamorphose the containing derivate rocks must unquestionably have attacked the contained igneous rocks, and that they have sometimes succeeded in altering them also more or less, seems clear from the circumstance, that the igneous inliers just mentioned are often different from any of those that occur in unmetamorphosed sedimentary rocks; which circumstance cannot be regarded as merely accidental.

In some localities, such as West Galway and Mayo, Ireland, there appear to be data for the above suggestions. In these places it has been observed that associated derivate and igneous rocks appear to have been acted on in similar ratio, as expressed in the following table: *

* This table only records changes that have been observed, *not, however*, all the changes rocks are capable of. The first *stages of metamorphism* of the Whiustones are so various, that *it is hard to give a general name for the rocks.*

UNALTERED ROCKS.	FIRST STAGE— METAMORPHISM.	SECOND STAGE— METAMORPHISM.	THIRD STAGE— METAMORPHISM.
Sandstone	Micacized sandstone...	Schist.....	Gneiss.
Quartzose sandstone ...	Quartzite	Mica-schist	Gneiss.
Shale	Argillite	{ Mica-schist	Gneiss.
Slate	Phyllitic schist	{ Chloritic schist	{
Slaty or shaly tuff	Phyllitic schist	{ Hornblende schist...	{ Hornblende gneiss.
Felsitic tuff	{ Felsitic schist	{ Talcose schist	{
Mealy or friable fel- stone	{ Steatyte schist	{ Steatyte.	{
Felstone	Ditto	{ Talc-schist.	{
	{ Schistoid felstone	Ditto.	Gneissyte.
	{ Granular felstone	Gneissoid felstone,	Granite, or granitic fel- stone.
Euryte	Schistoid euryte	Micacized granular fel- stone	{ Syenite.
Whinstone		Gneissoid euryte	{ Hornblende granite.
		{ Dioryte	{
		{ Syenite	{ Hornblende granite.

M. GNEISSYTE OR GNEISSITE; GNEISSOID FELSTONE.—A crystalline fissile aggregate of quartz, felspar, and mica, more or less foliated; always occurring in dykes or intrusive masses, associated with gneiss or schist.

NOTE.—Apparently this rock is the same as the rock described by Cotta, and called *Red Gneiss*.

Gneissyte generally does not occur among sedimentary rocks in the first stage of metamorphism* (*Schist series*), but among highly metamorphosed sedimentary rocks (*Gneiss series*); dykes and masses of it are often found.

Varieties in Composition.

a. Quartzose Gneissyte; b. Felsitic; c. Micaceous; d. Hornblendic; e. Syenyte Gneissyte; f. Chloritic or Ripidolitic; g. Talcose Gneissyte.

Varieties *d*, *e*, *f*, and *g* are probably metamorphosed eurytes. The difference between *d* and *e* is, that the former contains mica, in addition to amphibole, while in the latter there is only amphibole.

Structural Varieties.

h. Porphyritic, and i. Schistoid.

The latter variety splits easily along the planes of foliation, on account of one of the constituents, usually the mica or the quartz, being only very sparingly developed. It is the passage rock between gneissyte and schistoid felstone.

N. SCHISTOID FELSTONE OR FELSYTE.—A fissile aggregate of felspar and mica; generally a little

* *For the degrees of metamorphism of the derivate rocks see pages 37 and 76.*

quartz is present, but in such small quantities as to seem more an accessory than an essential. The rock mass is more or less foliated.

Some schistoid felstones have an unctuous feel, and change by pseudomorphic action into *steatyte schist*, or even *steatyte*.

NOTE.—The foliation in the metamorphic felstones seems to follow conspicuous structures in the original rocks. Some normal felstones are amorphous, others have a spheroidal or platy structure, while some are cleaved. In the amorphous felstones there are often lines apparently due to viscid fusion, while in the platy felstones the lines are parallel, or nearly so, and at other times oblique to the walls of the dykes. In the metamorphic felstones very similar structural peculiarities can be found; in some there is little or no foliation, but the rocks are banded (see *Granitic Felstone*, page 35), while in some there is well-marked foliation, either parallel or oblique to the walls of the dykes, or spheroidal.

Schistoid felstones may be named after conspicuous minerals locally essential; they may be *quartzose*, or *felspathic*, or *micaceous*, or *hornblendic*, or *steatitic*, &c. &c.

F. Whinstone; * *Greenstone*.—A crystalline aggregate of feldspar with either pyroxene, amphibole, or urolite. With these minerals, in forming subgroups and varieties, are associated ripidolite, mica, pyrite, marcasite. Under the group of Whinstone are

* The term Whinstone has been adopted in place of Greenstone, as the latter has different significations, according to the authority by which it is used. Jukes and many other British geologists include under that name not only the basic plutonic rocks, but also all the basic felstone, especially when of a green colour. Cotta includes under this name Dioryte and Diabase. Brongniart confines the term to Dioryte, while Naumann restricts it to Diabase. Whinstone seems to be a nearly universal name in England, Ireland, and Scotland, the only objection to it being that in places it is also applied to the basic volcanic rocks

included all the very basic Plutonic rocks. They are usually dark-coloured, greenish or blackish; however, a few are light-coloured. The whinstones are often vesicular or amygdaloidal, and have their associated tuffs, with which they are often interstratified.

A. DOLERYTE [Gr. *doleros*, deceptive].—A tough aggregate, principally a mixture of felspar and pyroxene. Nearly always of a dark colour.

a. *White-rock* (Jukes).—A doleryte of a nearly white colour, found associated with coal in Staffordshire and elsewhere. The dykes of "white-rock" can be traced into masses of ordinary dark-coloured doleryte.

b. *Aphanyte-doleryte*; *Basalt*.—Very compact; often called "Basalt;" the latter name, however, ought to be only used for a variety of volcanic rocks.

B. MELAPHYRE.—A crystalline aggregate of felspar and pyroxene, with or without amphibole; of a dark colour, greenish, brownish, or blackish, with more or less resinous lustre; may be compact, porphyritic, vesicular, or amygdaloidal, and often contains specks of opal or quartz, also zeolites. Before the blowpipe it fuses readily; fracture uneven.

Varieties in Composition.

a. *Eukryte*, or *Eukrite*.—A crystalline granular aggregate of anorthite and pyroxene, occasionally with some olivine, amphibole, and epidote; the latter mineral being due to decomposition.

Peridolyte.—A subvariety in which olivine is *always* present.

- b. Magnetic Melaphyre*; having magnetite as a constituent. This rock may always be distinguished from *magnetic diabase*, as the magnetic constituent of the latter is pyrrhotite.
- c. Micaceous Melaphyre*.—The mica often is in considerable quantities.

Structural Varieties.

- d. Porphyritic*, containing felspar crystals; *e. Amygduloidal*; *f. Vesicular*; and *g. Variolitic*, having spots of a different colour and texture, also perhaps of composition.
- C. PYROXENYTE (Dana).**—A compound of pyroxene and orthoclase.
- D. DIABASE.**—A crystalline granular aggregate of felspar, pyroxene, and ripidolite; colour usually greenish or purplish. The felspar is either oligoclase or labradorite, and frequently the latter. The pyroxene is diallage, and rarely ferruginous. The ripidolite is generally green, and in subordinate quantities; it is often only distinguishable with difficulty, from being in intimate admixture with the diallage.

Varieties in Composition.

- a. Pyroxenic, or Augitic Diabase*, having numerous and well-developed crystals of pyroxene, which, when weathered, give the rock a spotted appearance like a toad.
- b. Uralitic Diabase*, with numerous and well-developed crystals of uralite. In some subvarieties part of the pyroxene in the matrix is replaced by uralite. This rock, also, when weathered, has a spotted aspect.

- c. Labradoritic*; *d. Oligoclastic*; *e. Magnetic*, containing pyrrhotite; a micaceous subvariety, weathers freely into a magnetic sand; *f. Kersantyte* (Delesse), or *Micaceous*; and *g. Calcareous*.—"In a fine-grained or compact matrix of diabase rock are found small rounded grains of calcite, which do not appear to be the fillings-up of cavities."—*Cotta*.

Structural Varieties.

If diabase contains numerous and well-developed crystals of felspar, it is called *h. Porphyritic*; if there are roundish concretions of any mineral, it is called *i. Amygduloidal*; while it is *j. Vesicular*, if there are open cavities; *k. Variolitic* ("In the principal mass round concretions occur of a compact or radial-fibrous, or concentric felsite (labradorite)" —*Cotta*); and *l. Aphanyte*, or *Compact Diabase*.

E. GABBRO; *GRANITONE*; *DIALLAGE ROCK*—"consists of labradorite or saussurite, and diallage, or smaragdite irregularly combined; also sometimes all of these minerals together. It is very coarse-grained, fine-grained to compact, sometimes slaty or spotted."—*Cotta*.

Some Diallage rock, such as that of the Lizard, Cornwall, is a metamorphosed igneous rock.

- a. Euphotide*.—"A combination of felspar (saussurite) and diallage, with titaniferous and chromic magnetic iron-ore, pyrite, serpentine, and carbonates. The diallage often occurs as the variety smaragdite."—*Cotta*.
- b. Noryte*, or *Norite* (Scheerer), *Labrador Rock*.—"The Noryte of Scheerer, not of Esmark, is a compound of hypersthene or diallage,

labradorite, orthoclase (containing soda), and even some quartz."—*Cotta*.

Sterry Hunt has described rocks under the name *Anorthosyte*. Of these, some contain a felspar between andesine and anorthite, some between andesine and labradorite; while in others it is a pure labradorite.

Many of these rocks (*a* and *b*), if not all of them, may more properly belong to the metamorphic igneous rocks.

c. Hyperyte, or Hyperite; Hypersthenyte.

The hyperyte of Cotta and Dana is a coarse-grained to compact aggregate of labradorite and hypersthene.

Some varieties are said to occur as unaltered igneous rocks, but many of the coarse crystalline varieties would appear to belong to the metamorphic igneous rock series.

F. DIORYTE, or DIORITE [Gr. *dioros*, a clear distinction, in contradistinction to the doleryte, in which the minerals are indistinct].—A crystalline granular compound of felspar (not orthoclase) and amphibole. Sterry Hunt has called this rock *Anorthyte*.

Many diorytes are undoubtedly metamorphosed igneous rocks. In them the minerals are always prominently and distinctly visible; therefore it does not appear at all improbable that they were the rocks which first received this name. According to Dana, the felspar ought to be *albite*. He however seems to be nearly alone in this opinion.

a. Amphibolyte.—A homogeneous fine-grained compact mass, consisting of labradorite and

amphibole (*Dana*). According to Naumann, the latter ought greatly to predominate.

- b. *Aphanyte* [Gr. *aphanes*, unmanifest], or *Anamesyte* (Jukes).—A compact homogeneous variety of dioryte, without distinct grains.

Some dioryte-aphanytes are said to have as their felspar, orthoclase; in which case they are not, properly speaking, a homogeneous variety of *Dioryte*, but a homogeneous variety of *Syenite*.

- c. *Kersanton* (Rivière), *Micaceous Dioryte*.—Containing a great and conspicuous quantity of mica.

NOTE.—The highly micaceous diorytes, similarly as the highly micaceous melaphyres and diabases, would be classed by some geologists among the “mica traps.”

- d. *Napoleonite*, or *Orbicular Dioryte*.—A local variety, consisting, according to Delesse, of a combination of anorthite, blackish-green amphibole, with some quartz. The anorthite and amphibole form orbicular concentric layers round a kernel, each mineral forming different layers of alternate light and dark colour.

- G. **SYENITE, SYENITE**.—A crystalline aggregate of orthoclase and amphibole, with or without quartz. The quartz appears to be more an accessory than an essential.

NOTE. — Some syenites are undoubtedly metamorphosed igneous rocks, while others do not appear to have had such an origin.

The name Syenite has been given to various rocks. Haughton, Forbes, Cotta, Werner, and others describe it as a rock in which quartz is not a necessary ingredient; while Jukes, Dana, and

others define it as an aggregate of orthoclase, amphibole, and quartz. Some authorities erroneously class the hornblende granites among the syenites. Cotta says : "The orthoclase or microcline is usually the principal ingredient, and being in general red, it gives that colour to the whole rock. There are, however, syenites whose orthoclase is nearly white." In the syenites of West Galway, Ireland, of metamorphic origin, the orthoclase is usually white.

Varieties of Syenite are,—*a. Quartzose*, and *b. Titanitic*.

Whinstones, or Greenstones, supposed to be metamorphosed.

H. HORNBLENDE ROCK (Macculloch). — An aggregate of amphibole, felspar (usually not orthoclase), pyrite or marcasite, epidote, &c. &c. Some are so finely crystalline as to be nearly compact, while others contain crystals of amphibole, from one to four or five inches long.

Hornblende rock may in places be schistose, or nodular, or concretionary. Some of the schistose portions of this kind of rock, from Iarconnaught, Ireland, have been proved by Forbes to be derivate rocks; consequently such portions must be metamorphosed tuff. When Hornblende rock is nodular or concretionary, it may have spheroids from the size of a man's head to four or five feet in diameter, irregularly heaped up together, with the interstices filled with a schistose-looking stuff, that has a foliation rudely curling round the nodules; or the interstices may be occupied by a felsitic rock, or even with quartzitic stuff, or perhaps with two or more of these substances mixed together.

Spheroidal or concretionary structure occurs in some whinstones, and Scrope calls particular attention to it, in some of the augytes of Central France. To such a structure in the original (unmetamorphosed) rock, some of that in the Hornblende rock may possibly be due; nevertheless, the greater portion is probably due to the broken-up parts of the flow [the *Friction-breccia* of Cotta] that were rounded by attrition against one another; while the interstices were filled with the abraded or disintegrated portions of the rock, or by foreign matter washed into them by the water into which the flow was poured. This suggestion appears probable, more especially when the typical rock merges into the nodular variety, and the latter, through agglomerate or conglomerate, into schist and gneiss.

a. Actinolite Rock, or Glassy Actinolite Rock
(Krantz).

b. Tremolite Rock.

Hyaline acicular crystals of actinolite or tremolite, respectively mixed with a felspar (usually not orthoclase), or numerous nests of crystals may occur in a hornblendic felspathic matrix.

NOTE.—Actinolite and tremolite rocks are liable to merge into ophyte, or one of the allied rocks, as all the magnesian amphiboles seem to have a tendency to change by pseudomorphic action, partly or wholly into ophite, steatite, or such minerals.

c. Hyperyte, or Hypersthenyte (see *ante*, page 59).

d. Dioryte (see *ante*, page 59).

e. Syenyte (see *ante*, page 60).

é. Falso-syenyte, or Felsitic Syenyte, a variety of syenyte, in which the felspar (orthoclase)

predominates to the nearly total exclusion of the amphibole and other minerals.

f. Mico-hornblende Rock, or Micaceous Hornblende Rock.

Black, bronze, or white mica occurs as an essential. The flakes are sometimes of a considerable size, but more often they occur in small pockets or secretions. Usually they are regularly, although not very abundantly, distributed through the mass.

A mico-hornblende rock is easily distinguished, as the mica decomposes more readily than the other constituents, and gives the rocks a pitted aspect, if the mica is in bunches or pockets; while, if it is in large flakes, the weathered surface of the rock has often a somewhat graphic character, caused by the weathering away of the edges of the mica. Other varieties also have mica as an essential, scattered through the mass; these latter are often more or less schistoid.

G. Rocks due to Pseudomorphic Action, or to Weathering.—Some of these rocks, as will be evident from their description, belong in part to Order No. II. (*Derivate Rocks*); as, however, they are products of the igneous rocks, it seems expedient to describe them altogether.

A. OPHYTE, or OPHITE; SERPENTINE.—A compact rock, dull in fracture, with an unctuous feel. Colour dark green to blackish or brownish, reddish or variegated.

Dana divides massive serpentine into *Precious* and *Common Ophyte*. The first "is of a rich oil-green colour, of pale or dark shades, and translucent even when in thick pieces." The common ophyte is of dark shades of colour, and sub-translucent.

"The former has a hardness of from 2·5 to 3; and the latter often of 4 or beyond, owing to impurities."

Ophyte may originally have been either whinstone or felstone. It weathers into an impure ferruginous *meerschau*m of a dirty yellowish colour, giving the rock a peculiar burnt or baked aspect.

A variety is *Ophi-hornblende rock*, or a hornblende rock in which part of the felspar or part of the amphibole is changed into ophyte.

B. STEATYTE, Soap-stone.—An unctuous, soft, fissile, but, at the same time, tough rock, of a light greenish, greyish, or bluish colour.

Ophyte and *steatyte* often occur together in intrusive masses or dykes, associated with steatitic felstone or ophi-hornblende rock, also with hornblende schist or felsitic schist. Friable felstone, felstone tuff, whinstone tuff, &c., may change into steatyte.

Varieties are,—*a. Felspathic*; *b. Pyroxenic*; and *c. Amphibolic*.

Steatyte, as formed from felstone or whinstone, may contain a large percentage of felspathic tuff, or of pyroxene or amphibole.

C. EKLOGYTE, or EKLOGITE.—An aggregate of green smaragdite and red garnets. The smaragdite forms a finely crystalline matrix, in which the crystals of garnet are disseminated. It is often micaceous. Some varieties undoubtedly are pseudomorphs of hornblende rock.

NOTE.—Some of the fissile rocks, usually called Eklogyte, are metamorphic sedimentary rocks, and their description is given under the head of *Smaragdite Schist* (see page 93). The rock here described is an intrusive rock.

- D. EPIDOSYTE, EPIDOTE ROCK.** — An aggregate of felspar, amphibole, epidote, pyrite, or marcasite, &c. It occurs as dykes and intrusive masses, or as nodules and lentils in the metamorphic rocks.
- E. KAOLIN, Porcelain Clay** [Chinese, *Kau-ling*, high ridge, the name of the place where the clay was first manufactured].—A white, greyish-white, yellowish, sometimes brownish, bluish, or reddish, clay-like or mealy or compact substance, due to the decomposition of a very felsitic rock.

Baron Von Richthofen has proved that kaolin was originally manufactured by the Chinese from *Petrosilex*; but though the name at first indicated an artificial felsitic clay, now the term has generally been adopted for a clay due to the decomposition of a felsitic rock. *Kaolin* sometimes occurs in dykes, but more often it seems to have a bedded structure, when it more properly belongs to the second order of rocks (*Derivate rocks*).

- F. FULLER'S EARTH.** — "A substance resembling clay, somewhat greasy, but not in the smallest degree plastic, but falling to pieces in water; usually of a yellowish-green colour; is probably a product of the decomposition of basic igneous rocks."—*Cotta*.

Fuller's Earth sometimes occurs in dykes, at other times, similarly to kaolin, it seems to belong to the *Derivate rocks*.

- G. MEERSCHAUM, SEPIOLYTE.** — Compact, with a smooth feel, and fine earthy texture. Colour greyish-white, or with a faint yellowish or

reddish tinge; opaque. An earthy hydrated silicate of magnesia.

NOTE.—*Ophyte* weathers into an impure meerschaum. Dana points out that sepiolyte, in Asia Minor, occurs “in masses in stratified earthy or alluvial deposits;” and Cotta states it “forms separate beds, which are the result of a process of transmutation, probably of magnesite.”

H. MAGNESYTE.—An aggregate of the carbonates of magnesia and iron. It may contain felspar, mica, quartz, chrome, nickel, &c.

H. Tuff. Ingenite in part, derivate in part [Ital. *tuffo*, Gr. *tophos*].—The mechanical accompaniments of the Plutonic rocks, consisting of the dust, powder, cinders, fragments, blocks, and other rock *débris* ejected during an igneous eruption. Subsequently these were more or less stratified by water or air, and afterwards consolidated by pressure, heat, or cement; or, perhaps, by two or more combined.

NOTE.—The name tuff was originally applied to the mechanical accompaniments ejected during an igneous eruption, but afterwards it was erroneously used for any porous vesicular stone, often a purely sedimentary rock, while the term ash was used to denote the true tuffs. Ash, however, is an objectionable name, as an ash is the residue of any substance left after that substance has been burnt; while tuffs are never a pure ash, although portions of ash may form minor constituents. *Tuff* may be used as the name for the mechanical accompaniments of the Plutonic rocks, and *tufa* for those of the Volcanic rocks (*see* page 75).

Correctly speaking, no rock should be called a *tuff* unless the materials forming it were ejected in fragments from an igneous vent, and subsequently fell either on land or in water. In practice, however, this distinction cannot always be followed, many *tuffose* sandstones or *tuffose* shales, whose

origin is mainly due to the abrasion and disintegration of ingenite rocks, are undistinguishable in aspect or composition from true *tuff*. Forbes has pointed out, that when outbursts of ingenite rock are forced or have flowed into the sea, they may be "at once broken up into a state of division, more or less fine, in proportion to the greater or lesser cooling power of the water mass in immediate contact, and may be spread out into beds by the action of the waves." Many rocks thus formed, although not true tuffs, are yet not only undistinguishable from them, but in some cases (except under the microscope), are also undistinguishable in aspect and composition from igneous rocks. Tuffs may be stratified or unstratified; sometimes they even occur in dykes and pipes. As they, in part, are sedimentary rocks, fossils may occur in them.

Varieties due to Composition.

A. FELSTONE TUFF, FELSYTE or FELSITIC TUFF.—

An aggregate of felspathic or felsitic parts usually more or less flaky or mealy; colour reddish, greenish, greyish, or bluish, weathering yellowish-white. Often very greasy to the touch, which appears to be due to particles or portions changing into *steatite*.

As previously mentioned, portions of some felstones are very flaky or mealy in aspect; these it is difficult, except when studied *in situ*, to distinguish from felsyte tuff.

Varieties are, — *a. Quartzose*; *b. Calcareous*; *c. Pyritous*; *d. Cuprififerous*; and *e. Hematitic*.

B. WHINSTONE TUFF, Basic or Greenstone Tuff.—

A greenish, reddish, purplish, or greyish-bluish

rock, from coarse to fine and compact; often laminated, sometimes finely, at other times having alternate layers of fine and coarse materials; they may be conglomeritic or brecciated, when they often merge into *Agglomerate*.

Varieties are, — *a. Hornblendic*; *b. Pyroxenic*; *c. Chloritic*; *d. Calcareous*; *e. Pyritous*; *f. Cupriferous*; and *g. Hematitic*.

C. CALCAREOUS TUFF.—Tuff cemented together by a limy or calcareous matrix.

D. ARENACEOUS TUFF. E. ARGILLOUS TUFF.—These respectively contain a large percentage of sand and clay. Many of these rocks probably are not true tuff, but are due to the disintegration of igneous rocks, the detritus being mixed with sand or mud. Some arenaceous tuffs are cemented together by a siliceous paste.

F. HEMATITIC or FERRUGINOUS TUFF.—A reddish or yellowish ferruginous rock; sometimes argillous, at other times arenaceous. It may merge into an earthy limonite.

NOTE.—Some varieties of the Indian rock called *Lateryte*, from Kutch, seem to be a variety of hematitic tuff. Very similar rocks are found associated with doleryte in Antrim, Ireland: both in Ireland and Kutch the rock may be steatitic.

G. PYRITOUS TUFF. H. CUPRIFEROUS TUFF.—Containing a large percentage of pyrite, marcasite, or chalcopryrite, so as to give a character to the rock.

I. STEATITIC TUFF.—A greenish, greyish, bluish, or reddish mealy or flaky rock; having a greasy feel. It appears to be a pseudomorph.

*Varieties due to Structure.***J. PORPHYRITIC TUFF.**

K. NODULAR, with its varieties, *a. Concretionary*, and *b. Spheroidal*.

L. SHALY TUFF.—Fine-grained, compact, and deposited in thin laminae or layers. There are two varieties,—*a. Flaggy*, when the layers are between half an inch and four inches thick, the rock easily splitting into tabular masses; and *b. Slaty*, when there is a well-developed slaty cleavage.

M. CONGLOMERITIC TUFF, PLUTONIC CONGLOMERATE.
—Tuff containing blocks and fragments of different kinds of rock.

Its varieties are,—*a. Brecciated Tuff*, or *Plutonic Breccia*, and *b. Plutonic Agglomerate*.

Variety *a* includes those rocks that contain angular blocks and fragments, while Lyell has introduced the term *Agglomerate* to designate those large irregular unstratified accumulations of blocks and detritus thrown up by igneous eruptions.

Class III.—VOLCANIC ROCKS.

THE name of the class is derived from *Vulcanus*, the god of fire. They are eruptive or intrusive rocks, brought up by vulcanicity, and consolidated near or at the present surface of the globe. They are necessarily accompanied by tufa and such mechanically derived associates.

NOTE.—None of the rocks allied to the volcanic have been placed among the granitic rocks; nevertheless, it appears highly

probable that some varieties of *Liparyte* (Trachyte porphyry) ought to be classed among the *Elvanytes*. Cotta says of the liparytes, which are granitoid in aspect, that in some cases it is impossible to distinguish between them and *Quartz-porphry* (Elvanyte). "In these cases the only real difference consists in their geological connection with genuine trachytes or their petrographical transition into perlyte or pumice-stone."

I. Trachytic group.—Compact or granular, or splintery or cellular, crystalline felsitic or felspathic rock: some contain crystals of quartz, mica, and amphibole, or even pyroxene; the whole either confusedly united or imbedded in a felspathic paste. Some are porphyritic or even granitoid.

The Trachyte group embraces all the siliceo-felspathic volcanic rocks. Among them are also included such transitional basic felspathic volcanic rocks as the *Greystones* of Scrope, the *Trachydolerites* of Abich, or the *Hybrid rocks* of Durocher.

A. RHYOLYTE, or RHYOLITE.—"A compact enamel-like, or vitreous matrix enclosing grains or crystals of sanidine" (*rhyacolite*), "oligoclase, mica, or even quartz."—*Cotta*.

In Rhyolyte free quartz appears much more frequently than in common Trachyte; while it contains no amphibole or pyroxene, or at least, those minerals are rarely found in it. From this it is apparent that Rhyolyte is highly siliceous or felsitic Trachyte.

a. Liparyte, or Liparite; Trachyte-porphry.—"Is the name given to those rocks (prevalently felsitic and porphyritic with a compact matrix) which are geologically allied to the trachytes."—*Cotta*.

Liparyte has a compact felspathic matrix, containing crystals of felspar, and sometimes also

mica or quartz. As a rule it is much richer in silica than the trachytes hereafter described. Cotta says it only very rarely and exceptionally contains some traces of amphibole. In some the matrix is compact and somewhat shining; in others it is enamel-like, while in others it is dull. They may be platy or vesicular or pumiceous. Usually light-coloured; some are granitoid and undistinguishable except in their geological position and age from Elvanyte; the latter variety ought probably to be classed among the *Granitic rocks*. As among the Plutonic rocks, so also among the Volcanic rocks, there must be passage-rocks into true granite; among the Volcanic, however, these cannot be so prevalent as among the Plutonic, as the latter rocks have been usually exposed by the force of denudation, while rarely do we find the roots of the more recent eruptions exposed.

b. *Perlyte*, or *Perlite*, *Pearlstone*, *Pearlstone-porphry*.—"An enamel-like matrix containing round grains, several of which are constructed with concentric layers."—*Cotta*.

Perlyte may be granular, or sphærolitic (with compact or radial striped felsyte balls), or porphyritic, or pumiceous, or vitreous (with resinous lustre), or argillaceous.

c. *Obsidian* (so named after its discoverer, the Roman named *Obsidianus*).—Trachytic or volcanic glass.

Obsidian may be compact, or a mere glass, or porphyritic, with sanidine crystals, and sometimes also mica plates, or sphærolitic, in which case it is a passage-rock into perlyte.

d. *Pumice*, *Vesicular Obsidian* (Ital. *pomice*, akin

to *spuma*, froth).—Vesicular volcanic glass. Some varieties are so porous that they will float on water.

- e. *Phonolyte* or *Phonolite Clinkstone* (Gr. *phonè*, sound, and *lithos*, stone).—Compact, in places vitreous; due to cleavage-surfaces of felspar; usually dark greenish-grey. Often platy in texture, and rings when struck with a hammer. Weathers with a whitish crust, similarly to many of the *felstones*.

Phonolyte may be compact, vesicular, flaky, variolotic, porphyritic, or amygdaloidal.

- B. *TRACHYTE* (Gr. *trachys*, rough).—"A compound of sanidine, oligoclase (or even albite and labradorite), with some amphibole or pyroxene, and dark-coloured mica. A rough principal mass in which, as matrix, some of its mineral constituents are frequently distinctly and separately developed and imbedded."—*Cotta*.

Varieties in Composition.

- a. *Sanidine Trachyte*.—An aggregate of sanidine crystals, with some amphibole or mica. From coarse to fine, and from porphyritic to compact.
- b. *Drachenfels Trachyte*.—Containing oligoclase in addition to the sanidine, with some magnesia-mica, and amphibole, also pyroxene, magnetite, and titanite.
- c. *Domyte*, or *Domite* (after *Puy de Dôme*, Auvergne).—Oligoclase trachyte, having no sanidine. It also contains some amphibole or pyroxene, and dark-coloured mica.
- d. *Andesyte*, or *Andesite* (after the Andes).—Fine

or compact trachyte, sometimes vitreous; dark colour, with imbedded crystals.

e. *Trachydoleryte*, or *Trachydolerite*; *Greystone*.—

“A compound of oligoclase (or labradorite) with amphibole or pyroxene, some magnetite, and frequently also mica. These minerals lie imbedded in a grey or brown matrix.”—*Cotta*.

Trachydoleryte is the passage-rock between the Trachytes and Augytes.

Varieties in Texture.

f. *Granular Trachyte*; g. *Compact*; h. *Porphyry*; i. *Trachyte-lava* or *Vesicular Trachyte*; and j. *Alum-stone*.

The last does not appear to be a normal rock, but to be due to decomposition.

J. Augitic Group.—Aggregates of felspar with pyroxene and amphibole; they frequently contain mica and magnetite, while quartz is rarely present.

The Augitic group includes all the basic volcanic rocks; they may be compact, crystalline, granular, porphyritic, amygdaloidal, vesicular, or variolitic.

A. AUGYTE or AUGITE, DOLERYTE or DOLERITE, BASALT.—A crystalline granular aggregate of labradorite or nepheline and pyroxene, with some titaniferous magnetite; usually blackish or dark-coloured. In the compact mass there often occur prominently distinct grains or even crystals of olivine, labradorite, pyroxene, and magnetite.

NOTE.—The name *Basalt* is given above, as it is in such common use; it ought, however, to be solely confined to the compact varieties. See Doleryte among the Plutonic rocks.

Varieties in Composition.

- a. *Nephelite Augyte*, when this mineral is the felspar.
- b. *Hauynophyre* (Rammelsberg); hauyne being in the place of labradorite.
- c. *Allogovyte*, or *Allogovite* (Winkler), a reddish variety of labradoritic augyte; and
- d. *Common Augyte*, *Labradoritic Augyte*.

Varieties in Texture.

- e. *Anamesyte*, or *Anamesite* (Leonhard).
- f. *Basalt*, or *Compact Augyte*.

e and f seem to be different names for the same kind of rock, as both refer to those augytes that are so fine-grained and compact that the constituents are undistinguishable.

- g. *Porphyritic*, containing felspar crystals.
- h. *Amygdaloidal*, having almond-shaped concretions.
- i. *Basalt Lava*, or *Vesicular Augyte*, being scoriaceous or full of minute holes; and
- j. *Variolitic*, when there are dark grains in a lighter-coloured mass.

h and i always merge one into the other, as the latter was the original condition of all amygdaloids.

B. LEUCITYTE, or LEUCITITE, LEUCITE-ROCK.—A more or less distinct aggregate of leucite and pyroxene, with some magnetite.

Varieties are,—a. *Compact*; b. *Leucitophyre*, or *Porphyritic*, having felspar crystals; c. *Amygdaloidal*; and d. *Leucityte Lava*, or *Scorious* or *Vesicular Leucityte*.

K. Volcanic Tufa and Peperino (Ingenite in part, Derivate in part).—According to Scrope, the Italian geologists restrict the term *tufa* to the felspathic or trachyte aggregates, which are grey or whitish ; while the name *peperino* denotes the augitic varieties, which usually are brownish.

A. Tufa.—A trachytic aggregate of slag, ash, pieces of pumice and lava, with fragments of various other rocks. It may be arenaceous, argillaceous, conglomeritic, or brecciated. The principal fragments and particles are of trachyte.

Varieties are,—*a. Trachyte Tufa* ; *b. Pumiceous Tufa* or *Pumiceous Sand*, which have received the following local names : *Trass* (Rhine), *Tosca* (Sicily), and *Pausilippo Tufa* (Teneriffe) ; *c. Phonolyte* ; and *d. Pozzuolana*, a volcanic sand, very useful in the construction of mortar for hydraulic works.

B. PEPERINO.—A light porous rock ; augitic sand, scoria, cinders, &c., cemented together, the grains having a peppercorn-like appearance.

Varieties are,—*a. Augyte Peperino* ; *b. Leucityte* ; and *c. Palagonyte*, called after Palagonia, in Sicily.

C. VOLCANIC CONGLOMERATE, with its varieties, *a. Volcanic Breccia*, and *b. Volcanic Agglomerate*. Some of the mechanical accompaniments of the volcanic rocks consist of more or less stratified accumulations of blocks and fragments of volcanic and other rocks, usually more or less loose, but sometimes cemented together. If the contained blocks are round or roundish, the rock may be called *Volcanic*

Conglomerate; if they are more or less angular, a *Volcanic Breccia*; while if the accumulation is massive and without stratification, Lyell's name of *Volcanic Agglomerate* may be adopted.

Class IV.—TRANSITION OR METAMORPHIC SEDIMENTARY ROCKS.

THESE rocks, similarly to the metamorphosed igneous rocks, may occur in various degrees of change, from a rock scarcely altered to a rock undistinguishable from granite, in accordance with the intensity of the metamorphic action to which they were subjected.

The lowest degree of metamorphism seems to be, principally, induration, with the planes of the most conspicuous structure (whether lamination, cleavage, or jointing) glazed or micacized, while, at the same time, peculiar structures are developed. In finely laminated, or cleaved rocks, a crumpling takes place; in others a nodular or concretionary development; in some the joint-lines and the rock in their immediate vicinity, are silicified or hardened, so that on the weathered surfaces of the rocks are formed well-marked rectangular, rhombic, or oblique depressions. In the second degree, the rocks become typical schist; in the third gneiss; while in the fourth they become granitoid, and from that pass into typical granite.

Rocks usually pass in this order to the highest degree of change; however there are exceptions, as some rocks are more susceptible of change than *others*, on account of their mineral constituents.

Thus some sandstones, or even shales that contain the constituents of gneiss, may have a gneissoid aspect, while the associated rocks have only been changed into schist.

NOTE.—The following changes in sedimentary rocks were observed in co. Mayo, Ireland:—*Felspathic* and *micaceous sandstones*, changing into mica-schist and gneiss. This gneiss, however, was not typical; that is, the quartz, mica, and felspar were not arranged in leaves; nevertheless the rocks contained these three ingredients, and had an incipient foliation;—*quartzose sandstones* and *grits*, changing through quartz-schist, or quartzitic mica-schist into gneiss; *green tuffose shale* and *slate*, through hornblende schist, chlorite schist, or talc schist into basic gneiss; *argillaceous shale* and *slate*, through argillite into mica-schist or chloritic mica-schist, and from that into gneiss; *felstone* or *felsitic tuff*, through felsitic schist into steatitic schist, garnetiferous schist, or talcose schist.

L. Gneiss.—When typical, a crystalline granular aggregate of quartz, felspar, and mica; occurring in leaves or plates, more or less parallel to one another. Typical gneiss is rather uncommon, as usually various other minerals besides the quartz, felspar, and mica occur, disarranging the regularity of the leaves.

Varieties in Composition.

A. FELSPATHIC and FELSITIC GNEISS, having one or more felspars as the predominant ingredients.

These may be divided into,—*a. Orthoclase*, *b. Oligoclase*, *c. Alpinite* (Simler), and *d. Adularia-gneiss*.—Cotta.

c. is named after the *Alps*. It is principally made up of a variety of oligoclase, and is a gneissoid aggregate of quartz, oligoclase, and an undetermined flaky green mineral. The latter Cotta suggests “probably belonging to the mica species.”

A variety of felspathic gneiss, having adularia in the place of orthoclase, is mentioned by Cotta.

Felspathic gneiss is the passage-rock between gneiss and felsyte schist.

B. MICACEOUS GNEISS, containing much mica.

Its varieties are,—*a. Two-mica Gneiss*, *b. White Mica Gneiss*, *c. Black Mica Gneiss*.

Varieties might also be named after the different kinds of mica.

Micaceous gneiss is the passage-rock between gneiss and *mica-schist*.

C. QUARTZITIC GNEISS.—This is the passage-rock between gneiss and *quartz-schist*, or *quartzite*.

D. BASIC GNEISS has numerous varieties, according to the most prevailing accessory, or the mineral that gives character to the rock mass. The most particular are, *a. Hornblendic* or *Amphibolic*; *b. Chloritic* or *Ripidolitic*; *c. Talcose*, and *d. Protogene Gneiss*, when talc replaces the whole of the mica.

NOTE.—It is said by King (India), that to a gneissoid rock, composed of quartz, felspar, and talc, the name Protogene was first given, and not to an intrusive rock. Hornblendic gneiss is sometimes erroneously and unscientifically called syenitic gneiss, while chloritic gneiss and talcose gneiss have absurdly by some been classed together under the name of Protogene gneiss. (See note on the *Hornblendic Granite*, page 33.)

E. IOLITIC GNEISS, **F. HEMATITIC GNEISS**, and **G. GRAPHITIC GNEISS**, may be thus named respectively, if iolite, micaceous iron-ore, or graphite replace most of, or the whole of the mica.

Structural Varieties.

H. COMMON or **GREY GNEISS** may be divided into,—*a. Compact Gneiss*, *b. Slate-gneiss*, *c. Ribbed*

or *Lagen Gneiss*, and *d. Fibrous* or *Stangel Gneiss*.

Compact gneiss is usually a more or less indurated grit, or in some cases it may be a slightly altered *bedded* felstone. In *Slate gneiss* there is a distinct slaty cleavage in a fissile mass of small even-grained particles, the mica being seen on the cleavage-planes alone. *Ribbed gneiss* is when the constituents occur in thin layers, and form a riban; while in *Fibrous*, or *Stangel gneiss*, the ingredients are placed "in a fibrous manner towards one direction; so that a peculiar linear parallel conformation is produced." In a variety of fibrous gneiss the quartz particles are of an elongated almond-shape, and a cross section of the rock has a peculiar aspect.

I. OBLIQUE GNEISS, having as varieties,—*a. Curled Gneiss*, *b. Nodular Gneiss*, and *c. Conglomeritic Gneiss*.

The straight linear arrangement of the leaves of the constituents may be deflected by various causes; in some by oblique lamination, or by cleavage in the original rock; in others by a crumpled foliation, or spheroidal structure; also by the original lamination curling round contained nodules or blocks, if the rock before alteration was nodular or conglomeritic.

In ordinary gneiss, the direction of the foliation seems to be due to structure in the original rock; while in granitoid gneiss, it nearly always is linear, parallel, and regular, apparently being a more or less introduced structure. Conglomeritic, and some nodular rocks, appear capable of resisting this new arrangement longest, for in many rocks of these kinds, even when very granitoid, the foliation will

remain deflected or curled in the immediate vicinity of inclosed blocks or nodules; while in some granites regular or irregular masses, differently constituted to the rest of the rock, will occur.

In West Galway, Ireland, conglomeritic gneiss is usually associated with metamorphosed igneous rocks, and probably is metamorphosed conglomeritic tuff, or agglomerate.

J. PORPHYBITIC GNEISS; K. GRANITOID GNEISS.—

These subgroups are allied, nearly all porphyritic gneiss being also granitoid.

Gneiss may have large and prominent crystals of felspar disseminated through the mass. This usually occurs in the granitoid subgroup, but not always. In some granitoid gneiss there is a distinct foliation, while in others it is indistinct.

The degree of distinctness of the foliation of the granitoid gneiss appears due to the intensity of the metamorphic action to which it was exposed. Granitoid gneiss is the passage-rock between gneiss and granite.*

M. Schist [Gr. *schistos*, split or divided].—

A crystalline fissile aggregate, consisting, when typical, of two well-developed minerals, occurring in thin plates or leaves. Mica is usually one of these minerals; however, various others also occur, sometimes as accessories, but locally as essentials, thereby forming different subgroups and vari-

* Hereafter microscopists may prove that granitoid gneiss ought more properly to be classed with the Granitic rocks (Class No. 1), and taken out from the Transition rocks (Class No. 4), as in it the original structure is obliterated, except in a few rare exceptions. The mineral constituents seem to be altogether *crystalline*, no felsitic or feldspathic tuff occurring in *granitoid gneiss*.

eties. When mica is absent, quartz is nearly always present.

Schist is easily separated along its foliation-planes, while across them it is tough, and breaks with from an uneven to a hackly fracture.

A. ARGILLYTE, or ARGILLITE, *Argillous or Clay Schist.*

—A fissile more or less indurated argillous rock, having the planes of the most conspicuous structure in the original rock, glazed, mineralized, or micacized. Certain minerals, as chialtolite, phyllite, pyrite and such like, are developed in some, and peculiar structures in others ; thus forming varieties.

Varieties in Composition.

a. *Chialtolitic Schist.*

b. *Phyllitic Schist*, or *Ottrelitic Schist.*

c. *Carbonaceous Schist.*

d. *Alum-schist.*

e. *Spilyte* or *Spilite*, or *Calcareous Schist*, and

f. *Dolomitic Schist.*

Chialtolitic schists are mostly argillous rocks ; nevertheless some are micaceous. The crystals are usually disseminated through the mass quite irregularly, but in some varieties they are stellated, usually on the surfaces of the original structure of the rock. *Phyllitic schist* contains laminæ of phyllite (*Ottrelite*). *Carbonaceous schist* is rich in carbon, and often is pyritous ; of it *Alum-schist* seems to be a variety, being due to the decomposition of pyrite in pyrito-carbonaceous schist. *Spilyte* and *Dolomitic schist* are passage-rocks into schistose limestone and schistose dolomite.

Structural Varieties.

g. Folded or frilled Argillyte, h. Contorted, and i. Fibrous.

Variety *g* has a folding or crumpling that gives it a frilled aspect: this peculiar folding is only found in some of the schists, and evidently is solely due to the metamorphic action, as it does not follow any structure in the original rocks. In variety *i* the particles are so linear and parallel-arranged as to give the rock a woody aspect, a weathered block looking like the butt of a saw or some such tree.

Argillite is the passage-rock between normal clay-slate, or shale, and mica-schist. It occurs in various stages of metamorphism. Sometimes the whole mass appears homogeneous, differing only from clay-slate, or shale, by its superior lustre; nevertheless it is rarely without traits, more or less characteristic, of mica-schist; such as the crumpling, crushing, folding, twisting or contortion of the laminæ; also the hardening of the joint and other lines, with the development of minerals, especially on the surfaces of beds or on the lamination and cleavage-planes.

B. QUARTZYTE, or QUARTZITE, Quartz-schist.—Chiefly consists of quartz with some mica; the latter being most conspicuous on the planes of foliation.

Varieties in Composition.

a. Itacolumyte (Eschwege), *Pedra elastica* (Anchietà), *Micaceous Quartzyte*.—"A fine-grained and at the same time schistose compound of

quartz with some mica, talc, or chlorite.”—*Cotta*. Henwood says of the Itacolumyte of Minas Gerais, “quartzose talco-micaceous slate.”

These flexible quartzytes were first noted among the metamorphic sedimentary rocks of Brazil, and were named after Itacolumi Peak. Burton says there are three different kinds of rock named after this peak: 1st, the flexible quartzyte, or *Pedra elastica*, under which name it was described nearly three centuries ago by Padre Anchieta; 2nd, *Diamantine itacolumyte*, the matrix of the diamond, “a hard talcose rock of distinctly laminated quartz, white, red, or yellow, granular, with finely-disseminated points of mica;” and 3rd, in Minas the name is applied to “refractory sandstone grits, a fine crystalline rock evidently affected by intense heat.” The peak itself consists of none of the three kinds, although all are called after it.

Jukes thus describes Itacolumyte:—“A genuine unaltered sandstone, more or less micaceous, like other sandstones, but the mica in worn spangles, not in connected flakes.” The rock, of which this is a description, came from India. In it are lines that appear due to deposition; when placed under the microscope, it is found to be full of long drusy cavities, which lie in lines rudely parallel with the structure of the rock. The cavities open and shut when the slab is bent. According to Dana, Itacolumyte pertains to the talcose series, and is the matrix of the diamond; this evidently is Burton’s second kind of rock.

b. *Itabiryte* (Eschwege), *Jacotinga*.—Ferruginous quartz-schist, a variety of Itacolumyte, named after Itabira, in Brazil. Eschwege

who first described the rock, makes it the matrix of the diamond. Some very ferri-ferous varieties are said to be worked in Brazil as iron-ores; they are more or less auriferous. Itabirite, Dana says, "contains much specular iron-ore in grains or scales, or in the micaceous form." *Jacotinga* Henwood describes as an "auriferous micaceous iron-schist."

- c. *Felsitic Quartzite*—An aggregate of quartz and felsite, sometimes also felspar and a little mica; usually more or less massive, rarely well foliated.

Structural Varieties.

d. *Quartz Rock* (Jukes), *Granular Quartzite*.

e. *Fibrous Quartzite*, and

f. *Conglomeritic Quartzite*.

Variety *d* is fine-grained, homogeneous, and more or less saccharoid in aspect, often merging into felsitic quartzite (*c*). *e*. has an arrangement in long rude prisms like coarse wood; while *f*. contains pebbles, usually ovate, but sometimes more or less angular; this is evidently a metamorphosed conglomerate, the normal form of the pebbles having been elongated by the subsequent metamorphic action.

Quartzite or Quartz-schist (*B*) is an undoubted metamorphic sedimentary rock, originally a highly siliceous sandstone or grit. There are, however rocks described as Quartzite or Quartz-rock that seem not to be metamorphic rocks, but rather normal rocks, either deposited from a solution, or *perhaps* a variety of intrusive rock. Some of these

quartz-rocks appear to occur as masses in unaltered rocks.

Felsitic quartzyte is a remarkable variety; some would seem to be metamorphosed *Petrosilex*; while others merge into quartz-rock on the one hand, and into Felsitic granite or Felsite-rock on the other. Some also occur in bedded masses, when they may possibly be metamorphosed feldspathic sandstones.

C. FELSYTE SCHIST.—A felsitic fissile rock; colour whitish, greyish, or greenish; from compact to mealy or granular in texture.

NOTE.—Some of the compact varieties of Felsyte schist are identical with rocks named by Krantz as *Leptinyte* or *Whitestone*; on the other hand, Cotta classes *Leptinyte* with *Granulyte*, apparently considering both as intrusive rocks.

Varieties in Composition.

a. Micaceous, b. Quartzose, and c. Pyritous.

Varieties in Structure.

- d. Ribaned*, when the mineral constituents are in layers or seams, alternating with one another.
- e. Mealy*, having a leafy or scaly structure; and
- f. Gneissoid or granular*, a granular felsitic rock, slightly fissile, with grains of quartz disseminated through the mass, also a little mica.

NOTE.—Gneissoid or granular felsyte schist merges into felsitic quartzyte. In some localities it is an undoubted bedded or interstratified rock, while in others it is evidently intrusive. Perhaps in the first case it may be a metamorphosed stratified felstone or tuff (which could only be determined by a microscopical examination), while in the latter it may be a metamorphosed intrusive felstone. The latter variety of granular felsyte schist appears to answer the description of the rock called *granulyte* by Cotta.

D. MICA-SCHIST.—When typical, a foliated or fissile aggregate of quartz and mica. Usually the foliation is regular; nevertheless it may be crumpled, crushed, folded, twisted, contorted, curled, nodular, or spheroidal.

Varieties in Composition.

- a. Two-mica Schist, b. Chloritic, c. Tulose, d. Garnetiferous, e. Chiastolitic, f. Andalusitic, g. Amphibolic, h. Epidotic, i. Quartzose, j. Felsitic, k. Calcareous, l. Schorlaceous, m. Pyritous, n. Pyrrhotitic, o. Hematitic, and p. Graphitic.*

The varieties in composition are named after conspicuous minerals, locally essentials. Most of these rocks, however, are only passage-rocks into the other subgroups of schist; some even (such as *Epidotic mica-schist*) are not normal rocks, but due to a secondary change in the mineral constituents.

Structural Varieties.

- q. Fine Mica-schist*, where all the ingredients are small and even-grained.
r. Ribbed, with the minerals in layers or bands.
s. Fibrous, with a woodlike arrangement.
t. Nummoid, having in it small discs of quartz, giving the rock an appearance as if studded over with pieces of small coin.
u. Wavy.
v. Frilled, having the foliation folded or crumpled on itself.
w. Curled or Spheroidal.
x. Knotty or Concretionary.
y. Nodular; and
z. Gneissoid; the last-named variety being the passage-rock between mica-schist and gneiss.

E. BASIC SCHISTS.—Schist in which such minerals as amphibole, ripidolite, talc, and the like, replace the whole, or nearly the whole, of the mica, and become essentials of the rock.

Varieties are,—*a. Hornblende, or Amphibole Schist*; *b. Actinolite Schist*; *c. Chlorite or Ripidolite Schist*; *d. Chloritoid Schist* (Hunt); *e. Potstone*; *f. Talc Schist*; *g. Garnet Schist*; *h. Tourmaline, or Schorl Schist*; and *i. Rhætitic or Magnesia Schist*.

It is evident that with more mica, any of these rocks may pass into one of the varieties of mica-schist; or that one of the varieties of mica-schist, by a loss of its mica, may pass into one of the basic schists. Sterry Hunt describes a dark-coloured Canadian subvariety of chlorite-schist as largely composed of chloritoid, a mineral allied to ripidolite and phyllite; and Cotta describes potstone as a felt-like web of ripidolite, and rarely foliated. Perhaps these two varieties of schist would be more properly classed as varieties of argillyte. Rhætitic schist is described by Forbes as remarkable for the predominance of silicates of alumina and magnesia, rhætizite, iolite, chiastolite, &c.

The basic schists have numerous structural varieties. They are, however, very similar to those mentioned when describing mica-schist; therefore it is unnecessary to enumerate them here.

F. METALLIC SCHISTS.—Schists, in which a mineral ore replaces part or the whole of the mica, and gives a character to the rock mass.

The principal varieties are,—*a. Mico-iron Schist*, *b. Pyrite Schist*, and *c. Pyrrhotite Schist*: other ores may also give a character to a rock.

G. **BURNT SHALE.**—Altered argillous rocks, due to the burning of carbonaceous beds.

Varieties are,—*a. Burnt Clay*, *b. Rock Slag*, and *c. Porcelanyte*.

H. MINERALS FORMING ROCKS.

- a. Quartz and Jasper*, generally in veins.
- b. Corundum*, in subordinate layers.
- c. Apatite*, sometimes forms compact spheroidal masses.
- d. Magnesite*, in subordinate masses.
- e. Orthoclase*.
- f. Pistacite*; and
- g. Pycnite*, generally in veins.
- h. Lepidolite*, rarely forms independent rocks.
- i. Liévrte or Ilvaite*, in subordinate beds.
- j. Magnetite*.
- k. Anthracite*.
- l. Graphite*.

Normal graphite is pure carbon with a little oxide of iron mechanically mixed. Graphite occurs in beds and imbedded masses, laminæ, or scales, in granite, gneiss, schist, and schistose limestone. Of it Dana says, "It is, in some places, a result of the alteration by heat of the coals of the coal formation."

- m. Eisenkiesel*, "a concretionary rock, made up of red and brown iron-ore, and fragments of bright red jasper."—*Bauerman*.

n. Garnet Rock.

One variety is massive, of a milk-white colour, very hard and heavy, with a peculiar graining *through the mass*. Others are brownish or reddish, *more or less crystalline*; these usually pass into

a highly crystalline vuggy rock, that has as accessories, pyrite, marcasite, chalcopyrite, epidote, calcite, &c. : occurs in veins and subordinate masses.

N. **Slate** [according to Worcester, originally spelt "Sclate." Old Fr. *esclat*, a splinter, a shingle. Celtic *scorlt*, to split or cleave].—Originally shale or clay, into which a subsequent slaty cleavage has been introduced.

Some slates are said to belong to the Metamorphic series. The slaty structure, however, seems more connected with Sedimentary than with Ingenite rocks. Slates often occur associated with unmetamorphosed sedimentary rocks, and many metamorphic rocks are known to have been cleaved prior to their being metamorphosed. Slates are mentioned here, but they will be described hereafter among the Derivate rocks, as varieties of the subgroup *Argillous rocks* (see page 103).

O. **Calcareous Rocks** (Lat. *calx*, *calcis*, lime).

A. **SCHISTOSE LIMESTONE**.—A crystalline granular or nearly compact aggregate of calcite, with more or less ripidolite, mica, felsite, quartz, pyrite, galenite, &c. ; often more or less fissile, sometimes highly schistose; usually the foreign minerals are in more or less regular lines or layers, but in some varieties they are disseminated through the mass of the calcite.

a. *Cipollino*, or *Micaceous Limestone*.—Rich in mica, which gives the rock a schistose texture; this variety graduates into Spilyle.

b. *Calciiphyre* (Brongniart), or *Garnetiferous Limestone*, containing garnets and pyroxene or felspar.

- c. *Hemitrene*, or *Amphibolic Limestone*. In addition to the amphibole, it generally contains grammatite.
- d. *Felsitic Limestone*.—The felsite often occurs in highly crumpled parallel layers.
- e. *Statuary Marble*, or *Saccharoid Limestone*.—
“A white, fine-grained rock, resembling loaf sugar in colour and texture, working freely in any direction, not liable to splinter, slightly translucent, and capable of taking a polish. Flakes of mica or chlorite sometimes exist in it.”—*Jukes*.

Typical statuary marble should answer the above description, but the rock in mass is often more or less schistose, and also streaked with dark lines, caused by foreign minerals being associated with the calcite.

- f. *Granular Schistose Limestone*.—A granular crystalline, more or less schistose limestone; containing some mica, ripidolite, felsite, or even quartz. Colour white, pale grey, dove, light blue, greenish, reddish, purplish, or yellowish. It is often impregnated with microscopic crystals of pyrite, marcasite, galenite, or chalcopyrite.

P. Pseudomorph Calcareous Rocks.—Most of these rocks are pseudomorph metamorphic rocks; some, however, may be derived direct from the original rocks.

- A. **SCHISTOSE DOLOMYTE.**—A crystalline aggregate of calcite and dolomite, together with some ripidolite, talc, or mica: pyrite, marcasite, or chalcopyrite are usually present in greater or less quantities, but minutely disseminated

through the mass. The rock, though usually schistose, may be sometimes compact or granular; when compact, it breaks with a conchoidal fracture. The colours of these dolomytes are very various: a handsome variety is milk-white. Many of them weather into a micaceous ferriferous sand.

Some varieties are,—*a. Micaceous*; *b. Pyritous*; *c. Saccharoid*; and *d. Granular*.

The most marked varieties in composition are those due to an excess of mica, or one of the pyrites; and in structure, the saccharoid and the granular. The latter are somewhat similar in aspect to saccharoid and granular limestone, but harder, although more easily decomposed by meteoric action.

Ophyte and steatite occur associated with the calcareous rocks, but more especially with dolomite, they being due to pseudomorphic action. At Lissoughter, Galway, Ireland, blocks of dark green very pure serpentine, coated with a thick crust of rusty yellow impure meerschaum (locally called *Cream*), are found on or close to the surface of the ground; lower down, the solid rock was an opicalcyte, often micaceous or felsitic; while in depth the opicalcyte passed into a more or less schistose dolomite. Dana, Jukes, and others also mention dolomytes which have been noted as graduating into ophytes. It should be borne in mind, that the ophytes and steatites now to be described are pseudomorph sedimentary rocks, while those previously mentioned (*see* page 63) were pseudomorph igneous rocks.

B. OPHYTE [Gr. *ophis*, a serpent], *Serpentine*, *Verd antique*, *Ophiolyte* (Dana and Hunt).—When

pure, a compact, dark green aggregate of *serpentine*, usually associated with calcite, dolomite, and magnesite; and with such minerals as talc, ripidolite, magnesian mica, &c.; unctuous to the touch on fresh fractures.

The most common varieties are,—*a. Ophidolomyte*; *b. Ophimagnesyte*; *c. Ophicalcyte*; and *d. Ophyte Schist*.

The ophytes have been classed by Hunt: 1st, Dolomitic, 2nd, Magnesitic, or 3rd, Calcitic, according to whichever mineral gives a character to the rock-mass. To these may be added the schistose varieties, as all ophytes of this class become in places schistose, they being impregnated with, or having layers of, mica or some other conspicuous ingredient not necessary to a typical ophyte, but giving a more or less marked schistose character to the rock-mass.

C. STEATYTE, OR STEATITE [Gr. *stear*, *steatos*, fat].—

A grey, blue, green, reddish or whitish fissile aggregate of steatite, with some talc, ripidolite, amphibole, or magnesian mica: has a mealy aspect; very unctuous to the touch; cuts easily, but tough.

The principal varieties are,—*a. Felsitic*; *b. Hornblendic*; *c. Pyroxenic*; *d. Micaceous*; *e. Steatyte Schist* (this class of steatyte always graduates into a schistose rock); *f. Pyrallolyte* (Nordenskiöld), greyish to clouded milk white in colour, compact, translucent, steatitic rock. According to Hunt, it is identical in composition with talc.

Pyrallolyte occurs associated with ophyte, but *seems to be only a secondary rock filling up*

“shrinkage - fissures,” or open fault-lines. In Connemara, Ireland, there is a variety having dendrites disseminated through it. It is manufactured into ornaments, and called *Moss Serpentine*.

g. Onkosin (Kobell), *Agalmatolyte* (Sterry Hunt).

—A greenish or greyish corneoid steatitic rock. It, at first, was taken for ophyte.

D. MAGNESYTE.—An aggregate of carbonate of magnesia and carbonate of iron, but may contain felspar, mica, quartz, chrome, nickel, &c.

E. SMARAGDITE SCHIST.—An unctuous fissile rock, an aggregate of smaragdite and mica, but often containing felsite, quartz, and other foreign substances. One variety is *garnetiferous*, and is similar to, if not identical with, the micaceous eklogyte of some authors. Usually smaragdite schist is associated with ophyte, steatyte, or talc, but sometimes it forms independent beds.

PART III.

Order II.—*DERIVATE ROCKS.*

Class I.—*SUBAQUEOUS ROCKS.*

THE name of this class of rocks points to their having been deposited either mechanically or chemically in water [Lat. *sub*, under; *aqua*, water].—The materials of which they are composed have been derived from previously existing rocks that were denuded by marine, meteoric, or glacial abrasion, or by a chemical process.

A. Rocks for the most part mechanically formed.—This group includes all the rocks principally due to mechanical deposition. The detritus from which they were formed varies from an impalpable powder to pieces of considerable size, all being formed by one or more of the denuding forces,—marine action, meteoric abrasion, or glacial action. The first of these forces not only destroys, but it also reconstructs, while the detritus due to meteoric and glacial abrasion, prior to being formed into rocks, have to be transported by wind, “rain, and rivers,” or some other carrier, for deposition and rearrangement at the bottom of seas or other waters.

The mechanically-formed subaqueous rocks make two well-marked subgroups; namely, *Arenaceous* or *sandy*, and *Argillaceous* or *clayey*.

A. ARENACEOUS ROCKS [Lat. *arena*, sand].—Consisting for the most part of siliceous or quartzose particles and fragments, sometimes loosely piled together, but often united compactly, in a typical arenaceous rock by a siliceous matrix. Some sandstone varieties of these rocks may have a calcareous, ferruginous, or some other such base.

a. SAND, GRAVEL, and SHINGLE.—Of loose accumulations of water-worn fragments Page thus writes: "When the fragments are less than the size of a pea they are called *Sand*; when the pebbles vary from the size of a pea to that of a hen's egg, they are called *Gravel*; and when the fragments are larger, *Shingle*."

Usually the fragments in *sand*, *gravel*, or *shingle* are more or less rounded; nevertheless, sometimes they are angular. Sand, gravel, and shingle more generally occur as surface deposits; sometimes, however, they belong to much older formations.

NOTE.—The basal beds of the Permian system, and in some cases beds in the Carboniferous and Silurian systems, are a *gravel* or *shingle*, while sands are not uncommon in different groups of rocks.

Local Names for Gravel and Shingle.

Rubble, "rough angular gravel, either loose or compacted into stone."—*Jukes*. *Brash* (English), *Foundation* (Irish), the broken and angular fragments of any rock, lying on, and separating it from, the overlying drift.

b. CONGLOMERATE [Lat. *con*, together, and *glomerare*, to gather].—A fine or coarse matrix

containing numerous round or roundish blocks, pebbles, or fragments of one or more varieties of rocks.

Subvarieties in composition may be called (a.) *Calcareous*, (b.) *Hematitic*, (c.) *Cupriferous*, (d.) *Pyritous*, (e.) *Plumbeous*, (f.) *Shale Conglomerate*, having an indurated clayey or shaly matrix, (g.) *Slate Conglomerate*, where blocks and fragments are inclosed in a well-cleaved or slaty matrix.

In the *Metalliferous Conglomerates*, the matrix for the most part is a metallic ore. They may be *fer-riferous*, or containing iron-ore; *cupriferous*, or coppery; and *plumbeous*, or leady. These conglomerates are not uncommon in places in the Lower Carboniferous: some of the plumbeous in the New Red are said to be valuable lead ores.

Special names might also be given to subvarieties from the contained fragments, if the conglomerate only contains pieces of one kind of rock, or the fragments of one kind of rock predominate. Such names, however, would be too numerous to mention.

c. BRECCIA [Ital., a crumb or fragment].—An agglutination of angular fragments of one or more kinds of rocks.

The subvarieties of Breccia are somewhat similar to those of conglomerate; namely, (a.) *Calcareous*, (b.) *Hematitic*, (c.) *Cupriferous*, (d.) *Pyritous*. There is, however, a peculiar subvariety, (e.) *Shale Breccia*, when in a more or less calcareous base are inclosed pieces of shale.

These inliers do not appear originally to have

been fragments of a shale rock, but rather to have been pieces of clay that were caught up in the calcareous matrix, and subsequently became indurated and shaly.

- d. SANDSTONE and GRIT. — Sand consolidated by extreme pressure or by the aid of a cement. When typical the grains are quartzose, but in all of the subvarieties there are admixtures of one or more other substances. If the rock has a distinct granular structure, or is capable of being ground or separated into grains, it is called *Sandstone*; while, if the rock is hard and compact, the grains not being visible to the naked eye, or if the grains of sand are so firmly compressed or cemented together that it is impossible to separate them, then the rock is classed as a *Grit*.

The subvarieties of sandstone and grit are very numerous. In composition they may be—

- (a.) *Calcareous*, (b.) *Argillous*, (c.) *Felspathic*, (d.) *Micaceous*, (e.) *Hematitic*, (f.) *Cupriferrous*, (g.) *Pyritous*, and (h.) *Tuffose*.

The sandstones are more often argillous, felspathic, micaceous, and tuffose, than the grits.

- Of structural subvarieties there are—(i.) *Pebbly*, (j.) *Cleaved*, (k.) *Oblique* or *Obliquely foliated*, (l.) *Concretionary*, (m.) *Spheroidal*, (n.) *Nodular*, (o.) *Lenticular*; and of sandstone only (p.) *Friable*.

Sandstones necessarily are oftener oblique, concretionary, spheroidal, nodular, and lenticular, than grits. Friable sandstones have such a weak matrix that the grains separate and the rock easily returns

to its original condition of sand. Many of these have a concretionary or spheroidal structure, and contain hard spheres or concretions ; while some compact sandstones inclose spherical masses of friable sandstone, or even sand. A *quartzose* or *hard sandstone*, correctly speaking, is a grit : there are some grits so quartzose as scarcely to be distinguished from *quartzite*.

- e. FLAGSTONE, or FLAGS.—A grit or sandstone which was deposited in layers, from a quarter of an inch to three or four inches in thickness. The rock now splits, or is capable of being split, along the lamination or stratification planes, into tabular plates or flags.

The subvarieties are—(a.) *Arenaceous Flagstone*, and (b.) *Argillous*, according to whether they are sandy or clayey. Usually the layers are parallel to the true stratification-planes or the bedding, but not always, as in some cases the flags are due to oblique lamination.

NOTE.—In some rocks flaglike masses are due to cleavage, while in others to joint-planes. King (India) has proposed that the term *Flag* should be restricted to tabular masses due to bedding or stratification planes, while thin flat masses, caused by either cleavage or jointing, should be called *Slabs*.

- c. *Slabstone*, or *Slabs*.—Cleaved or finely parallel-jointed rocks, which split into tabular plates or *slabs*, from one inch to three or four inches in thickness. Slabs are rarely as strong as flags.

Local Terms for Sandstones and Grits.

Catsbrains, sandstones traversed in every direction by little branching veins of *calcite*. *Rotch*, *friable sandstone*. *Rock* (quarrymen), any very hard

grit. *Freestone*, or *Free* (Ulster), a stone that cuts easily in every direction. *Hazel* (North of England), hard grit. *Binder* (Cork) and *Post* (North of England), beds of grit in shale, slate, or clay. *Peldon* (South Staffordshire), *Calliard* or *Galliard*, and *Crowstone* (North of England), a hard, smooth, flinty grit. *Brownstone* (Munster), grit and sandstone. *Red free* (Ulster), New red sandstone. *Granât* (Munster), coarse quartzose grit. *Brass-binder* (Cork), a thin pyritous grit. *Brass-balls* (Munster), *Stone-mine* (Leinster), nodules of pyritous grit. *Catsheads* (Leinster), nodules of hard grits in shale. *Cornstone*, containing calcareous concretions.

B. ARGILLOUS ROCKS [Lat. *argilla*, clay], consisting for the most part of more or less argil, or clay. Other minerals and substances, however, often occur in the mass.

- a. CLAY, LOAM, MUD, and SILT.—“Perfectly pure *clay* is a hydrated silicate of alumina, and when pure very plastic. *Loam* is a soft and friable mixture of clay and sand, enough of the latter being present for the mass to be permeable by water, and to have no plasticity.”—*Jukes*. *Mud* and *silt* are more or less dry or moist mixtures of clay, loam, and sand; usually with decayed vegetable and animal matter.

The *subvarieties* of clay are very remarkable, and will require special descriptions. *Marl*, or *calcareous clay*, properly speaking, should be classed as a *subvariety* of clay. It seems preferable, however, to place it among the varieties of the argillous rocks, on account of its importance.

- a. *Kaolin*, or *Porcelain clay* (see *Kaolin* among the *Ingenite Rocks*); (b.) *Potter's*, or *Pipe-clay*, and (c.) *Brick-clay*.

Pure, or nearly pure clay, free from iron, usually of a whitish or light bluish-grey colour, is called *Potter's clay*. It is very plastic.

The best quality of *tiles* and *bricks* are made from *Potter's clay*; nevertheless, many impure clays are capable of being burned into bricks, and are commonly known by the name of *Brick-clay*. If clay is impure, owing to the presence of other silicates besides that of alumina, the bricks when burnt will be more or less ill-shaped and unsightly. In the neighbourhood of many bogs in Ireland is a deposit of dark sandy clay, locally called *Doab* (Connaught), used for making floors, or for plastering when mixed with a little lime.

- d. *Fire-clay*.—Clay containing much silica or fine sand, nearly or quite free from iron; often blackish from containing carbonaceous matter, which, however, does not prevent it from burning into bricks or tiles which stand the fire.

Of *Fire-clay* Jukes writes: "It is probable that in good fire-clays, the silica and alumina exist in just that definite proportion which would form a true silicate of alumina."

Local Terms for Fire-clay.

Seat, *Coal-seat*, *Coal-clay*, *Seat-clay*, *Under-clay*, *Buddagh*, *Soft-seat*, *Spavin*.—In Munster the term *Seat* is often used to denote the nearest bed of *clunch*, *grit*, or sandstone under a coal; while the *intervening fire-clay* is called *Soft-seat*, &c. In

Leinster a highly carbonaceous, soft, muddy-looking fire-clay is called *Buddagh*, and for a similar rock the term *Spavin* is used in Yorkshire.

- e. *Clunch*.—A tough more or less sandy indurated clay; characteristic of the Coal-measures and the Gault, but also found in other formations.

In Munster clunch is usually known as *Hard-seat* or *Seat-rock*, as it often occurs below a coal or fire-clay. These names, however, are not always restricted to it, for sometimes a grit or sandstone underlying a coal or fire-clay, may be so called. In Yorkshire and Lancashire a very hard siliceous variety is called *Ganister*.

- f. *Clay-stone*; (g.) *Clay-rock* (Jukes).

Clay-stone is a compact and tolerably solid mass, chiefly consisting of clay. It may be laminated, but not slaty; its fracture earthy. *Clay-rock* is highly indurated, or solidified clay or clay-stone; it is not laminated like *shale*, nor cleaved like *slate*. It may pass into shale by merging into a laminated mass; or into slate, by becoming cleaved.

- h. *Lateryte*, or *Laterite*, *Brick-stone* [Lat. *later*, a brick], a brick-coloured (reddish or brownish) arenaceous clayey rock, sometimes conglomeritic; soft before being exposed to atmospheric influences.

Lateryte is a *Tertiary rock* of India, which, while soft, is cut into large brick-shaped masses (hence the name), which on exposure to the air, harden. These are extensively used for building purposes. Some of the *Lateryte* from Kutch is evidently *tuffose*, or formed for the most part from the fine

débris of igneous rocks. A similar character belongs to rocks from other Indian localities, while other varieties seem not to contain any tuffose particles; usually it appears to be more or less associated with bedded or stratified igneous rocks. Lateryte may be very ferruginous, even so as to be classed as an *Earthy Limonite*. In Antrim, Ireland, associated with Igneous rocks of tertiary age, are rocks that seem to be identical with some of the Indian laterytes; while in Mayo, Ireland, there are somewhat similar rocks interbedded with Eurytes, of Upper Silurian age. Some varieties of Lateryte, both in Kutch and Antrim, appear to pass into a *Steatitic rock*, or even *Steatyte*.

- i. *Fuller's Earth* (see Fuller's earth among the Ingenite rocks).

Other subvarieties of clay are—(j.) *Bituminous*, (k.) *Saliferous*, (l.) *Variegated*, (m.) *Hematitic*, (n.) *Ferruginous*, and (o.) *Book*, or *leaf clay*, deposited in thin leaf-like laminæ.

- b. **MARL**, *Calcareous Clay*, an aggregate of clay, with more or less lime; earthy, compact, or fissile. Disintegrates when exposed to the atmosphere, and effervesces with acid.

Subvarieties in composition are—(a.) *Very calcareous*, the passage-rock into soft limestone; (b.) *Dolomitic*, (c.) *Very argillous*, (d.) *Arenaceous*, (e.) *Micaceous*, (f.) *Oil-slate*, or *Oelschiefer*, a bituminous dark-coloured subvariety; (g.) *Glaucinitic*, of a green colour; and (h.) *Gypseous*, containing strings or thin laminæ of gypsum.

In structure there are—(i.) *Shell marl*, and (j.) *Mealy*, or *Friable marl*. The latter, when

typical, is due to lime in solution, being deposited by the evaporation of water. On the other hand, *Shell marl* is partly due to the decomposition of shells, and partly to an admixture of mealy marl. Others are—

(k.) *Marl-stone*, a more or less indurated marl, without either lamination or cleavage; disintegrates when exposed to the atmosphere; (l.) *Tutenic*, or *Cone-in-cone marl*, that has a cone-in-cone structure; these concretions being called *tuten* in Germany; (m.) *Book*, or *Leaf marl*, a marl deposited in very thin laminæ; a cross section having the appearance of a closed book.

- c. SHALE [Ger. *schalen*, to peel or shell off].—A compact, fissile, clayey rock; its fissile texture due solely to the rock having originally been deposited in laminæ or leaves. It is very variable in colour, and usually contains many impurities.

Subvarieties in composition are—(a.) *Calcareous*, (b.) *Arenaceous*, (c.) *Micaceous*, (d.) *Bituminous*, (e.) *Carbonaceous*, (f.) *Hematitic*, and (g.) *Cupriferous*. Structural varieties are (h.) *Flaggy*, (i.) *Conglomeritic*, (k.) *Nodular*, (l.) *Concretionary*, (m.) *Spheroidal*, (n.) *Lenticular*, (o.) *Mudstone* (Murchison), shales which on being exposed to the atmosphere, “rapidly decompose, and are converted into their primitive state of mud;” and (p.) *Tutenic* (cone-in-cone).

- d. SLATE, *Sclate*, *Clay-slate* [originally spelt *sclate*, old French *esclat*, a splinter, a shingle; Celtic *Scorlt*, to split or cleave]. A shale or clay-rock in which a more or less perfect

slaty cleavage has been superinduced. The cleavage sometimes is much distorted.

Some slates (see *ante*, page 89) may be in part of metamorphic origin; most slates, however, are more nearly allied to the shales than to any other kind of rock. In all slaty or cleaved rocks there is a superinduced structure which gives the rocks a tendency to split into thin plates, whose direction is generally more or less oblique to the planes of the stratification; however, this may be perpendicular to, or even coincident with, these planes. Cleavage, as proved by Haughton and others, is due to the pressure, which has also produced the bending and folding of strata.

Slates may have been originally either shales or clay-rock, into which cleavage was introduced. If originally shale, there will be a lining or riban in the slate coinciding with the lines of deposition, while cleaved clay-rock without riban, is usually more homogeneous, and consequently generally makes the better and purer slate. The more argillaceous the rock, usually the more perfect the cleavage, while the slates have a submetallic ring. Good slate is called by the quarrymen *good metal*.

NOTE.—If cleavage in general is due to pressure, it should naturally be more prevalent among the older than the newer rocks; according to Agassiz, however, it may be sometimes due to induration, as that author mentions cleaved shales (*slates*), noted by him in Brazil, which he supposes to be part of the *Boulder-clay drift*. In them he records a perfect cleavage, which he suggests is due to the extreme tropical heat.

Subvarieties of slate are—(a.) *Roofing slate*, a pure or nearly pure slate, having smooth and perfect cleavage, allowing the rock to split into very thin plates, which at the same time are rigid, solid, and strong.

NOTE.—Some *so-called* slates, such as the *Stonefield slate* (which are sold for roofing purposes), are, correctly speaking, not slates, as such rocks split into plates not along the cleavage-planes but along the planes of lamination. Such miscalled slates are always coarse and unsightly, compared with true slates.

- b. *Slabs*, or *Slab-slates*.—Pure or nearly pure slate, in which cleavage is only partially developed; consequently the rocks will not split into plates thin or light enough for roofing purposes; it however splits in parallel more or less thick plates, and can be manufactured into slabs.
- c. *Pencil-slate*, *Pencil*, *Pinsill*.—Pure soft, but firm slate, which breaks easily into long spike-like prisms.

This fracture seems due to the cleavage, and two or three systems of close-lying joints, running in similar directions, and so placed as to divide up the rock into long, more or less regular narrow pieces. Lamination or foliation may take the place of one of the joint systems. These long narrow prisms are used as *pencils* for writing on slabs; whence the name.

- d. *Novaculyte*, or *Novaculite* [Lat. *novacula*, a razor], *Whetstone*, *Whetslate*, *Hone*, *Oilstone*.—A highly siliceous clay-slate or shale, perfectly compact and homogeneous; usually only indistinctly cleaved; with a conchoidal or semiconchoidal fracture along the grain of the rock, and from uneven to hackly across it.

NOTE.—Many of the *Novaculytes* seem more allied to the Unmetamorphic than to the Metamorphic argillaceous rocks; nevertheless they often occur in the vicinity of outburst of Igneous rocks, thereby suggesting that the hardening and silicification is partly or wholly due to vulcanism.

Other subvarieties of slate are—(e.) *Calcareous*, (f.) *Carbonaceous* (these often afford an even roofing-slate, but are more or less soft and affected by nodules and seams (*Bulls'-eyes* and *Rucks*) of pyrite), (g.) *Arenaceous*, (h.) *Tuffose*, (i.) *Ribbed*, and (k.) *Conglomeritic*.

The last has, in a slaty matrix, fragments, often of quartz, rarely exceeding the size of a large pea or small marble, but usually about the size of shot. This rock is sometimes erroneously called Porphyry.

Local Names for Shale and Slate.

Rock-bind, *Stone-bind*, *Fekes* (Scotland), and *Blue-rock* (Leinster), arenaceous shales. *Bind*, *Killas* (Cornish) and *Slig* or *Sliggeen* (Irish), either shale or slate. *Bury* (Kilkenny) and *Flucan* (Cornish), soft shale or clay. *Metal*, compact flinty shale. *Plate*, flaggy shale-bed. *Shingle* and *Gravel*, names used in Ireland for shale or slate that breaks up into small angular pieces; usually they cannot be blasted, but must be worked by the pick, bar, or wedge.

C. FAULT-ROCK.—Fault-rock is usually a mechanically formed rock, but in many instances it is also in part chemically formed. It may be conglomeritic, brecciated, arenaceous, argillous, shaly, or schistose (metamorphosed). Some of the metamorphosed fault-rocks are peculiar, especially the brecciated, calcareous, and dolomitic rocks.

NOTE.—Fault-rocks, properly speaking, are not subaqueous; nevertheless, it seems more appropriate to mention them here than elsewhere.

B. Rocks for the most part chemically formed.—*Such rocks*, when typical, are nearly solely due

to chemical action ; they are more common in veins, nodules, and patches, than in beds or interstratified masses. Chemically-formed rocks are always insignificant, when compared with the associated mechanically-formed rocks, and often the former graduate into the latter.

- A. HALYTE, ROCK-SALT [Gr. *hals*, salt ; *lithos*, stone].
—A crystalline, granular, translucent or transparent rock, consisting almost entirely of chloride of sodium ; colourless, or of grey, yellow, reddish, occasionally bluish, or greenish colour ; soluble in water ; taste saline. Of Halyte, Cotta writes : “ In Nature it almost always contains sulphate of lime, chloride of calcium, chloride of magnesium, and other salts ; frequently admixtures of bitumen, clay, or boracite.”

Varieties are—*a. Granular*, *b. Sparry*, and *c. Fibrous*, or with a wood-like aspect.

Halyte always merges into impure varieties, the impurities having a more or less mechanical origin.

- B. GYPSUM [Gr. *ge*, earth ; *epso*, I boil].—A hydrated sulphate of lime ; usually crystalline, but sometimes compact or fibrous ; soft ; generally white or whitish.

- a. Alabaster* [after *Alabastron*, a town in Egypt].
- b. Granular Gypsum*, always almost white, somewhat translucent ; granular and finely crystalline ; sometimes it is fissile, each bed being composed of many layers of little crystals, slightly differing in colour and texture.
- c. Compact gypsum*, rare.
- d. Fibrous*, or having a wood-like structure.

e. Selenyte, Spathic or Sparry gypsum, transparent varieties.

f. Tripestone, "a variety both of texture and composition. It is formed of thin layers of pure white gypsum, alternating with grey argillaceous gypsum, the whole twisted or crumpled to resemble a ruff." — Cotta. Gypsum merges into impure, more or less mechanically formed varieties.

C. ANHYDRYTE, or ANHYDRITE.—"A granular or compact aggregate of anhydrous sulphate of lime; harder than gypsum; white, grey, or blue."—*Cotta.*

The varieties of anhydrite are very analogous to those of gypsum, the more marked being—*a. Granular, b. Compact, and c. Fibrous.*

D. DOLOMYTE, or DOLOMITE [after M. Dolomieu].—A granular crystalline aggregate of dolomite associated with some calcite; more or less saccharoid; does not effervesce, or only slightly with dilute acid. Dolomite weathers into a ferruginous sand, *Dolomitic sand.*

NOTE.—Some dolomytes are of purely chemical origin, while most dolomytes are partly chemically and partly mechanically, or even perhaps organically, formed. Only the first belongs to this group (B), the others are described in group (C), page 120.

E. QUARTZ.—Pure or nearly pure silica.

Varieties of Quartz are—*a. Chert* [Celtic *cairt* or *chairt*, rough bark; as chert forms a rough bark or coat to the beds of limestone], *b. Flint, c. Menilyte, or Menilite.* Hard, compact, containing more or less carbonaceous matter; breaks with from a conchoidal to a semiconchoidal fracture; colour white, grey, yellowish, brown, black, and sometimes blue.

Chert, *Flint*, and *Menilyte* occur in layers of nodules, thin beds, and concretions; sometimes, however, as is especially the case with chert, they form massive strata. *Menilyte* is found in the Tertiary limestone. *Flint* is characteristic of some chalk, while chert is mostly found in the Carboniferous and older limestone, and in some shales. Jukes writes of these rocks, "Almost all large masses of limestone have their flints or siliceous concretions. Pure siliceous concretions occur even in the fresh-water limestones and gypsum beds of Montmartre."

d. Jasper and *e. Agate* (particoloured varieties).
f. BasarYTE, or *Basanite*, *Touchstone*, or *Lapis Lydius* (Pliny), a velvet-black cherty variety, used on account of its hardness and black colour for trying the purity of the precious metals.

F. LIMONITE ROCK.—A fissile, porous, compact or fibrous, earthy or arenaceous hydrated oxide of iron, bedded or in veins; reddish, yellowish, or blackish in colour.

Varieties are—*a. Limonitic Shale*; *b. Oolitic Brown Ore*; *c. Reniform Iron Ore*; *d. Pea-iron Ore*; *e. Bog-iron Ore*: the latter having sub-varieties, (*a.*) *Black Bog-iron Ore*; (*b.*) *Yellow Bog-iron Ore*, or *Yellow Ochre*; and (*c.*) *Sparry Bog-iron Ore*.

Limonite Rock sometimes occurs interstratified among rocks of Secondary age,* but other vari-

* In co. Clare, Ireland, it occurs in beds among Cambro-silurian rocks, while in Limerick and Queen's counties it has been found associated with black shales near the junction of the Carboniferous limestone and the lower shales of the Coal-measures. In Ulster it occurs interbedded with Whinstone of Tertiary age.

eties, such as *c*, *d*, and *e*, usually occur associated with surface accumulations, such as drift or peat. Cotta writes of *Oolitic Brown Ore*, that it "occurs in the form of layers in many formations." *Bog-iron Ore*, and its sub-varieties, are due to the decomposition of iron ores (often pyrite or marcasite), or minerals, or rocks containing a large percentage of iron. *Black Bog-iron Ore* in general is rich in manganese, while the *yellow* and *sparry* subvarieties are respectively argillaceous and siliceous.

G. HEMATITE ROCK.—A granular or compact aggregate of hematite, with arenaceous or argillaceous matter; usually reddish in colour, but sometimes blackish or rich brown.

Varieties are—*a. Red-iron Mould*, or *Red Ochre*; *b. Reddle* or *Raddle*; *c. Sparry Red-iron stone*; *d. Specular Iron*.

Of these varieties, the principal are *a*, *b*, and *c*, while *d* seems to be rather rarely a rock-mass.

e. Topanhoacanga, or *Moorshead Rock*.—This Brazil variety is found among the surface deposits, and consists of fragments of specular iron, micaceous iron, and magnetite in a ferruginous matrix. In this rock there are various accessories; even, on rare occasions, grains of native gold.

H. SPHEROSIDERITE, or SPHEROSIDERITE, Clay-iron Stone.—A crystalline aggregate of siderite combined with argillaceous and carbonaceous matter. Impure varieties are arenaceous.

Sphaerosiderite, or *Clay-iron Stone*, occurs in thin beds, layers, and nodules, in beds of clay, clunch, and shale. It is typical of the coal-measures, but

is also found associated with other rocks. An important Coal-measure variety, very rich in carbonaceous matter, is said by Bristow to be called *Black-band*. Some beds of the latter are so rich in carbon, that they can be burned by themselves, and might be classed among the coals.

I. MINERALS OCCURRING AS ROCKS.—Some minerals occur in subordinate masses interstratified with the sedimentary rocks; these, however, are not of very frequent occurrence, minerals in mass more often being found in veins, nodules, or irregular accumulations.

a. Asphalte [Gr. *asphaltos*], *Bitumen*, *Mineral Pitch*.—Colour brownish, blackish, or black; lustre like that of black pitch, odour bituminous, melts at from 90° to 100°, and burns with a bright flame.

Of this rock Dana writes: "Asphaltum belongs to rocks of no particular age; the most abundant deposits are superficial."

b. Pyrite occurs in thin beds, layers, and as nodules.

c. Cinnabar.—"Occurs in beds in slate-rock and shale, and rarely in granite or porphyry."
—*Dana*. This observer also states that this mineral occurs as rock-masses, both in unaltered and metamorphosed sedimentary rocks.

d. Boracite "occurs in beds of anhydrite, gypsum, and salt."—*Dana*.

e. Trona, "in North Africa, forming a rock which is even used for building purposes."
—*Cotta*.

f. Aragonite, with its subvariety *Flosferri*. "The

most common repositories of aragonite are beds of gypsum, and beds of iron ore (where it occurs in coralloidal forms, and is denominated *Flosferri*), basalt, and igneous rocks."—*Dana*.

- g. *Manganese Ores*, generally in the older rocks, but also in surface deposits.
- h. *Rhodochrosite* occurs generally in veins, but at Glendree, Clare, Ireland, it forms part of the surface-deposits.
- i. *Bole*, in subordinate masses, in limestone.
- j. *Opal*, or *Vitryte*, in very subordinate masses.
- k. *Fluorite*, and l. *Barite*, rarely found in beds or layers, often occur as gangue in mineral veins.
- m. *Ankerite*, sometimes as an independent rock.
- n. *Malachite*, usually in veins.
- o. *Melanite*, in small accumulations.
- p. *Galmey*, or *Calamine*, as aggregates in dolomite.
- q. *Zincite*, r. *Galenite*, s. *Stibnite*, and t. *Arsenopyrite*, rarely occur as subordinate layers, but more generally in veins.
- u. *Sulphur*, in concretions and layers, principally in marl and limestone, also in the vicinity of volcanoes.
- v. *Wavellite*, in subordinate beds or layers.
- w. *Chalcopyrite*, usually in veins.

C. Rocks partly mechanically, partly chemically, and partly organically formed.—These rocks are always more or less crystalline, and usually occur as subordinate beds, interstratified with sedimentary rocks, but not always.

A. LAMINATED OR STRATIFIED COAL.—A more or less fissile crystalline aggregate of carbonaceous

matter; arenaceous or argillaceous to a greater or less degree.

Most coals do not belong to the Subaqueous rocks; some, however, have been formed mechanically from carbonaceous matter, that was first denuded, and afterwards deposited in the sea, or other waters.

NOTE.—At the present day the accumulation of mechanically-formed coal can be observed. Wherever *peat-bogs* or even hard *coal-beds* are exposed to denudation, more especially to marine abrasion, they are gradually carried away to be deposited in seas, lakes, &c., forming stratified more or less arenaceous or argillaceous carbonaceous beds, which eventually must be indurated, forming a hard coal. Some of these deposits in the vicinity of the Irish coast are of considerable thickness.

Varieties are—*a. Cannel Coal*, a compact, semi-fissile coal; breaks with a semiconchoidal smooth fracture, with little or no lustre; colour dull black or greyish black.

Cannel Coal is bituminous and often caking. Of it Dana writes: "On distillation it affords, after drying, 40 to 66 of volatile matter, and the material volatilized includes a large proportion of burning and lubricating oils. It graduates into oil-producing coaly shales, the more compact of which it much resembles."

NOTE.—Some varieties of anthracyte have erroneously been classed as Cannel coal, on account of their smooth, close texture.

Subvarieties are—(a.) *Parrot Coal*, (b.) *Horn Coal*.

A Scotch subvariety is called *Parrot Coal*, because it burns with a crackling noise; while a Welsh is named *Horn Coal*, as it emits, when burning, an odour like that of burnt horn. An earthy subvariety is called, in Yorkshire, *Clod*.

And (c.) *Torbanyte*, *Boghead Cannel*, *Boghead Mineral* (after *Torbane Hill*, Scotland).—A dark brown variety of cannel coal; yellowish streak, without lustre, and having a sub-conchoidal fracture.

Torbanyte yields over 60 per cent. of volatile matter, and is used for the production of burning and lubricating oils, paraffin, and illuminating gas.

b. *Splint Coal*, *Splent Coal*.—A hard, laminated, bituminous coal; not easily broken or kindled, though, when lighted, it affords a clear, lasting fire.

According to Page, the name “Splint” or “Splent,” is derived from the coal splitting (or *splenting*) up “in large flaggy or board-like laminæ.”

c. *Culm*, a fissile, flaggy, shaly, flaky, or scaly variety of non-bituminous coal or *Anthracyte*.

Culm sometimes occurs in independent beds, but more often it is found associated with *Anthracyte*. The flaky or scaly nature of some culm may possibly be due to the growth and decay of organic matter in layers, but in others undoubtedly it is due to carbonaceous matter having been deposited in water.

d. *Bituminous Shale*, and e. *Carbonaceous Shale*.—Respectively bituminous or non-bituminous shaly aggregates of coal, clay, sand, and such-like; more or less crystalline; colour black, or blackish-brown.

Both the bituminous and carbonaceous shales *graduate* on the one hand into argillaceous shale, and on the other into coal. The former merges

into *Cannel*, and the latter into *Culm*. The sub-varieties a and b are bituminous, while c, d, and e are non-bituminous. Some bituminous shales are impure coals; that is, they are capable of being used as fuel, while the poorer kind will not continue to burn in the fire, but become ash, the "slate" of the coal-merchant. Burning and lubricating oils, illuminating gas, &c., can be procured from many of the *Bituminous shales*. In general the *Carbonaceous shales* are not of much value; a few, however, when mixed with anthracyte, make a strong and lasting fire.

Local Terms for Bituminous Shale.

- a. *Batt* and *Bass* (Staffordshire), (b.) *Dauks*.

Local Terms for Carbonaceous Shale.

- c. *Kelve* (Leinster), (d.) *Pindy* (Cork), (e.) *Slaty Culm* (Limerick and Clare).

NOTE.—The other coals, such as *Peat*, *Lignyte*, and *Black Coal*, are classed and described among the SUBAERIAL ROCKS (page 122); they having accumulated on the surface of the earth, prior to having been buried and covered by more recent deposits.

B. LIMESTONE.—A more or less crystalline aggregate of calcite, in association with a greater or less quantity of carbonaceous, argillaceous, and arenaceous matter; compact, porous, fissile, amorphous, or oolitic; very variable in colour.

Some limestones are made up almost entirely, others only partially, of shells, fragments of shells, corals, madrepores, and such-like; others for the most part are due to the precipitation of lime held in solution by water; while some in a great measure are mechanically formed, preëxisting calcareous

rocks having been denuded away and the detritus deposited in water. Limestones in general, however, appear to be more or less due to a combination of two or more of these modes of formation.

Limestone may contain magnesia, silica, alumina, bitumen, ores of iron, &c., either as mechanical admixtures or as chemical deposits, in conjunction with the carbonate of lime; and the presence of such minerals occasions many varieties in colour as well as composition. Of limestone Jukes writes: "Varieties of limestone occur in different localities, both geographical and geological, peculiar forms of it being often confined to particular geological formations over wide areas; so that it is much more frequently possible to say what geological formation a specimen was derived from, by the examination of its lithological characters, in the case of limestone, than in that of any other rock."

NOTE.—The colour of many, if not of most, limestones seems due, in a great measure, to the ores of iron chemically or mechanically contained therein. Dark blue or blackish limestones, when followed to a good depth, are nearly always found to be of a greenish colour. Red and variegated limestones in depth are generally grey or dove-colour, while all yellowish limestones, when in their normal state, have a greenish shade, no matter how pale.

- a. *Compact Limestone*.—Homogeneous and fine-grained; fracture sometimes uneven, but generally conchoidal, either earthy or smooth; may be argillaceous, siliceous, magnesian, or ferruginous.
- b. *Crystalline Limestone*.—An aggregate of crystals of calcite, fine or coarse, in association with argil, silica, and the like.

Typical Crystalline limestone belongs to the me-

tamorphic rock series; nevertheless, most sedimentary limestones are more or less crystalline, some being aggregates of large well-developed crystals. Jukes found the limestone forming the "great barrier coral reef," Australia, "to have a crystalline structure internally."

c. *Marble* [Lat. *marmor*].—Any limestone capable of being cut and of taking a fine polish.

Marbles may be *unicoloured*, such as pure black, white, grey, &c.; or they may be *particoloured*, deriving their colour from contained minerals or animal remains. Homogeneous rocks, both as to texture and hardness, cut more evenly and take finer polish than others, and make the better marbles.

The names by which marbles are known are very various; some are called after the localities in which they are found; others after their colour or the shells they contain, or any other peculiarity that may give a character to a rock.

d. *Oolyte*, or *Oolite* [Gr. *oon*, an egg, and *lithos*, a stone], *Roestone*.—Spheroidal concretionary structure, having the appearance of the egg or roe of a fish.

Subvarieties are—(a.) *Pisolyte*, or *Pisolite* [Lat. *pisum*, a pea], *Peastone*, which has a resemblance to an agglutination of peas; and (b.) *Dolomitic Oolyte*, there being minute spheroids of dolomite in a limy matrix.

Each concretion is formed of numerous concentric layers, being sometimes hollow at the centre, or it may inclose a minute particle of sand or other mineral substance.

NOTE.—Oolitic rocks are typical of one of the *Secondary geological groups*, which has been given the name of the *Oolitic formation*. This structure, however, occurs in other groups, but more especially in the Carboniferous limestone.

- e. *Chalk* [Lat. *calx*, lime].—Usually a white or whitish fine-grained rock, but sometimes greyish, or even red; often earthy or pulverulent, but sometimes hard and compact.

Subvarieties are—(a.) *Indurated chalk*, called *White limestone* in Ulster. Jukes thus describes it: “Considerably harder and firmer than the friable rock which is commonly known as chalk.” (b.) *Chalk-rock* (Whitaker), a siliceous variety; (c.) *Pisolitic chalk*, or *Maestricht*, which has a minutely spheroidal structure; (d.) *Glaucconitic chalk*, greenish from containing glauconite; and (e.) *Ferruginous* or *Red chalk*, coloured red by oxide of iron.

- f. *Calc-sinter*, or *Calcareous Tufa*, a loose and friable variety of calcite deposited from water charged with lime.

Subvarieties are—(a.) *Travertine*, (b.) *Stalactyte*, (c.) *Stalagmyte*, and (d.) *Rock-meal*. When compact, hard, and semi-crystalline, it is called *Travertine*; if deposited in long mammillated and pendent masses from the roof of a cave or fissure, *Stalactyte* [Gr. *stalactis*, that drops]; while the boss formed on the floor of the cave or fissure from the lime in the water which dropped, is named *Stalagma* [Gr. *stalagma*, a drop]. *Rock-meal* is a white and light calc-sinter that becomes powder on the slightest pressure.

- g. *Coral-reef Limestone*.—Limestones in course of formation, consisting of corals, shells, and

calcareous sand or mud, often more or less crystalline.

- h. *Brecciated Limestone, Limestone Breccia*, and
- i. *Limestone Conglomerate, or Puddingstone*.—A rock that has an appearance as if fragments of limestone had been dropped into a limestone matrix.

If the contained pieces are of different colours from the base, and the latter contrasts well, the result is often a handsome rock; the inliers are usually purer than the matrix, and weather more freely, giving the exposed surfaces of the rock a cellular tissue. If the inliers are round or roundish, the rock is called *Puddingstone Limestone*.

- j. *Rubbly Limestone*.—A limestone made up of round or roundish irregular lumps and aggregates of lumps of various sizes, separated from one another by shaly partings, the latter being more or less calcareous.*

Limestones may also be called *Pebbly** or *Conglomeritic, Tuffose, Shaly, Slaty, Geodic, Cellular, or Porous*, if any of these structures extend over a considerable portion of the rock and give it a marked character.

- k. *Lithographyte, Lithographic Stone* [Gr. *lithos*, a stone, and *grapho*, I write].—A very even-grained, compact, fine limestone, usually buff or drab-colour, employed in lithography.

Other varieties are, — l. *Siliceous Limestone*, m. *Cherty*, n. *Argillous*, o. *Arenaceous*, p. *Fer-*

* At Oughterard, Galway, Ireland, there are remarkable beds of *Pebbly limestone*, full of quartz fragments, generally of the size of large shot, and rarely larger than a pea.

ruginous, *q.* *Bituminous*, *r.* *Stinkstone*, or *Swinestone*, or *Fetid Limestone*.

If silica is diffused throughout the lime, the rock is *siliceous*; but if the silica is in separate nodules, concretions, or layers, the rock is *cherty*; while, if particles of sand are present, the rock is *arenaceous*. Cherty and argillaceous limestone, as also some kinds of siliceous limestone, when weathered, lose these calcareous portions, the residue being called *Rotten-stone*. Many bituminous limestones, when struck or rubbed, emit an odour of sulphuretted hydrogen gas, and these have been called by the different names of *Stink-stone*, *Swine-stone*, and *Fetid Limestone*.

Important varieties are,—*s.* *Dolomitic*, and *t.* *Hydraulic Limestone*, or *Cement-stone*.

Hydraulic limestones are more or less dolomitic; they contain from 10 to 30 per cent. of silica, alumina, and magnesia. Some contain ores of iron, which are rather prejudicial than otherwise to their economic value.

Hydraulic limestone, when burnt, yields a lime that does not slake when water is added, but forms a mortar which will set under water. According to Dana, some hydraulic limestones should be classed as *Dolomytes*.

C. DOLOMYTE, *Dolomite* (after M. Dolomieu), *Magnesian Limestone*.—An aggregate of dolomite (bitter spar), combined with some calcite; usually also with more or less iron, some silica, alumina, &c.

The mineral Dolomite contains—carbonate of lime, 54·35; carbonate of magnesia, 45·65 (Dana).

The rock Dolomite, according to Cotta, must contain 23 per cent. of carbonate of magnesia; while those calcareous rocks in which there is less than this amount should belong to the limestones.

Dolomite is often geodic and vuggy, or containing numerous drusy cavities. The geodes and the lining of the cavities are usually crystalline (*Bitter spar*), and not uncommonly contain one or more isolated crystal or crystals of quartz (*Rock crystal*).

NOTE.—Dolomite belongs to both the Ingenite and the Derivate orders of rocks; as some are due to metamorphism, while others were formed by the contemporaneous deposition of the carbonates of lime and magnesia, forming a Dolomite or Magnesian limestone.

Dolomite varies in texture. It may be, *a. Granular, b. Oolitic, c. Compact, d. Porous, e. Cellular, f. Concretionary, g. Brecciated, or h. Rubbly.* It also varies in composition, and may be, *i. Ferruginous, j. Argillous, or earthy; k. Arenaceous, or sandy; l. Siliceous, m. Cherty, and n. Tuffose;* they being analogous to the previously described varieties of limestone. It weathers into *o. Dolomitic sand.*

NOTE.—Cotta appears to be of opinion that *Dolomite* is never oolitic. This however seems erroneous, as among the Carboniferous limestone in the cos. Limerick, Mayo, &c., Ireland, subordinate beds of *Oolitic dolomite* occur. In Limerick minute spheroids of dolomite are disseminated in a calcitic matrix; while in Mayo the spheroids, as well as the base, are dolomitic or dolomite.

Class II.—SUBAËRIAL ROCKS.

[Lat. *sub*, under; *aer*, lower air].—Rocks that were formed on the surface of the earth. Most of these, however, subsequently were covered by, or buried under, newer-formed strata, which may be either subaqueous, subaërial, or igneous.

D. Mechanically, chemically, and organically-formed Rocks.—Subaërial rocks are due to mechanical, chemical, or organic action, or two or more of these combined. The major part, however, including the principal rocks of the group, are more or less due to the growth and decay of organic matter, united with chemical action.

A. COAL.—This geologically includes and signifies all rocks, recent or otherwise, that can be ignited and kept burning without the addition of any other substance.

NOTE.—The burning properties of some *coals* can be improved by the addition of other substances, as of water to anthracite; nevertheless, all rocks that are classed as coal should be capable of burning by themselves.

a. PEAT, Turf, Bog [Celtic? *peat* or *puit*, a soft miry substance].—An aggregate of organic matter more or less solidified, or indurated and mineralized; colour whitish-brown, yellowish-brown, blackish or yellowish white.

Subvarieties are—(a.) *White Turf*, (b.) *Brown Turf*, (c.) *Black*, or *Stone Turf*, (d.) *Gas* or *Candle Turf*.

Peat, or Bog, varies in appearance and texture, according to the position in which it is situated,

and the rate at which the accumulation was made. On low, flat land it grows much more rapidly than on hills or hill-slopes; consequently on the low land the peat is quite different from that on the hills, and in Ireland they are classed as *Low-bog*, or *Bawn*,* and *Mountain-bog*.

The Low-bog gives (a.) *White Turf*, (b.) *Brown Turf*, and (c.) *Black*, or *Stone Turf*; while the Mountain-bog only gives (b.) *Brown*, and (c.) *Black Turf*. On both there is an unprofitable surface or "clearing" (more or less living organic matter), which varies in thickness; on the former ranging from three to six feet in depth, while on the latter it rarely exceeds twelve inches in thickness.

Peat may be *fissile*, but more generally it is *felt-like*, the vegetable remains being interwoven, and forming a tough mass. The fissile texture of subaërial peat is due to vegetable growth;† each annual growth forming a separate layer. This is more common in some bogs than in others.

White turf is a nearly pure, if not an entirely pure, organic substance, and when burnt has little or no ash. *Brown turf* is always more or less

* *Bawn* (Anglicè *white*), so called from the white appearance the dead moss, grass, &c., have during the major part of the year, compared with the colour of the grass land and tillage in the vicinity.

† There are fissile peats in which the texture is caused by lamination, peat masses being denuded and subsequently deposited. Such peats, however, are not Subaërial, but are Subaqueous Rocks, and they are often interstratified, or mixed with clay, sand, or the like. Extreme heat or frost will break up peat, and leave it ready to be denuded and carried away by wind, or "rain and rivers," to be deposited in hollows, lakes, and such other reservoirs.

mineralized; while *Black*, or *Stone turf*, is a chemico-organic rock, often containing such minerals as pyrite, marcasite, and the like; when burnt, it leaves a considerable residue or ash; sometimes it is semi-crystalline, and often it is scarcely distinguishable from *Lignyte*.

NOTE.—The residue or ash of peat attains its maximum in the deepest portions of the bog; thus *Stone turf* will have more ash than the overlying *Brown turf*, and the latter than the *White turf*; while the subordinate parts of each have respectively more ash than the portions above them. The plants growing on the surface collect their inorganic food from the atmosphere, and after their decay, the mineral substances are being continually carried downwards by the water that percolates through the mass, the lower portions thereby becoming the most impregnated: some impurities also come up in water from springs.

Gas peat, or *Candle turf*, is not very common. It was found in a mountain bog on the island of Valencia, Ireland; was of a dirty yellowish-white colour; had the consistency of soap; when dried was very inflammable, and burned with a clear, bright, steady flame.

Peat accumulations are usually surface rocks, that is, rocks still growing at the surface, and being added to at the present day; it has, however, been found under drift and other surface deposits. *Æolian drift* may be blown over it; on a sunken coast-line it often is found under *Marine drift*; while in some places it has been found under *Boulder-clay drift*.

NOTE. — Cotta mentions peat in Germany "covered by diluvial loam." Oldham records peat "under a considerable depth of drift," near Nenagh, co. Tipperary, Ireland. Other Irish localities are the Boleynendorrish valley, near Gort, co. Galway, under 25 feet of boulder-clay drift; and Newtown, Queen's co., under a thickness of 96 feet. The latter is three

feet thick, compact, solid, semi-crystalline, very bituminous, and blazing like a candle when lighted. It might almost be classed as *lignyte*.

b. Lignyte, or Lignite, Brown Coal [Lat. *lignum*, wood]. — Fissile or compact; woody or earthy; usually brown or black; streak brown; very inflammable, burning with much smoke and smell. A non-caking coal.

a. Woody Lignyte, (b.) Compact Lignyte.

Fremy has divided lignyte into two kinds; 1st, Lignyte that still displays woody structure, and 2nd, Lignyte exhibiting the aspect and compactness of soft coal.

c. Jet [*Jayet, Gagates*, after *Gagas*, a place in Lydia, Asia Minor].—"A black variety of brown coal, compact in texture, and taking a good polish."—*Dana*.

NOTE.—Jukes considers jet as a subvariety of cannel coal, while Page supposes it "is rather a species of amber than coal."

c. Black Coal, Steinkohle, Common Coal, Pit Coal.
—A black, brownish-black, or greyish-black, compact or semi-compact mass, occasionally iridescent; lustre dull to brilliant, and either earthy, resinous, or submetallic; opaque; fracture conchoidal to uneven.

Subvarieties are—(a.) *Caking Coal*, (b.) *Non-caking Coal*, (c.) *Cherry or soft Coal*.

Mineral coal, according to Dana, should be "compact, massive, without crystalline structure or cleavage; sometimes breaking with a degree of regularity, but from a jointed rather than a cleavage

structure." Black coal may be a caking or a non-caking coal; the latter is sometimes called *Cherry* or *soft coal*. *Caking coal* softens and becomes pasty or semi-viscid in the fire, also it is inclined to form clinkers or to become welded together. *Cherry* or *soft coal* burns freely without softening, or any appearance of incipient fusion.

d. *Anthracyte*, or *Anthracite*, *Stone Coal* [Gr. *anthrax*, carbon].—Non-bituminous coal; lustre bright, often submetallic, and frequently iridescent; conchoidal, sharp-edged, shining fracture, or breaking readily into small cubical lumps.

Anthracyte appears not to be a normal rock, but to be an altered bituminous coal, from which the bituminous qualities have been extracted or expelled; sometimes it is an associate of Metamorphic rocks, but it also is found interstratified and associated with unaltered rock.

NOTE. — In Munster, Leinster, and Connaught, Ireland, anthracyte is interstratified with fossiliferous grits, shales, clunch, and fire-clay, the ordinary associates of bituminous coal, and these strata are not altered, except perhaps that they are more siliceous and harder than the ordinary coal-measure rocks in other countries. In places they are also cleaved.

A rock allied to Anthracyte is *Native coke*. It is undoubtedly an altered rock, due to the irruption of igneous rocks into beds of coal. Native coke is harder and more solid than artificial coke.

B. SURFACE DEPOSITS AND ACCUMULATIONS.—These may be peaty, clayey, sandy, gravelly, shingly, or a combination of two or more.

a. *Boulder-clay Drift*, a clayey or sandy-clayey matrix, sometimes calcareous, inclosing

more or less rounded, polished, grooved, scratched and etched blocks and fragments of one or more kinds of rocks.

Subvarieties are—(a.) *Calcareous* or *Corn-gravel*, (b.) *Argillous*, or *Till*, and (c.) *Arenaceous*.

Boulder-clay drift in general partakes more or less of the nature of the subjacent rocks, with a greater or less admixture of blocks and fragments of foreign derivation. The calcareous variety in Munster is called *Corn-gravel*, being used as a manure for corn-land, while the red argillous variety in Ulster is known as *Till*.

b. *Moraine Drift*, *Boulder Drift*, an accumulation having a more or less sandy, clayey, or gravelly matrix, containing angular or sub-angular, rarely rounded blocks and fragments of the adjacent rocks, foreign materials being in subordinate quantities.

Subvarieties are—(a.) *Calcareous*, (b.) *Argillous*, (c.) *Arenaceous*, (d.) *Rocky*.

Moraine drift when in low valleys often graduates into shingle, gravel, and sand, while on hill-slopes it may pass into *Rocky moraine drift*, full of large angular blocks, often tons in weight.

c. *Sand, gravel, and shingle*, usually more or less stratified; sometimes containing thin beds or layers of clay; at times the materials are heaped confusedly together.

Subvarieties are—(a.) *Esker* [Celtic, ridge], *Kaim* [Celtic, *Kam*, winding], *A's* or *Os Drift* (Scandinavian), (b.) *Post-drift Gravel*.

NOTE.—*Boulder-clay* and *Moraine drifts* are due to glacial or ice action, and therefore may be considered subaërial accumulations, even although some of the first may have been deposited in the sea or a lake. *Esker drift* seems to be either of the previously mentioned drifts, well washed, seemingly by marine or tidal currents. Some writers, while describing *Asar* and *Kaims*, evidently have included with them the “drumlins” or ridges of boulder-clay drift, which are quite distinct. The subvarieties (a and b), though not Subaërial Rocks, are yet described here, on account of their connection with the other *Surface deposits and accumulations*. (a) is probably of marine origin, while (b) may be marine, lacustrine, or fluviatile.

c. *Beaches*.—Sand, gravel, or shingle ; often shelly.

Beaches usually occur in more or less level terraces, on account of their marking the margins of present or ancient seas or lakes.

d. *Æolian Drift* or *Sand*, *Quartzose* or *Rabbit Sand* ; (e.) *Shell* or *Calcareous Sand*.—Fine incoherent sand : the first subvariety, when normal, is entirely composed of siliceous particles. In the neighbourhood of the sea it is more or less mixed with shells, corals, and such-like, or may graduate into *shell sand*, a calcareous accumulation of the débris of corals, shells, madrepores, and the like, sometimes containing as much as ninety per cent. of limy matter.

All these sands are also called *Blowing sands*, and are of various thicknesses up to hundreds of feet. Both *Æolian drift* and *Shell sand* may have been originally subaqueous, but their present arrangement, form, and character, are due to subaërial action. Some collections of *Æolian drift*, such as that of *the Sahara*, in Africa, seem to have been originally produced by marine action, as they occupy lately-

raised sea-basins. Other accumulations may be caused by meteoric abrasion; frost, heat, and chemical action disintegrating arenaceous rocks. But large quantities of it would appear to be glacier-formed, being due to the grinding of ice-bound rocks one against another, or the bottom and sides of valleys, forming sand or silt, that subsequently was carried down by a glacial river and deposited either in a lake or sea, at or near the mouth of a river. Such sands, when afterwards raised to form dry land, if in sheltered situations, would in time be clothed with vegetable mould; but if in exposed situations, they would be wafted hither and thither by every gust of wind, forming the ridges and hillocks known as *Sand-dunes* (Celtic, *doon*, hill).

At or near the mouth of most of the valleys in Ireland, which in prehistoric times were occupied by ice-streams, whether inland or on the seaboard, these accumulations of sand are found. In the plain of Limerick, at Killonaa, opposite the mouth of the valley called Glencoloo, these sands have been proved to be over 200 feet in thickness.

f. *Bergmehl*, *Mountain-meal*. — An exceedingly fine earth, covering extensive tracts of country in Sweden, composed of microscopic siliceous shells, said by T. Rymer Jones to be due to a lake deposition.

d. *Guano*.—Accumulations of animal excrement and remains; more or less compact; sometimes semi-crystalline; very fetid; yellow, brown, or blackish-brown in colour.

e. *Soil*, *Vegetable Mould*, earthy, sandy, or peaty. —Due to the combination of meteoric action with vegetable growth and decay, along with the work of certain animals.

Any portion of the surface of the ground, if deprived of its vegetable soil, unless subject to extremes of heat or cold, or some such obstructive action, will in time regain its coating of soil. At the first, meteoric action disintegrates and ameliorates the surface, to prepare it for the growth and decay of vegetables; after which, as pointed out by Darwin and others, the operations of earthworms, ants, and such-like, accelerate the formation of soil.

Local Names for some Wine-growing Soils.

Albarizo (Sicily), a yellowish-whitish chalk-soil, consisting of lime, argil, and oxide of iron; an absorbent, spongy substance, loose, always fresh and open, not caking. *Barro* (Sicily), sand mixed with a clayey earth and gravel. *Arena* (Sicily), sand on which the vine thrives well and produces abundantly, but the grapes make a thin wine. *Bugeo* (Sicily), a blackish aggregate of clay, vegetable earth, and gravel. *Pedra Molla* (Madeira), soft rock-*débris*, which never becomes a fine mould, but is generally in a crumbly state, like small coal; in this surface-deposit the vines are planted.

NOTE.—All rocks, whether tenacious, like granite, grit, limestone, and the like, or fragile, like drift and the other surface accumulations, weather more or less freely when exposed to the atmosphere. If circumstances allowed such weathered portions to remain, a soil of a greater or less thickness would gradually form. In many localities, however, the detritus, as fast as it is formed, is carried away either by wind or rain and rivers, while in most other places more or less is removed prior to a protecting envelope being formed by the growth and decay of vegetation; and even after the envelope is complete, some of the earth, continually being brought to the surface by earthworms and the like, will be removed by wind and water.

*f. Meteoric Drift, Clay, Sand, Gravel, and Shingle,
Meteorites.*

Under peculiar circumstances, in some localities, surface rocks that can scarcely be classed with soil, are formed by meteoric abrasion ; shaly slates, some limestones, and certain ingenite rocks weather into clays ; arenaceous and some ingenite rocks will produce sands, while certain limestones, conglomerates, breccias, and the like, weather into a coarse shingle or gravel. Meteorites in a few places, as in Greenland, form surface deposits.

NOTE.—In Brazil, Agassiz and others have noted the vast thickness of meteoric drift formed *in situ* by extreme heat ; while in Ireland, in many places on the extensive crags or flats of carboniferous limestone, a coarse shingle, more or less angular, is formed by the weathering of the rock.

g. Ice may be compact, granular, fibrous, or laminated.

Firn, or *Névé*, is the accumulation above the “snow-line,” which becomes consolidated into granular ice. *Glacier ice* is indistinctly granular, with its lamination more or less contorted or even effaced. Ice that forms in crevasses is compact, while *Cave ice* may have interstratified beds of sand in it.



ALPHABETICAL INDEX to Local, Duplicate, and other Rock names, that do not appear in the Classified List, Part I.; the Local names being printed in *Italics*. In the second column are the names of the Rocks, under which the others will be found in Parts II. and III.

Agalmatolyte	Onkosin	93
<i>Albarizo</i>	Soil	130
Amphibolic Gneiss	Hornblendic Gneiss... ..	78
Amphibolic Limestone... ..	Hemitrene	90
Amphibole Schist... ..	Hornblende Schist	87
Anamesyte	Aphanyte-Dioryte	60
Anorthosite	Noryte	59
Anorthyte	Dioryte	59
<i>Arena</i>	Soil	130
Argillous Schist	Argillyte	81
<i>A's Drift</i>	Esker Drift	127
Augitic Diabase	Pyroxenic Diabase	57
Ball Elvanyte	Pyromeride	43
Ball Porphyry	Pyromeride	43
<i>Barro</i>	Soil	130
Basalt	Aphanyte-Doleryte... ..	56
Basalt	Augyte	73
Base, or Mother-rock	Elvanyte	40
Base, or Mother-rock	Granitoid Elvanyte... ..	43
Basic Tuff	Whinstone Tuff	67
<i>Bass</i>	Batt	115
<i>Bind</i>	Shale... ..	106
<i>Binder</i>	Grit	99
Bitumen	Asphalte	111
Bituminous Marl	Oil-slate	102
Black Turf	Stone Turf	123
<i>Blowing Sand</i>	<i>Æolian Drift</i>	128

Blue Rock	Shale	106
Bog	Peat	122
<i>Boghead Cannel</i>	Torbanyte... ..	114
<i>Boghead Mineral</i>	Torbanyte	114
Boulder Drift	Moraine Drift	127
<i>Brash</i>	Sand and Gravel	95
<i>Brass Balls</i>	Grit	99
<i>Brass Binders</i>	Grit	99
Brick-stone	Lateryte	101
Brown Coal	Lignyte	125
Brown Stone	Grit	99
<i>Buddagh</i>	Fire-clay	100
<i>Bugeo</i>	Soil	130
<i>Bull's-eyes</i>	Slate... ..	106
<i>Bury</i>	Shale... ..	106
Calamine	Galmey	112
Calcareous Clay	Marl	102
Calcareous Sand	Shell Sand	128
Calcareous Schist	Spilyte	81
Calcareous Tufa	Calcsinter	118
<i>Calliard</i>	Grit	99
<i>Candle-turf</i>	Gas Turf	122
<i>Catsbrain</i>	Sandstone	98
<i>Catshead</i>	Grits... ..	99
Cellular Limestone	119
Cement-stone	Hydraulic Limestone	120
Clay-ironstone	Sphaerosiderite... ..	110
Clay-schist	Argillyte	81
Clay-slate	Slate... ..	103
Clinkstone	Phonolyte	72
<i>Clod</i>	Cannel Coal	113
<i>Coal-clay</i>	Fire-clay	100
<i>Coal-seat</i>	100
Common Coal	Black Coal	125
Common Ophyte	Ophyte	63
Compact Augyte	Basalt	74
Concretionary Mica-schist	Knotty Mica-schist	86
Cone-in-cone Marl	Tutenic Marl	103
<i>Corn Gravel</i>	Calcareous Boulder-clay	127
	Drift	127
<i>Cream</i>	Ophyte	90
<i>Crow-stone</i>	Grit	99
<i>Diallage Rock</i>	Gabbro	58
<i>zb</i>	Clay	100

Doleryte	Augyte	73
Epidote Rock	Epidosyte... ..	65
<i>Fekes</i>	Shale	106
Felstone Tuff	Felsyte Tuff	67
Felsite Rock	Felsitic Granite	30
Felsyte	Felstone	44
Felsyte	Petrosilex	45
Ferruginous Oligoclase Granite	Hematitic Oligoclase Granite	33
Ferruginous Tuff	Hematitic Tuff... ..	68
Fetid Limestone	Stinkstone	120
Flag	Flagstone	98
Flaggy Euryte	Slab Euryte	50
Flos Ferri	Aragonite	112
Flucan	Shale	106
Foliated Granite	Gneissoid Granite	34
Foundation	Gravel	95
Freestone	Sandstone	99
Friable Marl... ..	Mealy Marl	102
Friction Breccia	62
Friiled Argillyte	Folded Argillyte	82
<i>Furbogh Granite</i>	Titanitic Granite	33
<i>Galliard</i>	Grit	99
<i>Galway Granite</i>	Porphyritic Oligoclase Gra- nite	34
<i>Ganister</i>	Clunch	101
Garnetiferous Limestone	Calciiphyre	89
Geodic Limestone	119
Glassy Actinolite Rock	Actinolite Rock	62
Gneissic Granite	Gneissoid Granite	34
Gneissoid Felstone	Gneissyte	54
<i>Granåt</i>	Sandstone	99
Granitic Porphyry	Elvanyte	39
Granitone	Gabbro	58
Granular Felsyte Schist	Gneissoid Felsyte Schist... ..	85
Granular Quartzzyte	Quartz Rock	84
<i>Gravel</i>	Shale	106
Greenstone	Whinstone	55
Greenstone Tuff	Whinstone Tuff	67
Grey Gneiss	Common Gneiss	78
Greystone	Trachy-doleryte	73
<i>Hardseat</i>	Clunch	101
<i>Hazel</i>	Sandstone	99
Hone	Novaculyte	103
Hybrid Rocks	40

Hypersthenyte	Hyperyte	59
Jacotinga	Itabirite	83
Jasper	Quartz	88
Kaim Drift	Esker Drift	127
Killas	Shale	106
Labrador Rock	Noryte	58
Labradoritic Augyte	Common Augyte	74
Lagen Gneiss	Ribanded Gneiss	79
Lapis Lydius	Basanyte	109
Leaf Clay	Book Clay	102
Leaf Marl	Book Marl	103
Leinster Granite	Intrusive Granite	27
Leptynite	Felsyte Schist	85
Leucite Rock	Leucilyte	74
Limestone Breccia	Brecciated Limestone	119
Lithographic Stone	Lithographyte	119
Maestricht	Pisolitic Chalk... ..	118
Magnesia Schist	Rhætzitic Schist	87
Magnesian Limestone	Dolomite	120
Mealy Felstone	Friable Felstone	50
Metal	Shale	106
Mica-trap	48
Micaceous Diabase	Kersantite	58
Micaceous Dioryte	Kersanton	60
Micaceous Felstone	Minette	47
Micaceous Hornblende Rock	Mico-hornblende Rock	63
Micaceous Limestone	Cipollino	89
Micaceous Quartzite	Itacolumyte	82
Mineral Pitch	Asphalte	111
Moorhead Rock	Topanhoaanga	110
Moss Serpentine	Pyrallolyte	93
Mountain Meal	Bergmehl	129
Obicular Dioryte	Napoleonite	60
Oelschiefer	Oil-slate	102
Oilstone	Novaculyte	105
Oney Granite	Intrusive Oligoclase Granite	34
Ophiolyte	Ophyte	91
Orthophyre	Felstone Porphyry	51
Os, or A's Drift	Esker Drift	127
Ottrelitic Schist	Phyllitic Schist	81
Oughterard Granite	Intrusive Granite	27
Pausilippo Tufa	Pumiceous Tufa	75
Pearl-stone	Perlyte	71
one	Pisolyte	117

Pebbly Limestone	119
<i>Pedra elastica</i>	Itacolumyte	...	82
<i>Pedra molla</i>	Soil	...	130
<i>Peldon</i>	Grit	...	99
<i>Pencil</i>	Pencil-slate	...	105
Peridolyte	Eukyte	...	56
Phyllitic Argillyte	Phyllistic Schist	...	81
<i>Pinsill</i>	Pencil-slate	...	105
Pipe-clay	Potter's Clay	...	100
Pitchstone Felsyte	Felstone Glass	...	46
Pit Coal	Black Coal	...	125
<i>Plate</i>	Shale	...	106
Plutonic Breccia	Brecciated Tuff	...	69
Plutonic Conglomerate	Conglomeritic Tuff	...	69
Pontellaryte	Compact Elvanyte	...	42
Porcelain Clay	Kaolin	...	65
Porous Limestone	119
<i>Porphyrite</i>	Euryte	...	49
Porphyritic Leucilyte	Leucitophyry	...	74
Porphyry	Felstone Porphyry	...	51
<i>Post</i>	Grit	...	99
Precious Ophyte	Ophyte	...	63
Puddingstone Limestone	Brecciated Limestone	...	119
Pumiceous Sand	Pumiceous Tufa	...	75
Pyrito-carbonaceous Schist	Carbonaceous Schist	...	81
Quartz Porphyry	Elvanyte	...	39
Quartz Schist	Quartzyte	...	82
Quartziferous Felstone	Quartzitic Felstone	...	46
Quartziferous Porphyry	Elvanyte	...	39
Quartzitic Granite	Greissen	...	30
Quartzose Felstone	Quartzitic Felstone	...	46
Quartzose Sand	Æolian Drift	...	128
<i>Rabbit Sand</i>	Æolian Drift	...	128
Raddle	Keddlle	...	110
Red Chalk	Ferruginous Chalk	...	118
<i>Red Free</i>	Sandstone	...	99
Red Gneiss	Gneissyte	...	54
Red Ochre	Red Iron-mould	...	110
Ribaned Quartzitic Felstone	Striped Quartzitic Felstone	...	47
Ripidolitic Elvanyte	Chloritic Elvanyte	...	42
Ripidolitic Euryte	Chloritic Euryte	...	49
Ripidolitic Gneiss	Chloritic Gneiss	...	78
Ripidolitic Gneissyte	Chloritic Gneissyte	...	54
Ripidolitic Granite	Chloritic Granite	...	33

Epikrinitic Quartzitic Felstone	Chloritic Quartzitic Felstone	47
Epikrinitic Schist	Chloritic Schist	87
Rock	Grit	98
Rock-kind	Shale	106
Rock salt	Halvite	107
Rock-salt	Quartzite	117
Rock	Sandstone	98
Rotenstone	120
Rubite	Gravel	95
Rubite	Slate	106
Saccharite Limestone	Statuary Marble	90
Scandinavian Granite	Basic or Oligoclase Granite	32
Schistoid Felstone	Foliated Felstone	51
Schorl Schist	Tourmaline Schist	87
Scorious Leucityte	Leucityte Lava	74
Seat	Fire-clay	100
Seat-clay	Fire-clay	100
Sepiolite	Meerschaum	65
Serpentine	Ophyte	63, 91
Shaly Limestone	119
Shingle	Shale	106
Shingly Felstone	Rubby Felstone	50
Schistoid Felstone	Foliated Felstone	51
Siliceous Elvanyte	Quartzitic Elvanyte	41
Slabs	Slab Slate	98
Slaty Felstone	Cleaved Felstone	51
Slaty Limestone	119
Slig	Shale	106
Sliggeen	Shale	106
Soapstone	Steatyte	64
Soft Coal	Cherry Coal	125
Soft-seat	Fire-clay	100
Sparry Gypsum	Selenyte	108
Spathic Gypsum	Selenyte	108
Spavin	Fire-clay	101
Spheroidal Mica-schist	Curled Mica-schist	86
Splint Coal	Splint Coal	114
Spotted Quartzitic Felstone	Variolitic Quartzitic Fel-	
	stone	47
Strangel Gneiss	Fibrous Gneiss	79
Striped Felsyte	Ribbed Petrosilex	46
Stone-coal	Anthracite	126
Stone-turf	Black Turf	123
Stone-kind	Shale	106

<i>Stone-mine</i>	Grit	99
Stonefield Slate	Roofing Slate	105
<i>Swine-stone</i>	Stinkstone	120
Thickly-jointed Felstone	Rubbly Felstone	50
<i>Till</i>	Argillous Boulder - clay Drift	127
<i>Tosca</i>	Pumiceous Tufa	75
Touchstone	Basanyte	109
Trachyte Porphyry	Liperyte	70
<i>Trass</i>	Pumiceous Tufa	75
Tuffose Limestone	119
<i>Turf</i>	Peat	122
<i>Underclay</i>	Fire-clay	100
Vegetable Mould	Soil	129
Verd-antique	Ophyte	91
Vesicular Augyte	Basalt Lava	74
Vesicular Lencityte	Leucityte Lava	74
Vesicular Obsidian	Pumice	71
Vesicular Trachyte	Trachyte Lava... ..	73
Vitryte	Opal	112
Whetslate	Novaculyte	105
Whetstone	Novaculyte	105
White Limestone... ..	Indurated Chalk	118
White-stone	Felsyte Schist	85
Yellow Ochre	Yellow Bog-iron Ore	109





Interesting and Useful Works

PUBLISHED OR SOLD BY

ROBERT HARDWICKE,

192, PICCADILLY,

W.

ANY WORKS IN THIS CATALOGUE MAY BE HAD
FROM ANY BOOKSELLER.





Interesting and Useful Works

PUBLISHED OR SOLD BY

ROBERT HARDWICKE,

192, PICCADILLY,

W.

**ANY WORKS IN THIS CATALOGUE MAY BE HAD
FROM ANY BOOKSELLER.**



**For Facility of Reference the Works are arranged under
Subjects, with Index to Authors.**



CONTENTS.

	PAGE
Sowerby's English Botany.....	5-10
Works on Ferns	10-11
Works on Fungi	12-13
Works on Astronomy	20
Natural History, Popular Science, and General Science.....	9-21
Publications of the Ray Society	33
Genealogical, Heraldic, and County History	25-26
Miscellaneous Literature.....	26-30
Hardwicke's Elementary Books.....	22
Mackenzie's Educational Books	23
Medical and Surgical	31-33
Hardwicke's Science-Gossip	56-64
The Monthly Microscopical Journal	48-57
Popular Science Review Contributors and Articles, Vol. 1	41-4



INDEX TO AUTHORS.

—0—

	PAGE		PAGE
Adams, L. A., F.S.A.	43	Curry, F., M.A., F.R.S.	40
Alder and Hancock	49	Daniel, P. A.	30
Alderson, Sir James	81	Darby, Rev. W. A., F.R.A.S.	20
Allman, Prof., F.R.S.	39, 40	Darwin, C., F.R.S.	39
Annandale, Thomas, F.R.C.S.	31	Davies, Thos.	15
Ansted, Prof., F.R.S.	17, 42	Davis, Dr. J. H., F.R.C.P.	33, 34
Anstie, Dr.	43	Dawkins, W. B., F.R.S.	45
Armitage, Dr.	24	Debus, H., F.R.S.	43
Baird, Dr., F.L.S.	40	De Quatrefages, A.	18
Baker, J. G., F.L.S.	10	Dick, Dr. H.	34
Barker, Dr., S.	28	Divers, Dr.	44
Barnard, H.	24	Dougall, J. D.	27
Barrett, W. F.	44	Douglas, J. W.	39
Barwell, R., F.R.C.S.	32	Dowie, James	28
Bastian, Dr., F.R.S.	46	Dyer, W. T., B.A.	11
Bate, Spence, F.R.S.	45	Ellis, Robert, M.R.C.S.	34
Beale, Dr. Lionel, F.R.S.	44	Byton, Charlotte	16
Beckstein, Dr.	20	Fleischmann, A., M.R.C.S.	27
Belcher, Rev. H.	30	Flower, W. H., F.R.S.	40
Bennett, J. H., M.D., F.R.S.	46	Forbes, David, F.R.S.	46
Beigel, Dr. H., L.R.C.P.	32	Forbes, Prof., F.R.S.	45
Bell, Joseph, F.R.C.S.	32	Foster, Prof.	44
Blackwall, J., F.L.S.	39	Fraser, Dr. P.	34
Bond, Dr. F., F.C.S.	43	Frazer, Dr. W.	34
Boner, Charles	16, 43	Fripp, Dr. H., F.R.S.	44
Bowerbank, Dr., F.R.S., F.L.S.	39	Fry, Herbert	26
Bowring, Sir John, L.L.D.	26	Galton, J. C., M.A., F.L.S.	46
Brande, Prof., D.C.L., F.R.S.L.	31	Gangee, J.	21
Brearey, F. W.	46	Glaisher, James, F.R.S.	44
Breen, James	42	Goodfellow, Dr. S. J., F.R.C.P.	34
Briot, M. C.	24	Gore, G., F.R.S.	42
Brown, I. B., F.R.C.S.	32	Gorham, M.R.C.S.	34
Browne, Mrs.	24	Gosse, P. H., F.R.S.	42
Brown, Robert, D.C.L., F.R.S.	40	Gray, Dr. J. R., F.R.S.	45
Buckman, Prof., F.L.S., F.G.S.	18, 42	Grove, Dr. John, M.R.C.S.	34, 35
Burmeister, Prof.	39	Gulliver, Prof., F.R.S.	44
Burnaby, George J.	20	Günther, Dr., F.R.S.	39
Canton, E., F.R.C.S.	33	Habershon, Dr. S. O., F.R.C.P.	35
Carpenter, Dr., F.R.S.	39	Hanks, H., L.R.C.P., Edin.	28
Carruthers, W., F.L.S.	45	Harris, Wm.,	24
Carter, R. B., F.R.C.S.	33, 46	Henslow, Rev. G., F.L.S.	43
Chamisso, A. von	26	Hickman, W., M.B., F.R.C.S.	35
Chambers, G. F., F.R.A.S.	42	Hicks, Dr. B., F.R.S.	42
Church, H. J., F.C.S.	33	Hincks, Rev. T., M.A.	46
Church, Prof.	45	Hofmeister, Dr., F.R.S.	39
Cleland, Prof.	46	Hogg, J., F.L.S.	45, 42
Cobbold, Dr. T.S., F.R.S.	43	Hooker, Dr., F.R.S.	45
Cooke, M. C.	12, 13, 14	Hooker, Sir W. J., K.H.	10
Coote, Holmes, F.R.C.S.	33	Houghton, Rev. W., F.L.S.	42
Cormac, J. R., M.D. (Cam.) F.R.S.E.	33	Howden, P., V.S.	42
Coulson, W., F.R.C.S.	38	Hull, E., F.G.S.	42
Crookes, William, F.R.S.	42, 44	Hulme, R. T., M.R.C.S., F.L.S.	42
Cross, Dr.	28	Hunt, Robert, F.R.S.	42

INDEX.

	PAGE		PAGE
Huxley, Prof., F.R.S.	39	Samuelson, James	42
Jones, Prof. T. R., F.G.S.	43	Sawyer, James M. B.	37
Jones, Dr. W. H., M.R.C.S.	35	Shaible, Dr. C. H., Ph.D.	28
K. K. (illuminated books)	30	Schleiden, Prof.	12
Kinahan, G. H.	17	Sclater, Dr., F.R.S.	40
Langley, J. Baxter, M.R.C.S., F.L.S.	35	Scott, John	39
Lankester, Dr., F.R.S.	13, 15, 29, 30	Seemann, Dr. B., Ph.D., F.L.S., F.R.G.S.	43
Lankester, E. Ray	42	Sheen, J. B.	27
Lankester, Mrs.	5, 11, 12	Sheppard, Dr. E.	27
Lawson, Dr.	18, 29, 35, 41	Simmonds, P. L.	21
Le Hardy, Miss	27	Sims, Dr. J. Marion	37
Leighton, Rev. W. A., M.A.	40	Smith, David, M.D.	37
Lewes, G. H.	42	Smith, John, A.L.S.	11
Liebig, Baron	36	Smith, W. G., F.L.S.	13
Lord, J. K.	16	Sorby, H. C., F.R.S.	44
Lowth, G. T.	30	Sowerby, J. E.	5, 12
Lukis, Captain J. H.	27	Spear, J.	24
Mackie, S. J., F.G.S.	43	Spicer, Rev. W. W., M.A.	14
Marsh, Dr. J. C. L., M.R.C.P.	27	Steenstrup, Prof.	39
Masters, Dr. M., F.L.S.	40	Strickland, H. E., F.R.S.	...
Meyen, Prof.	40	Syme, J. T. B., F.L.S.	5
Miller, W. A., M.D., V.P.R.S.	46	Symonds, Rev. W. S.	16, 43
Milne, Alex., M.D.	36	Tate, B., F.R.G.S.	13
Milton, J. L.	18, 36	Taylor, Dr. A. S., F.R.S., F.R.C.P.	31
Mivart, Geo. F.G.S.	46	Taylor, J. E., F.G.S.	17, 19
Monckhoven, D. Van, Ph.D.	19	Teevan, W. F., F.R.C.S.	38
Morris, Dr. E., F.R.C.S.	36	Trimmen, H., M.B., F.L.S.	11
Morris, Prof., F.R.S.	47	Trousseau, Prof. A.	38
Nelson, David, M.D.	17	Turner, Mansfield	24
Norton, A. T., F.R.C.S.	36	Unwin, W. C.	46
Nunn, T. W., F.R.C.S.	37	Vincent, John	19
Ogle, W., M.D.	46	Voelcker, Prof., F.C.S.	47
Parker, W. K., F. S.	40	Waite, S. C.	21
Patterson, Robt., F.R.S.	43	Walford, Rev. E.	25
Pengelly, W., F.R.S.	47	Walker, C. V., F.R.S.	43
Proctor, R. A., F.R.A.S.	20, 44	Whitley, N., C.E.	46
Procter, Wm., M.D.	37	Williamson, Prof., F.R.S.	39
Pullar, Alfred, M.D.	36	Winn, J. M., M.D.	38
Ransonnette, Baron	28	Wood, Jno., F.R.C.S.	37
Robson, J. E.	15	Zander, Dr. A.	33
Richardson, Dr. B. W., F.R.S.	43	Zerffi, Prof.	30
Sanderson, Dr. J. B., F.R.S.	37		



A CATALOGUE OF BOOKS

PUBLISHED OR SOLD BY

ROBERT HARDWICKE,

192, PICCADILLY, LONDON, W.

Sowerby's English Botany:

Containing a Description and Life-size Drawing of every British Plant. Edited and brought up to the present standard of scientific knowledge, by T. BOSWELL SYME, LL.D., F.L.S., &c. With Popular Descriptions of the Uses, History, and Traditions of each Plant, by Mrs. LANKESTER, Author of "Wild Flowers Worth Notice," "The British Ferns," &c. The Figures by J. E. SOWERBY, JAMES SOWERBY, F.L.S., J. DE C. SOWERBY, F.L.S., and J. W. SALTER, A.L.S.

"Under the editorship of T. Boswell Syme, F.L.S., assisted by Mrs. Lankester, 'Sowerby's English Botany,' when finished, will be exhaustive of the subject, and worthy of the branch of science it illustrates..... In turning over the charmingly executed hand-coloured plates of British plants which encumber these volumes with riches, the reader cannot help being struck with the beauty of many of the humblest flowering weeds we tread on with careless step. We cannot dwell upon many of the individuals grouped in the splendid bouquet of flowers presented in these pages, and it will be sufficient to state that the work is pledged to contain a figure of every wild flower indigenous to these isles."—*The Times*.

"Will be the most complete Flora of Great Britain ever brought out. This great work will find a place wherever botanical science is cultivated, and the study of our native plants, with all their fascinating associations, held dear."—*Athenæum*.

"Nothing can exceed the beauty and accuracy of the coloured figures. They are drawn life-size—an advantage which every young amateur will recognize who has vainly puzzled over drawings in which a celandine is as big as a poppy—they are enriched with delicate delineations of fruit, petal, anther, and any organ which happens to be remarkable in its form—and not a few plates are altogether new..... A clear, bold, distinctive type enables the reader to take in at a glance the arrangement and divisions of every page. And Mrs. Lankester has added to the technical description by the editor an extremely interesting popular sketch, which follows in smaller type. The English, French, and German popular names are given, and, wherever that delicate and difficult step is at all practicable, their derivation also. Medical properties, superstitions, and fancies, and poetic tributes and illusions follow. In short, there is nothing more left to be desired."—*Guardian*.

"Without question, this is the standard work on Botany, and indispensable to every botanist..... The plates are most accurate and beautiful, and the entire work cannot be too strongly recommended to all who are interested in botany."—*Illustrated News*.

* * Subscribers to this great National Undertaking may commence at any time without buying all the back numbers at once.

Prospectuses and Specimens gratis.

Sowerby's English Botany, Volume 1 contains

Aconite	Coralwort	Ladies' Smocks	Pheasant's eyes
Alyssums	Corydalis	Larkspurs	Poppies
Anemones	Cresses	Lilies	Radishes
Awlwort	Crowfoots	Madwort	Rape
Banetary	Cuckoo flower	Marigold	Rockets
Barberry	Earthsmoke	Mayflower	Rues
Barrenwort	Figwort	Monkshood	Scurvy grasses
Bearsfoot	Flixweed	Mousetails	Spearworts
Buttercup	Fumitories	Mustards	Stocks
Cabbages	Gilliflowers	Nailwort	Toothwort
Caltrops	Headwark	Nasturtiums	Traveller's joy
Candytuft	Hellebores	Naveus	Turnips
Celandines	Horseradish	Old man's beard	Wallflowers
Charlocks	Hutchinsia	Pasque flower	Whitlow grasses
Coleseed	Jack-by-the-hedge	Peony	Woad
Colewort	Kingcup	Pepperworts	Wolfsbane
Columbine			

All the Plants ranked under the orders Ranunculaceæ, Berberidaceæ, Nymphæaceæ, Papaveraceæ, and Cruciferæ.

Sowerby's English Botany, Volume 2 contains the

Aaron's beard	Cyphel	Pansies	Spindle tree
Alder	Dyer's weed	Pearlworts	Spurreys
All-seed	Flaxes	Pinks	Stitchworts
Balsams	Fuller's herb	Purslane	Storks' bills
Bladder-nut	Heartsease	Ragged Robin	Sun dews
Brusewort	Herb Roberts	Red rot	Sun rose
Buckthorns	Holly	Rocket	Sycamore
Campions	Limes	Rock roses	Tamarisk
Carnation	Limewort	Rose of heaven	Tutsan
Catchflies	Lychnis	St. John's worts	Violets
Chickweeds	Mallows	Sandworts	Water blinks
Claytonia	Maples	Sea heath	Waterworts
Corn cockle	Mignonettes	Soapwort	Wild Williams
Crane's bills	Milkworts	Sorrels	Yarr
Cuckoo flower	Moenchia		

All the Plants ranked under the orders Resedaceæ, Cistaceæ, Violaceæ, Droseraceæ, Polygalaceæ, Frankeniaceæ, Carophyllaceæ, Portulacaceæ, Tamaricæ, Elatinaceæ, Hypericaceæ, Malvaceæ, Tiliaceæ, Linaceæ, Geraniaceæ, Ilicinæ, Celastraceæ, Rhamnaceæ, Sapindaceæ.

Sowerby's English Botany, Volume 3 contains the

Agrimonies	Burnets	Gean	May
Apples	Cherries	Gorse	Meadow-sweet
Avens	Cinquefoils	Greenweeds	Medicks
Beams	Cloudberry	Hawthorns	Medlar
Birdsfoots	Clovers	Honestalks	Mellilot
Blackberry	Cotoneaster	Ladies-finger	Mountain-ashes
Blackthorn	Dewberry	Lady's mantles	Nonsuchs
Brambles	Dropwort	Lamb-toe	Oxytropis
<i>Briars</i>	Eglantine	Liquorice	Parley pier
	Fenugreek	Lucerneæ	Peat
	Furzes	Marl-grass	Peas

Sowerby's English Botany, Volume 3 (*continued*) contains the

Plum	Saintfoin	Sweetbriars	Vetches
Queen of the meadow	Service-trees	Tares	Vetchlings
Raspberries	Shamrock	Tongue under	Waxen wood
Rest-harrows	Sibbaldia	tongue	Whins
Roebuck-berry	Silver-weed	Tormentils	Whitebeams
Roses	Spinaea	Trefoils	Whitethorns
	Strawberries		

All the Plants ranked under the orders Leguminiferæ and Rosaceæ.

Sowerby's English Botany, Volume 4 contains the

Alexanders	Earth-nuts	Isnardia	Pennywort
Angelicas	Elders	Ivy	Purslane
Anise	Eryngo	Lamb's-lettuces	Rose-root
Astrantia	Evening primroses	Linnaea	Samphires
Bald-money	Fennels	Livelong	Sanicle
Bay, rose	Golden moss	London Prides	Saxifrages
Bedstraws	Gooseberry	Loose strife	Scabiouses
Bladder-seed	Goosegrass	Lovage	Squinancywort
Bryony	Goutweed	Madders	Stonecrops
Caraways	Grass of Parnassus	Mare's tail	Stoneworts
Carrots	Grass-poly	Masterwort	Sulphur-wort
Celery	Guelder-rose	Meal tree	Teasels
Cheese-rennet	Hare's-ears	Milfolia	Thorough-wax
Chervils	Hartwort	Mistletoe	Tillæa
Cicely	Hemlocks	Moschatel	Valerians
Coriander	Herb, Bennet	Mugwort	Venus's comb
Cornel	Herb, Gerard	Navelwort	Wayfaring tree
Currants	Hog's fennels	Nightshades	Willow herbs
Danewort	Holley, sea	Orpines	Woodbines
Dogwood	Honeysuckle	Parsleys	Woodruffs
Dropworts	House-leek	Parsnips	

All the Plants ranked under the orders Lythraceæ, Onagraceæ, Cucurbitaceæ, Grossulariaceæ, Crassulaceæ, Saxifragaceæ, Umbelliferæ, Araliaceæ, Cornaceæ, Loranthaceæ, Caprifoliaceæ, Rubiaceæ, Valerianaceæ, and Dipsaceæ.

Sowerby's English Botany, Volume 5 contains the

Aster	Flecampane	Heliotrope	Salsify
Blue-bottle	Everlastings	Hemp-agrimony	Samphire
Burdocks	Feverfew	Inula	Saw-worts
Bur-Marygolds	Fleabanes	Knapweeds	Sneeze-worts
Butter-bur	Fleaworts	Lettuces	Southernwood
Cat's Ears	Galin-oga	Leopards'-banes	Sow-thistle
Chamomiles	Goats'-beards	Marigold	Spikenard
Coltsfoots	Golden-rods	Mayweeds	Star-thistles
Corn-flower	Goldylocks	Mugwort	Succorys
Cotton-weed	Groundsels	Nipple-wort	Tansy
Cudweeds	Hawk-bits	Ox-eye	Thistles
Daisy	Hawks'-beards	Ox-tongue	Wormwoods
Dandelions	Hawkweeds	Ragworts	Yarrows

All the Plants ranked under the order Compositæ.

Sowerby's English Botany, Volume 6 contains the

Andromeda	Banny	Henne-belle	Rosemary
Arbutus	Butter-and-eggs	Higtaper	Sheep's-bit
Ashes	Canterbury-bell	Jacob's ladder	Snappdragons
Azalea	Centaurys	King's-taper	Speedwells
Bartsias	Cicendias	Ling	Strangle-weed
Bearberrys	Convolvulus	Lobelias	Strawberry-tree
Belladonna	Cow wheats	Louseworts	Tea-plant
Bell-flowers	Cranberry	Menziesias	Thorn-apple
Betonys	Devale	Moneywort	Throat-wort
Bilberrys	Dodders	Monkey-flower	Toadflaxes
Bindweeds	Euphrasiae	Mother-of-thou-	Tooth-wort
Bird's-nest	Eyebrights	sands	Torch-blade
Bitter-sweet	Felwort	Mudwort	Valerian
Bleaberry	Figworts	Mulleins	Venus's looking-
Bog bean	Fleullins	Nightshades	glass
Broom-rapes	Foxglove	Pennyweed	Vervian
Brooklime	Gentians	Periwinkles	Violet
Brown-rapes	Hag-taper	Primprint	Whortleberry
Brownworts	Hare-bell	Privets	Witches' thimbles
Buck beans	Heaths	Rabbit's-mouth	Winter-greens
Bull-dogs	Heather	Rampions	Yellow-rattles
Bullock's-wort	Henbane		

All the Plants ranked under the orders Campanulaceæ, Ericaceæ, Jasmminaceæ, Apocynaceæ, Gentianaceæ, Polemoniaceæ, Convolvulaceæ, Solanaceæ, Scrophulariaceæ, Orabanchaceæ, and Verbenaceæ.

Sowerby's English Botany, Volume 7 contains the

Alkanets	Comfrees	Knawels	Pimpernels
Amaranth	Cowslips	Lavenders	Plantains
Archangel	Creeping Jenny	Loosestrifes	Primroses
Balms	Cyclamen	Lungworts	Rib-grass
Basil	Dead-nettles	Madwort	Ruptureworts
Betony	Forget-me-nots	Marjoram	Saltwort
Bladderworts	Germanders	Mints	Skull-caps
Borage	Gipsey worts	Moneywort	Sea lavenders
Brook-weed	Gromwells	Motherwort	Self-heal
Bugles	Ground ivy	Nettles	Strapwort
Buglosses	— pine	Oxlips	Thrifts
Butterworts	Hemp-nettles	Oyster-plant	Thymes
Calamints	Horehounds	Pennyroyal	Water violets
Chickweed	Hounds' tongues	Peppermint	Woundwort
Clarys	Illecebrum		

All the Plants ranked under the orders Labiatae, Boraginaceæ, Lentibulariaceæ, Primulaceæ, Plumbaginaceæ, Plantaginaceæ, Paronychiaceæ, and Amarantaceæ.

Sowerby's English Botany, Volume 8 contains the

Alder	Bistorts	Elms	Junipers
Allgocd	Box	Fir	Knotgrasses
Asarabaca	Buckthorn	Goosefoots	Mercury
Aspen	Buckwheats	Hazel	Mezereon
Beech	Chestnut	Hemp	Myrtle
Beet	Crowberry	Hop	Nettles
Rinches	Docks	Hornbeam	Oaks
orts	Dog's mercurys	Hornworts	Oraches

Sowerby's English Botany, Volume 8 (*continued*) contains the

Osiers	Poplars	Samphires	Spurges
Pellitory of the wall	Purslane	Seablites	Starworts
Pepper	Rhubarb	Sea beet	Willows
Persicarias	Sallows	Sorrels	Yew
Pine	Saltworts		

All the Plants ranked under the orders *Chenopodiaceæ*, *Polygonaceæ*, *Eleganceæ*, *Thymelaceæ*, *Santalaceæ*, *Aristolochiaceæ*, *Empetraceæ*, *Euphorbiaceæ*, *Callitrichaceæ*, *Ceratophyllaceæ*, *Urticaceæ*, *Amentifereæ*, and *Coniferæ*.

Sowerby's English Botany, Volume 9 contains the

Arrow-grasses	Epipogium	Lily of the valley	Snowdrop
Arrow-head	Flowering rush	Lloydia	Snowflakes
Asparagus	Fritillary	Martagon lilies	Solomon's seals
Asphodels	Frog-bit	Naïas	Squils
Bryony	Garlics	Narcissuses	Star of Bethels
Bur-reeds	Gladiolus	Orchises	Sweet-flag
Butchers' broom	Grass-wracks	Pondweeds	Thrumwort
Cats' tails	Helleborine	Ramsons	Trichonema
Chives	Herb-Paris	Ruppias	Tulip
Coral-root	Hyacinths	Saffrons	Tway-blades
Crocuses	Iris	Scheuchzeria	Water plantains
Cuckow-pints	Lady's slipper	Semethis	Water-soldier
Daffodils	Lady's tresses	Sisyrinchium	Water thyme
Duckweeds	Leeks	Smilicina	

All the Plants ranked under the orders *Typhaceæ*, *Araceæ*, *Lemnaceæ*, *Naiadaceæ*, *Alismaceæ*, *Hydrocharidaceæ*, *Orchidaceæ*, *Iridaceæ*, *Amaryllidaceæ*, *Diascoriaceæ*, and *Liliaceæ*.

Sowerby's English Botany, Volume 10 contains the

Pipewort	Sharp - flowered	Blismuses	Prickly Sedges
Wood Rushes	Rush	Club Rushes	Grey Sedge
Three-leaved Rush	Shining - fruited	Bull Rushes	Spiked Sedges
Clustered Rush	Rush	Cotton Grasses	Axillary Sedge
Three - flowered	Lesser - jointed	Kobresia	White Sedges
Rush	Rush	Dioicious Sedges	Haresfoot Sedge
Two - flowered	Capitate Rush	Flea Sedge	Black Sedge
Rush	Toad Rushes	Rock Sedge	Alpine Sedges
Sea Rushes	Mud Rush	Few - flowered	Hoary Sedge
Common Rush	Heath Rush	Sedge	Tufted Sedge
Soft Rush	Round - fruited	Curved Sedge	Mountain Sedges
Diffuse Rush	Rush	Marsh Sedges	Water Sedges
Hard Rush	Brown Cyperus	Brown Sedges	Heath Sedges
Northern Rush	Galingale	Sea Sedge	Mud Sedges
Thread Rush	Black Schoenus	Panicked Sedges	Fingered Sedges
Blunt - flowered	Fen Sedge	Paradoxical Sedge	Silvery Sedges
Rush	Beak Sedges	Great Sedge	And other Sedges

And all Plants ranked under the Orders *Juncaceæ* and *Cyperaceæ*

Sowerby's English Botany, Volume 11 contains the

Alpine Grasses	Cut-Grass	Lyme Grass	Quaking Grass
Barley Meadows	Darnel	Marram	Reeds
Bent Grasses	Dog's-tail Grasses	Mat Grass	Ribbon Grass
Bog Hair Grass	Fesque Grasses	Meadow Grasses	Rye Grasses
Bristle Grasses	Finger Grass	Melic Grass	Small Reeds
Brome Grasses	Fox-tail Grasses	Millet Grass	Timothy Grasses
Canary-Grass	Hair Grasses	Moor Grass	Vernal Grass
Cocks'-foot Grass	Hard Grass	Nit Grass	Wall Barley
Cord Grasses	Hares'-tail Grass	Oats	Whorl Grass
Conch Grasses	Heath Grasses	Oat Grass	Yorkshire Fog
Creeping Grasses	Holy Grass	Panic Grass	

All the Plants ranked under the Order Gramineæ.

Sowerby's English Botany.

The Prices of the Volumes are:—

	Bound cloth			Half morocco			Morocco elegant				
	£.	s.	d.	£.	s.	d.	£.	s.	d.		
Vol. 1. (Seven Parts)....	1	18	0	..	2	2	0	..	2	8	6
Vol. 2. ditto	1	18	0	..	2	2	0	..	2	8	6
Vol. 3. (Eight Parts)....	2	3	0	..	2	7	0	..	2	13	6
Vol. 4. (Nine Parts)	2	8	0	..	2	12	0	..	2	18	6
Vol. 5. (Eight Parts)....	2	3	0	..	2	7	0	..	2	13	6
Vol. 6. (Seven Parts)....	1	18	0	..	2	2	0	..	2	8	6
Vol. 7. ditto	1	18	0	..	2	2	0	..	2	8	6
Vol. 8. (Ten Parts)	2	13	0	..	2	17	0	..	3	3	6
Vol. 9. (Seven Parts)....	1	18	0	..	2	2	0	..	2	8	6
Vol. 10. ditto	1	18	0	..	2	2	0	..	2	8	6
Vol. 11. (Six Parts)	1	13	0	..	1	17	0	..	2	3	6

Or, the Eleven Volumes, £22. 8s. in cloth ; £24. 12s. in half morocco ; and £28. 3s. 6d. whole morocco.

Every Known Fern.

Synopsis Filicum ; including Osmundaceæ, Schizæaceæ, Marattiaceæ, and Ophioglossaceæ, accompanied by Figures representing the essential characteristics of each Genus. By the late Sir W. J. HOOKER, K.H., and JOHN GILBERT BAKER, F.L.S., Assistant Curator of the Kew Gardens. Price £1. 2s. 6d. plain ; £1. 8s. coloured by hand.

This is the cheapest and most complete book which describes all the Ferns throughout the world. So full and clear are its details that anybody who will take the trouble to master a few botanical terms can with certainty name any fern he may meet with in any part of the world, whether growing wild or under cultivation in garden or hothouse.

"Sir W. Hooker began collecting a Fern Herbarium in 1811. For this study an enthusiastic and illustrious botanist had all his life extraordinary advantages and positions he occupied—first as Professor of Botany at Glasgow, and then as Curator of the Kew Gardens—were made the most of, not merely by amiable character which made his pupils and friends willing assistants, but at Kew

possessed the finest herbarium of ferns, and, thanks to Mr John Smith, the best collection of ferns in cultivation in the world. The names of travellers and botanists to whom he acknowledges obligations fill a page and a half of his preface. When Sir W. Hooker died, his son, Dr. Hooker, requested Mr. J. G. Baker to carry it out, possessing as he did full access to the specimens, and being supplied with Sir William's notes and annotations. The result is the publication of such a work as the author designed—a *vade mecum* for the students, cultivators, and collectors of ferns."—*Athenæum*.

Ferns, British and Foreign.

Their History, Organography, Classification, Nomenclature, and Culture, with Directions showing which are the best adapted for the Hothouse, Greenhouse, Open Air Fernery, or Wardian Case. With an Index of Genera, Species, and Synonyms. By JOHN SMITH, A.L.S., late Curator of the Royal Gardens, Kew. With 250 Woodcuts. Crown 8vo. cloth, fully illustrated, price 6s.

Mr. Smith is acknowledged to be one of the first authorities on Ferns, having been engaged nearly half a century in arranging them at Kew.

The Fern Collector's Album.

A descriptive Folio for the reception of Natural Specimens; containing on the right-hand page a description of each Fern printed in colours, the opposite page being left blank for the collector to affix the dried specimen; forming, when filled, an elegant and complete collection of this interesting family of plants. Handsomely bound, price One Guinea, size 11 $\frac{3}{4}$ in. by 8 $\frac{1}{2}$ in. A Large Edition, size 17 $\frac{1}{2}$ in. by 11 in., without descriptive Letterpress, One Guinea.

The British Ferns.

(A Plain and Easy Account of). Together with their Classification, Arrangement of Genera, Structure, and Functions, directions for out-door and in-door Cultivation, &c. By MRS. LANKESTER. Fully illustrated, price 4s. coloured by hand; 2s. 6d. plain.

"Not only plain and easy, but elegantly illustrated."—*Athenæum*.

The Flora of Middlesex.

A Topographical and Historical Account of the Plants found in the County; with Sketches of its Physical Geography and Climate, and of the progress of Middlesex Botany during the last three Centuries. By HENRY TRIMEN, M.B. (Lond.), F.L.S., Botanical Department, British Museum, and Lecturer on Botany, St. Mary's Hospital; and WILLIAM T. THISTELTON DYER, B.A., late Junior Student, Christ's Church, Oxford; Professor of Natural History, Royal Agricultural College, Cirencester. With a Map of Botanical Districts. Crown 8vo. cloth, price 12s. 6d.

The object of this book is to give a complete and accurate catalogue of the plants which have at any time been recorded to grow in Middlesex, either as natives or in a more or less completely naturalized state; to indicate the special localities where they have been found, and to trace the history of their discovery. The existence of many of these is attested only by records in scarce or little-known books, or by the original specimens preserved in old collections.

"A carefully and conscientiously compiled volume.... The authors have done their best to render it complete and useful to botanists. The biographical and historical notes and the diligent research displayed amongst scarce publications and British Museum manuscripts, render it manifest that the authors have been engaged in a labour of love, and in a sympathetic record of the scientific diligence of not a few unremembered but most devoted botanists...."—*Athenæum*.

"So many tastes and interests are catered for in this volume, that our authors are sure to get full meed of praise and thanks from all sorts and descriptions of plant-lovers, and we too are happy to aid in the general chorus."—*Gardeners' Chronicle*.

The London Catalogue of British Plants.

Published under the direction of the London Botanical Exchange Club, adapted for marking Desiderata in Exchanges of Specimens; for an Index Catalogue to British Herbaria; for Indicating the Species of Local Districts; and for a Guide to Collectors, by showing the rarity or frequency of the several species. Second Edition, 8vo. sewed, price 6d.

Wild Flowers worth Notice.

A Selection from the British Flora of some of our Native Plants which are most attractive for their Beauty, Uses, or Associations. By Mrs. LANKESTER. Illustrated by J. E. SOWERBY. Fcp. 8vo. cloth, coloured by hand, price 4s.

"We could while away a long summer day talking of the pleasant things suggested by this little book. Although all intelligent persons cannot become botanists, not to know the wild flowers of our country is to be ignorant both of our country and ourselves. And this little book will, as a pocket companion during holiday rambles—the descriptions and plates being both good—destroy this ignorance in reference to at least a hundred plants. After mastering it, the student will be not a little astounded at his own learning, when he surveys it in the systematic chapter of contents."—*Athenæum*.

Microscopic Fungi.

(Rust, Smut, Mildew, and Mould). An Introduction to the Study of Microscopic Fungi. By M. C. COOKE, Author of "The British Fungi." Fcap. 8vo. nearly 300 coloured Figures. 2nd Edition, with Appendix of New Species, price 6s.

"There is a thoroughness about Mr. Cooke's writings which always makes his communications welcome. He is not content to gather information from cyclopædias, classify and adapt, and then give a new form to the thoughts of others. On the contrary, he strikes out a new course of study, and after a laborious use of analysis, produces an entirely original work, one on which nothing of the kind had been before attempted."—*Wesleyan Times*.

The British Fungi.

(A Plain and Easy Account of). With especial reference to the Esculent and other Economic Species. By M. C. COOKE. New and revised Edition, with Coloured Plates of 40 Species. Fcap. price 6s.

"A very readable volume upon the lowest and least generally understood race of plants. For popular purposes the book could not have been better done."—*Athenæum*.

Mushrooms and Toadstools:

How to Distinguish easily the Difference between Edible and Poisonous Fungi; with two large sheets, containing Figures of 29 Edible and 31 Poisonous Species, drawn the natural size, and coloured from living specimens. By WORTHINGTON G. SMITH, F.L.S., &c. In sheets, with book, price 6s.; on canvas in cloth case for pocket, 10s. 6d.; on canvas, with rollers and varnished, for hanging up, 10s. 6d.

"Before the appearance of Mr. Smith's book scarcely more than three or four kinds were freely accepted by prudent cooks. We have now before us the list of esculent mushrooms recommended, nay, warranted by Mr. Smith, of all of which he and his friends have freely partaken, with the most satisfactory results, and many of which are far more delicious in flavour and toothsome in substance than the meadow and horse mushrooms, generally known as the mushrooms of commerce."—*Pail Mall Gazette*.

The British Reptiles.

A Plain and Easy Account of the Lizards, Snakes, Newts, Toads, Frogs, and Tortoises indigenous to Great Britain. By M. C. COOKE, Author of "The British Fungi," &c. Fcap. 8vo. cloth, fully illustrated, 4s. plain; 6s. coloured.

British Mollusks;

Or, Slugs and Snails, Land and Fresh-water. A Plain and Easy Account of the Land and Fresh-water Mollusks of Great Britain, containing Descriptions, Figures, and a Familiar Account of the Habits of each Species. By RALPH TATE, F.R.G.S. Fcap. 8vo. cloth, fully illustrated, 4s. plain; 6s. coloured.

Schleiden's Principles of Scientific Botany.

Or, Botany as an Inductive Science. Translated by Dr. LANKESTER. Hundreds of Woodcuts, and 6 pages of Figures, beautifully engraved on Steel. Demy 8vo. price 10s. 6d.

Every Botanical Library should possess this work, as it contains the principles upon which all structural botany is based.

A Manual of Structural Botany.

By M. C. COOKE. New Edition, with Chemical Notation. Illustrated by more than 200 Woodcuts, price 1s. ; bound, 1s. 6d.

"Condensed yet clear, comprehensive but brief."—*Globe*.

"We are confidently able to recommend the little volume to public favour, its very low price bringing it within the range of all purchasers."—*Era*.

A Manual of Botanic Terms.

By M. C. COOKE. New Edition. Greatly enlarged and including the recent Teratological terms. With more than 300 illustrations. Fcp. 8vo. cloth, price 2s. 6d.

"We do not hesitate to say that by a careful use of this book a sound knowledge of the theoretical portion of botany may be obtained without tedious labour,"—*Mining Journal*.

The Collector's Handy-Book of Algæ, Diatoms, Desmids, Fungi, Lichens, Mosses, &c.

With Instructions for their Preparation and the Formation of an Herbarium. Translated and Edited by the Rev. W. W. SPIGER, M.A. With 114 Illustrations. Fcp. 8vo. cloth, price 2s. 6d.

"The publisher of this work well deserves the thanks of all amateur naturalists, and especially of those who have not an unlimited amount of money to expend on their favourite pursuits. The practical hints for collecting, mounting, and preserving, are so varied and abundant that the collector can desire no better *vade mecum*."—*Naturalists' Circular*.

Economic Products from the Vegetable Kingdom.

Arranged under their respective Natural Orders, with the Names of the Plants, and their parts used in each case. Demy 8vo. price 1s. 6d.

Collection Catalogue for Naturalists.

A Ruled Book for keeping a permanent record of Objects in any branch of Natural History, with Appendix for recording interesting particulars, and lettered pages for general Index. Strongly bound, 200 pages, 7s. 6d. ; 300 pages, 10s. ; and 2s. 6d. extra for every additional 100 pages. Working Catalogues, 1s. 6d. each.

A really good catalogue, upon a simple and inexpensive plan, has long been a desideratum with naturalists and collectors. Few will originate or prepare a catalogue for themselves, and it is a frequent matter of regret, that large and otherwise valuable collections are rendered comparatively useless for want of such a history. Again, how many naturalists pass through life, enriched with knowledge gleaned from Nature's inexhaustible store, but whose knowledge, alas! dies with them, for they leave behind them no record with their collections. This catalogue commends itself to all naturalists, whether they be ornithologists, entomologists, botanists, conchologists, or mineralogists.

In addition to the ruled and printed form of catalogue, there is a ruled and paged appendix, and an Index lettered and ruled. The appendix is intended to receive such notes and observations as are too long to be recorded in the column for remarks ; and the index at the end of the work is intended to furnish an easy reference to the contents of the Catalogue.

Botanical Labels,

(A Series of.) For labelling Herbaria adapted to the names in the London Catalogue of Plants and the Manuals of Professor Babington and Dr. Hooker, with extra labels for all new species and varieties recorded in the recent volumes of the Journal of Botany, and the Exchange Club Reports. By JOHN E. ROBSON. With Index. In all 3576 labels, like specimen. Complete, Price 5s.

VIOLACEÆ.

Viola lactea, Sm.

Cream-coloured Violet.—*Heaths, E. England.*

Loc.

TEM.

COL.

The Preparation and Mounting of Microscopic Objects.

By THOMAS DAVIES. Fcp. cloth, price 2s. 6d.

This manual comprises all the most approved methods of mounting, together with the result of the Author's experience and that of many of his friends in every department of Microscopic Manipulation; and as it is intended to assist the beginner as well as the advanced student, the very rudiments of the art have not been omitted.

"Nothing is more difficult to those who handle a microscope for the first time than to get their objects in a fit state for exhibition and preparation. They will, therefore, feel greatly indebted to Mr. Davies for a little book on 'The Preparation and Mounting of Microscopic Objects.' It is clear, full, and practical; and it soon reveals to the careful student the valuable fact that a great deal may be done with very simple appliances. We recommend it to young microscopists as a book which supplies a felt deficiency."—*Guardian*.

Half-Hours with the Microscope.

By EDWIN LANKESTER, M.D., F.R.S. Illustrated by 250 Drawings from Nature by TUFFEN WEST.

CONTENTS.

Half an hour on Structure.

Half an hour in the Garden.

Half an hour in the Country.

Half an hour at the Pond-side.

Half an hour at the Sea-side.

Half an hour In-doors.

Appendix.—The Preparation and Mounting of Objects.

Third Edition, much enlarged, with full description of the various parts of the Instrument, price 2s. 6d. plain; 4s. coloured.

"The beautiful little volume before us cannot be otherwise than welcome. It is, in fact, a very complete manual for the amateur microscopist.... The 'Half-Hours' are filled with clear and agreeable descriptions, whilst eight plates, executed with the most beautiful minuteness and sharpness, exhibit no less than 250 objects with the utmost attainable distinctness."—*Critic*.

Hardwicke's Science-Gossip:

A Monthly Medium of Interchange and Correspondence for Students and Lovers of Nature. Monthly, 4d.; Annual Volume, in cloth, price 5s. See pages 56 to 64.

At Home in the Wilderness:

Being Adventures and Experiences in Uncivilized Regions, in which it is shown where and when to encamp; how to equip and manage a train of pack mules; break, gear, and saddle wild horses; cross streams, build log shanties, trenail a raft, dig out a canoe or build it with bark or hide, manage dog sleighs, tramp on snow shoes, &c. By J. KEAST LORD, late of the British North American Boundary Commission. Crown 8vo. cloth, price 6s.

In the Plain and on the Mountain.

A Guide for Pedestrians and Mountain Tourists in the Plain and on the Mountain. By CHARLES BONER, author of "Chamois Hunting in Bavaria," "Forest Creatures," &c. With illustrations of dress requisites, &c. Fcp. 8vo. price 2s.

"A little book which compresses into a very small space a great deal of good advice."—*Pall Mall Gazette*.

"We recommend Mr. Boner's book to all travellers, either on mountain or plain."—*Athenæum*.

The Book of Knots.

Illustrated by 172 Examples, showing the manner of making every Knot, Tie, and Splice. Price 2s. 6d.

"It is an honourable characteristic of our literature that it contains numerous admirable and complete treatises on many special subjects. Mr. Robert Hardwicke, publisher of many pleasant and useful works, has sent forth a 'Book of Knots,' by Tom Bowling, illustrated with one hundred and seventy-two diagrams, showing the manner of making every knot, tie, and splice, for the moderate price of half a crown, or nearly six knots a penny."—*All the Year Round*, May 23, 1868.

Old Bones; or, Notes for Young Naturalists.

By the Rev. W. S. SYMONDS, Rector of Pendock, Author of "Stones of the Valley," &c. With References to the Typical Specimens in the British Museum. Second Edition, much improved and enlarged, fully illustrated, fcp. 8vo. price 2s. 6d.

"The plan pursued by Mr. Symonds is a very simple one. He adopts the classification of Professor Owen, and carries the young naturalist from family to family, beginning with man and ending with the lowest fishes, making his own remarks as he goes on. We recommend these notes. The volume is neatly got up, and deserves a sale amongst the class for whom it is intended."—*Athenæum*.

Notes on the Geology of North Shropshire.

By CHARLOTTE EYTON. Now ready, fcp. 8vo. cloth, price 3s. 6d.

Geological Stories;

A Series of Autobiographies in Chronological Order—Being the Autobiography of—A Piece of Granite; A Piece of Quartz; A Piece of Slate; A Piece of Limestone; A Piece of Sandstone; A Piece of Coal; A Piece of Rock-Salt; A Piece of Jet; A Piece of Chalk; A Piece of Purbeck Marble; A Lump of Clay; A Piece of Lignite; The Craggs; A Boulder; A Gravel Pit. By J. E. TAYLOR, F.G.S., Author of "Half-hours at the Seaside," &c. Large fcap. 8vo., fully illustrated, price ~~4s.~~ 4s.

The Handy-book of Rock Names.

By G. H. KINAHAN, of the Irish Geological Survey, fcap. 8vo., price 4s.

On the Being and Attributes of the Godhead,

As evidenced by Creation. By DAVID NELSON, M.D., Edinburgh, Author of "The Principles of Health and Disease," &c. Demy 8vo., price 10s. 6d.

The Applications of Geology to the Arts and Manufactures.

Six Lectures delivered before the Society of Arts. By Professor D. T. ANSTEAD, M.A., F.R.S. Fcap. 8vo. cloth, illustrated, price 4s.

I. On the Formation of Natural Soils by Derivation from Rocks, and on the Improvement of Soils by the admixture of Minerals.

II. On Natural and Artificial Springs, and on the various sources of Water Supply for Towns and Cities, in connection with the Geological Structure of the Vicinity.

III. On Mineral Materials used for the Purposes of Construction: Plastic and Incoherent Materials (Clays and Sands).

IV. On Mineral Materials (*continued*): Building Stones and Slates, and their Relative Value under given Circumstances of Exposure, and on Methods of Quarrying.

V. On Stratified Deposits of Minerals, as Coal and Iron Ore, usually obtained by Mining Operations, and on Mining Methods for such Deposits.

VI. On Metalliferous Veins for Lodes and their Contents, and on the Extraction of Metalliferous Minerals from Lodes.

"The science of geology largely engages the attention of the public; but persons are frequently deterred from the study of it by the dry and tedious style of writers dedicating their talent to its exposition. Professor Ansted has supplied a book which meets the public want.... The chapters relating to agricultural geology have an especial interest for farmers; but the whole are deeply interesting and worthy of recommendation."—*News of the World*.

"Professor Ansted takes in hand a subject of some difficulty, but of universal importance. In this he describes the different kinds of mineral veins, their contents, and the methods adopted to extract those contents, and thus fitly concludes his labours in elucidation of the practical application of geological knowledge to economic purposes, especially in connection with the arts."—*Daily News*.

Transactions of the Victoria Institute

Or, Philosophical Society of Great Britain and Ireland. Twenty parts, at 3s. 6d. to 5s. each. Lists on application.

Metamorphoses of Man and Animals.

Describing the changes which Mammals, Batrachians, Insects, Myriapods, Crustaceæ, Annelids, and Zoophytes undergo whilst in the egg; also the series of Metamorphoses which these beings are subject to in after-life. Alternate Generation, Parthenogenesis, and general Reproduction treated *in extenso*. With Notes, giving references to the works of Naturalists who have written upon the subject. By A. DE QUATREFAGES. Translated by HENRY LAWSON, M.D. Crown 8vo. cloth, price 6s.

"We have already said enough to show that the essay which Dr. Lawson has introduced to us in an English garb is one which marks a new era in the history of Embryology, and which presents to both general and scientific readers information which has been hitherto confined to the realms of dusty periodicals of all languages. 'The Metamorphoses' is a work which tends to elevate the science of Biology, and deserves the attention of all classes of cultivated readers."—*London Review*.

Science and Practice in Farm Cultivation.

By Professor BUCKMAN, F.L.S., F.G.S.

CONTENTS:

- | | |
|-----------------------------------|---------------------------------|
| 1. How to Grow Good Root Crops. | 4. How to Grow Good Corn Crops. |
| 2. How to Grow Good Grass Crops. | 5. How to Grow Good Hedges. |
| 3. How to Grow Good Clover Crops. | 6. How to Grow Good Timber. |
| | 7. How to Grow Good Orchards. |

Fully Illustrated. Complete in One Volume, cloth, 7s. 6d.

The Stream of Life on our Globe.

Its Archives, Traditions, and Laws, as revealed by Modern Discoveries in Geology and Palæontology. A Sketch in Untechnical Language of the Beginning and Growth of Life, and the Physiological Laws which govern its progress and operations. By J. L. MILTON, M.R.C.S. Second Edition, crown 8vo. cloth, pp. 624, price 6s.

CONTENTS:

- | | |
|--|---|
| The Beginning of Life. | Glance at the Laws of Life. |
| England long, long ago. | Life in the Blood. |
| The First Dwellers on Earth. | Life in the Nerves. |
| The First Builders. | Life of a Giant. |
| The First Wanderers. | Life of Men of Genius. |
| The First Colonists of Sacred History. | Influence of Smoking on Life and Race. |
| The First Language. | Life in the Stars and Planets; or, |
| The First Alphabet. | Coloured Stars and their Inhabitants, &c. &c. |
| The Battle of Life. | |

"A very agreeably-written record of some of the newest and most remarkable discoveries in geology, language, and physiology. The language is always untechnical and picturesque. It has the merit of inspiring interest in subjects often treated in a manner to repel the ordinary reader."—*Lancet*.

Country Cottages.

A series of Designs for an Improved Class of Dwellings for Agricultural Labourers. By JOHN VINCENT, Architect. With numerous Plans, Elevations, &c. New Edition, folio, fully illustrated, price 12s.

Photographic Optics :

Including the Description of Lenses and Enlarging Apparatus. By D. VAN MONCKHOVEN, Ph.D. Crown 8vo. cloth, with more than 200 illustrations, price 7s. 6d.

BOOK I.—PHOTOGRAPHIC LENSES.

Preliminary Ideas.	Spherical Aberrations.
Refraction of Light.	Description of Photographic Objectives.
Chromatics.	Employment of Photographic Objectives.
Lenses.	
Aberrations.	

BOOK II.—APPARATUS FOR ENLARGEMENTS.

On the Negative intended for Enlargement.	Setting-up Enlarging Apparatus in Winter.
Description of Apparatus for Enlargements.	Setting-up of Movable Dyalitic Apparatus.
Theory of Formation of Enlarged Image in Woodward's Apparatus.	Application of Heliostat to Enlarging Apparatus.
Imperfections of Woodward's Apparatus.	Setting-up of Heliostat with Enlarging Apparatus.
The Dyalitic Apparatus.	Parallel Solar-Light Apparatus of Bertsch.
Description and Setting up of Dyalitic Apparatus, and of all Apparatus for Enlargements.	Indirect Enlargement by the Sun or by Diffused Light.
Management of Enlarging Apparatus.	Application of Artificial Light to Enlarging Apparatus.

Half-Hours at the Seaside :

Or, Recreations with Marine Objects. By J. E. TAYLOR, F.G.S., Author of "Geological Stories," &c. Small 8vo. with about 150 Illustrations, price 4s. plain ; 6s. coloured.

CONTENTS :

Half an Hour with the Waves.	Half an Hour with Sea Anemones
Half an Hour with Preparations.	Half an Hour with Sea Mats and Squirts.
Half an Hour with Seaweeds.	Half an Hour with Sea Urchins and Starfish.
Half an Hour with Sponges.	Half an Hour with Shell Fish.
Half an Hour with Seaworms.	Half an Hour with Crustaceans.
Half an Hour with Corallines.	
Half an Hour with the Jelly Fish.	

Half-Hours with the Telescope :

Being a Popular Guide to the Use of the Telescope as a means of Amusement and Instruction. Adapted to inexpensive Instruments. By R. A. PROCTOR, B.A., F.R.A.S. Fcap. 8vo. cloth, with Illustrations on Stone and Wood, price 2s. 6d.

CONTENTS :

Half an Hour on Structure of Instrument.	Half an Hour with Boötes, Scorpio, Ophiuchus, &c.
Half an Hour with Orion, Lepus, Taurus, &c.	Half an Hour with Andromeda, Cygnus, &c.
Half an Hour with Lyra, Hercules, Corvus, Crater, &c.	Half-hours with the Planets.
	Half-hours with the Sun, Moon, &c.

"It is crammed with starry plates on wood and stone, and among the celestial phenomena described or figured, by far the larger number may be profitably examined with small telescopes, and none are beyond the range of a good three-inch achromatic. The work also treats of the construction of telescopes, the nature and use of star maps and other subjects connected with the requirements of amateurs.... The book is full of 'useful and interesting information,' and will form a valuable companion to the various admirable handbooks for which Mr. Hardwicke enjoys so good a reputation."—*Illustrated Times*.

Half-hours with the Stars:

A Plain and Easy Guide to the Knowledge of the Constellations, showing, in 12 Maps, the Position of the Principal Star-Groups Night after Night throughout the Year, with Introduction and a separate Explanation of each Map. True for every Year. By RICHARD A. PROCTOR, B.A., F.R.A.S., late Scholar of St. John's College, Cambridge, and Mathematical Scholar of King's College, London; Author of "Saturn and its System," "Half-hours with the Telescope," "The Handbook of the Stars," "Sun-views of the Earth," &c. &c. Second Edition, demy 4to. price 5s.

"Nothing so well calculated to give a rapid and thorough knowledge of the position of the stars in the firmament has ever been designed or published hitherto. Mr. Proctor's 'Half-hours with the Stars' will become a text-book in all schools, and an invaluable aid to all teachers of the young."—*Weekly Times*.

The Astronomical Observer:

A Handbook for the Observatory and the Common Telescope. By W. A. DARBY, M.A., F.R.A.S., Rector of St. Luke's, Manchester. Embraces 965 Nebulae, Clusters, and Double Stars. Royal 8vo. cloth, price 7s. 6d.

"I think the design of the work has been well carried out. The catalogue will no doubt be very acceptable to the amateur observer desirous of obtaining a knowledge of practical astronomy, and it will also be useful in the library of the regular observatory."—*From the Earl of Rosse, K.P., F.R.S., &c.*

Chamber and Cage Birds.

Their Management, Habits, Diseases, Breeding, and Methods of Teaching them. Translated from the last German Edition of Dr. Bichstein's Chamber Birds, by W. E. SHUCKARD. New Edition, by GEO. J. BARNESBY, Judge of Show Birds, Derby. Price 3s. 6d.

Science and Commerce.

Their Influence on our Manufactures. A Series of Statistical Essays and Lectures describing the Progressive Discoveries of Science, the Advance of British Commerce, and the Activity of our Principal Manufactures in the Nineteenth Century. By P. L. SIMMONDS, Editor of *The Journal of Applied Science*, Honorary and Corresponding Member of Various Foreign and Colonial Societies. 600 pp. Fcap. 8vo. price 6s.

CONTENTS.

Obligations of Commerce to Science and the Vegetable Products Imported.	Silk Trade and Manufacture :
Mineral and Animal Substances entering into Commerce.	On Silk Cultivation and Supply in India.
The Industrial and Manufacturing Uses of Shells.	Production of Silk.
The Progress of Science in the 19th Century.	Iron Manufacture.
The Cotton Manufacture :	Progress of our Mineral Industries.
Production and Consumption of Cotton.	Glass Manufacture.
Our Cotton Supplies.	Earthenware and Pottery.
The Cotton Trade.	Dyes and Colouring Stuffs :
The Woollen Manufacture :	Dyes obtained from the Animal Kingdom.
The Wools of Commerce.	Mineral Colours and Dyes.
Wool and the Woollen Trade.	Vegetable Colours and Dyes.
Our Wool Supplies.	The Manufacture of, and Trade in, Precious Metals and Fancy Articles.
Colonial Wools.	The Grocery and Allied Trades :
The British Woollen Manufactures.	The Commerce in Groceries.
A—The Worsted Manufacture.	The Leather Trade and Tanning Substances :
B—The Woollen Manufacture.	The Leather Manufacture.
C—Shoddy Fabrics.	New Paper-making Materials and Progress of the Paper Manufacture.
Chief Mineral and Vegetable Products of Commerce.	On Nuts : their Produce and Uses.

The Cattle Plague.

With Official Reports of the International Veterinary Congresses, held in Hamburg, 1863, and in Vienna, 1865. By JOHN GAMGEE, Demy 8vo. 860 pp. price 21s.

Horse Warranty.

A Plain and Comprehensive Guide to the Various Points to be noted, showing which are essential and which are unimportant. With Forms of Warranty. By PETER HOWDEN, V.S. Fcap. 8vo. price 3s. 6d.

Graceful Riding.

A Pocket Manual for Equestrians. By S. C. WAITE. With Illustrations. Fcap. 8vo. cloth, price 2s. 6d.

"In the school, on the road, on the course, or across country, this little book will be invaluable; and we heartily recommend it."—Morning Post.

One Hundred Double Acrostics.

A New Year's Gift. Edited by "Myself." 16mo. cloth, price 2s. 6d.

Hardwicke's Elementary Books. Price Twopence each.

Under the above title is presented to the public a complete Library of Elementary Works adapted for the use of the people.

Thousands of people at present skilled as handicraftsmen, and as workers in the various arts and sciences of life, plod on from day to day, with some vague notion that they can improve their own particular calling; but it generally ends in nothing, because they are ignorant of the first principles of those laws which regulate the things in which they are engaged. It is a rare occurrence to meet with workmen who know anything beyond what they picked up in their apprenticeship, or obtained by imitating others more skilled than themselves.

It will be the aim of HARDWICKE'S ELEMENTARY BOOKS to teach these first principles.

No labour or expense will be spared to make the information of a thoroughly reliable character; and where advisable, a free use of authentic illustrations will be brought to bear.

"Hardwicke's series of 'Elementary Books of Science' at present includes Optics, Hydraulics, Hydrostatics, Geography, Chemistry, Mechanics, and the parts vary in price from twopence to sixpence. Each part is a very admirable epitome of the subject it treats, and there is more reliable information in any one of these little pamphlets of a few pence than there is in many a costly volume. The woodcuts are in outline, or only slightly shaded, and their subjects are exceedingly well selected."—*London Review*.

NOW READY.**Mechanics.**

Fully illustrated by nearly 100 cuts. Two parts, 2d. each; complete, 4d.; bound in cloth, 6d.

Hydrostatics.

Fully illustrated. Complete, 2d.; in cloth, 4d.

Chemistry.

Three parts, 2d. each; complete, 6d.; cloth, 8d.

Hydraulics.

Fully illustrated. Complete, 2d.; cloth, 4d.

Optics.

Fully illustrated. Complete, 4d.; cloth, 6d.

Pneumatics.

Fully illustrated, 2d.

Other Works of a similar character are in preparation, and will shortly be announced.

Mackenzie's Educational Books.

Intended for Schools or Self-instruction.

ALREADY PUBLISHED.

Mackenzie's Tables.

Commercial, Arithmetical, Miscellaneous, and Artificers'. Calculations in Bricklaying, Carpentry, Lathing, Masonry, Paper-hanging, Paving, Painting, Plastering, Slating, Tiling, Well-sinking, Digging, &c. &c. Fractions and Decimals. Forms of Receipts and Bills. Calculations on Man, Steam, Railways, Power, Light, Wind, &c. Language and Alphabets. Calendar of the Church. Scripture Money, Principal Foreign Moneys and Measures. Geographical and Astronomical Tables, &c. &c. Complete, price 2d. ; cloth, 6d.

Murray's English Grammar.

Complete, word for word with the Shilling Editions. Price 2d. ; cloth, 4d.

Mavor's Spelling.

With numerous Cuts. Price 4d. ; or, two parts, 2d. each.

Walkinghame's Arithmetic.

Same as the Half-Crown Edition. Price 4d. ; or, two parts, 2d each ; cloth, 6d.

Short-Hand.

With Phrases and Exercises, to gain facility in the use of all the characters, by which perfection may soon be attained. Complete price 2d.

Phrenology.

Explained and Exemplified. Complete, price 2d.

Bookkeeping.

By Single Entry, with explanations of Subsidiary Books, being a useful system for the Wholesale and Retail Shopkeeper. Complete, price 2d.

Elements of Arithmetic.

From the French of M. C. BRIOT. Translated by J. SPEAR, Esq.
Crown 8vo. cloth, price 4s.

"The little book before us is a translation of a French school arithmetic, and we notice it especially in reference to the metric system, which is clearly explained in its pages, and put forward in the introduction as one of the prominent features of the work."—*Morning Star*.

Mangnall's Questions, Complete, 1s.

The Cheap Edition of this valuable School Book is now ready. It has been carefully revised and brought up to the present time. It is well printed and strongly bound.

"Published in a compact form, neatly bound, and being condensed without being abridged, comes before us in a greatly improved form. Few books contain so much information in so small a space."—*Portsmouth Guardian*.

Method for Teaching Plain Needlework in Schools.

By a LADY. Price 2s. 6d. Illustrated by Diagrams and Samplers.

This useful method is based on *steps* which are *gradual, well defined*, and clear to the *perception*. It is calculated to insure the improvement of each individual child, and while it offers the necessary instruction to the less talented pupil, it enables the more clever one to attain the highest degree of perfection.

Oral Training Lessons in Natural Science and General Knowledge.

Embracing the subjects of Astronomy, Anatomy, Physiology, Chemistry, Mathematical Geography, Natural Philosophy, the Arts, History, Development of Words, &c., intended for Teachers of Public Schools, and also for Private Instruction. By H. BARNARD, Principal of Lincoln's School, Minneapolis. Crown 8vo. price 2s.

Education and Employment of the Blind.

What it has been, is, and ought to be. By T. R. ARMITAGE, M.D. Demy 8vo. cloth, price 2s. 6d.

Institutions and Charities for the Blind

In the United Kingdom (a Guide to), together with Lists of Books and Appliances for their use; a Catalogue of Books published upon the subject of the Blind; and a List of Foreign Institutions, &c. By MANSFIELD TURNER and WILLIAM HARRIS. Demy 8vo. price 3s.

Dedicated by Express Permission to H.R.H. the PRINCE of WALES.

The County Families;

Or, Royal Manual of the Titled and Untitled Aristocracy of the Three Kingdoms. It contains a complete Peerage, Baronetage, Knightage, and Dictionary of the Landed Commoners of England, Scotland, Wales, and Ireland, and gives a brief notice of the Descent, Birth, Marriage, Education, and Appointments of each Person, his Heir Apparent or Presumptive, a Record of the Offices which he has held, together with his Town Address and Country Residences. By EDWARD WALFORD, M.A., late Scholar of Balliol College, Oxford. 1,200 pages, 11,000 families. Published Annually. Price £2. 10s.

"What would the gossips of old have given for a book which opened to them the recesses of every County Family in the Three Kingdoms—we will not say every recess, for here and there we observe what may be termed 'The Blue Beard family cupboard,' omissions which are not made evidently without very good cause! There are, for example, some awkward blanks of parentage to be found. Very many have no fathers; at all events none such as they cared to name. In some places the particulars of marriages are omitted, possibly with prudence. This work, however, will serve other purposes besides those of mere curiosity, envy, or malice. It is just the book for the lady of the house to have at hand when making up the county dinner, as it gives exactly that information which punctilious and particular people are so desirous of obtaining—the exact standing of every person in the county. To the business man, *The County Families* stands in the place of directory and biographical dictionary. The fund of information it affords respecting the upper ten thousand must give it a place in the lawyer's library; and to the money-lender, who is so interested in finding out the difference between a gentleman and a 'gent,' between heirs-at-law and younger sons, Mr. Walford has been a real benefactor. In this splendid volume he has managed to meet an universal want, one which cannot fail to be felt by the lady in her drawing-room, the peer in his library, the tradesman in his counting-house, and the gentleman in his club."—*Times*.

"It possesses advantages which no other work of the kind that we know of has offered hitherto. Containing all that is to be found in others, it furnishes information respecting families of distinction which are not to be found in the latter. It will prove to be invaluable in the library and drawing-room."—*Spectator*.

"To produce such a work in the perfection which characterizes 'County Families' must have been an almost Herculean task. It is sufficient for us to say that accuracy even in the minutest details appears to have been the aim of Mr. Walford, and the errors are so few and slight that they may readily be passed over."—*Weekly Register*.

By the same Author, published annually,

**The Shilling Peerage,
The Shilling Baronetage,
The Shilling Knightage, and
The Shilling House of Commons,**

Giving the Birth, Accession, and Marriage of each Personage his Heir Apparent or Presumptive, Family Name, Political Bias and Patronage; as also a brief Notice of the Offices which he has hitherto held, his Town Address and Country Residence

The Complete Peerage, Baronetage, Knightage, and House of Commons.

In One Volume. Half-bound, with coloured edges, for marking the different divisions. Price 5s.

How to Address Titled People.

Companion to the Writing-Desk ; or, How to Address, Begin, and End Letters to Titled and Official Personages, together with Tables of Precedence, copious List of Abbreviations, Rules for Punctuation, and other useful information. New Edition. Royal 32mo. price 1s.

"A word, and more than a word of praise is due. Full information on every subject of importance to correspondents is afforded in it. The instructions on official points will not fail to be of importance to many persons unable to obtain the proper information from even much larger works of the same kind."—*Court Journal*.

"This is one of the most useful little books we have for a long time seen."—*Eve*.

The Royal Guide to London Charities.

Showing, in Alphabetical Order, the Name, Date of Foundation, Address, Object, Annual Income, Number of People benefited, Mode of Application to and Chief Officers of every Institution in London. By HERBERT FRY. Published Annually, price 1s. 6d.

Peter Schlemihl.

From the German of ADALBERT VON CHAMISSE. Translated by Sir JOHN BOWRING, LL.D., &c. Crown 8vo. cloth, with Illustrations by George Cruikshank, price 2s. 6d. ; the Illustrations on India paper, price 5s.

Hardwicke's Shilling Handy-Book of London.

An Easy and Comprehensive Guide to Everything worth Seeing and Hearing. Royal 32mo., cloth, price 1s.

CONTENTS :

Bazaars.	Omnibuses.
Ball-rooms.	Palaces.
Cathedrals.	Parks.
Dining-rooms.	Passport Offices.
Exhibitions.	Picture Galleries' Regulations.
Mansions of Nobility.	Popular Entertainments.
Markets.	Police-courts.
Money-order Offices.	Prisons.
Monuments and Statues.	Railway Stations.
Museums.	Steamboats.
Music-halls and Concert-rooms.	Theatres.
Out-door Amusements.	Telegraph Offices, &c.

Shooting Simplified.

A Concise Treatise on Guns and Shooting. By JAMES DALZIELL DOUGALL. Second Edition, Re-written and Enlarged, with a Special Chapter on Breech-Loaders. Fcap. 8vo. cloth, price 6s.

Wines and other Fermented Liquors.

From the Earliest Ages to the Present Time. Dedicated to all Consumers in the United Kingdom. By JAMES RICHMOND SHEEN. Fcap. 8vo. cloth, price 5s.

The Common Sense of the Water Cure.

A Popular Description of Life and Treatment at a Hydropathic Establishment. By Captain J. H. LUKIS, late of the 61st Regiment, and the North Durham Militia. Crown 8vo. price 5s.

"We have seldom read a more amusing book than this. Capt. Lukis is a clever well-bred gentleman, who has found the greatest pleasure of his life in the strict discipline of hydropathy, and in this volume he chats pleasantly about his own experiences, and puts in a very popular form the chief arguments in favour of the water cure. There is not a dull page in the book."—*Morning Herald*.

Special Therapeutics.

An Investigation into the Treatment of Acute and Chronic Disease by the Application of Water, the Hot-Air Bath, and Inhalation. By J. C. LORY MARSH, M.D., M.R.C.P. Crown 8vo. price 3s. 6d.

Bathing: How to do it, When to do it, and Where to do it.

By EDGAR SHEPPARD, M.D., Medical Superintendent of the Male Department, Colney Hatch Asylum. Third Edition. 8vo. sewed, price 1s.

Plain and Practical Medical Precepts.

Second Edition, revised and much enlarged. By ALFRED FLEISCHMAN, M.R.C.S. On a large sheet, price 4d.

The Home Nurse.

A Manual for the Sick Room. By ESTHER LE HARDY. Second Edition, fcap. 8vo. cloth, price 2s. 6d.

Air.	Practical Duties.	Setons, Issues.
Cleanliness.	Moral Duties.	Blisters and Plaisters.
Dress.	The Patient.	Chamber of Death.
Diet and Cookery.	Visitors.	&c. &c.

"In our notice of the first edition we expressed our approbation of the manner in which she had performed her task; and we are gratified to see that the useful lessons she inculcated both in regard to nursing and medical attendance have met with such general approval as to require another edition of her unpretending but really valuable volume."—*Lancet*.

On Teething of Infants.

Its prevalent Errors, Neglects and Dangers; their influence on the Health, and as causes of death of Children. Including the dangers of Teething Powders, Soothing Syrups, &c. By HENRY HANKS, L.R.C.P. Edinburgh, M.R.C.S. England, &c. Illustrated by Cases. Fcap. 8vo. cloth, 3s. 6d.

The Domestic Management of Infants and Children.

In Health and Sickness. By S. BARKER, M.D., Brighton. 8vo. price 5s. Also,

The Diet of Infancy and Childhood.

By S. BARKER, M.D. Demy 8vo. sewed, price 1s.

The Gastric Regions and the Victualling Department.

By AN OLD MILITIA SURGEON. The whole outward and inward man, from the crown of his head to the corns on his little toes, all tell the sad tale of the Gastric Regions' Wrongs. Crown 8vo. cloth, price 2s. 6d.

"This is a most useful, and by no means a dull or heavy book. . . . The old Militia Surgeon gives some most useful advice, in a pleasant practical manner, respecting different varieties of food and their effects upon the system."—*Observer*

The Foot and its Covering.

With Dr. Camper's work on "The Best Form of Shoe," translated from the German. By JAMES DOWIE. New Edition. Fcap. 8vo. cloth, illustrated, price 2s. 6d.

Auvergne :

Its Thermo-Mineral Springs, Climate, and Scenery. A new Salutory Resort for Invalids. By ROBERT CROSS, M.D. With three Tinted Lithographs, price 4s.

Sketches of Ceylon.

Sketches of the Inhabitants, Animal Life, and Vegetation, in the Lowlands and High Mountains of Ceylon, as well as the Submarine Scenery near the Coast, taken in a Diving Bell. By Baron EUGENE DE RANSONNETT. With 26 large Chromo-lithographs, taken from Life by the Author. Folio, £2. 10s.

First Help in Accidents.

Being a Surgical Guide in the absence or before the arrival of Medical Assistance. For the use of the Public, especially for Members of both Military and Naval Services, Volunteers, and Travellers. By CHARLES H. SCHABLE, M.D., Ph.D., Royal Military Academy, Woolwich. Fully illustrated.

First Help in Accidents (*continued.*)

Bites.	Choking.	Hanging.
Bleeding.	Cold.	Poisoning.
Broken Bones.	Dislocations.	Scalds.
Bruises.	Drowning.	Sprains.
Burns.	Exhaustion.	Suffocation.

And other Accidents where instant aid is needful. In sup. royal 32mo. cloth, price 2s. 6d.

"A most useful and interesting little book, which is besides prettily got up, and contains some accurate and nice woodcuts. In 200 small pages we find, condensed, whatever can be done in case of accidents until the arrival of a professional helper; in fact, the work is a short but complete manual, and will prove of great service to military men, volunteers, and tourists. . . . The instructions are plain and to the point, and there is a welcome absence of technical terms."—*Spectator*.

A Manual of Popular Physiology :

Being an Attempt to Explain the Science of Life in Untechnical Language. By HENRY LAWSON, M.D., Co-Lecturer on Physiology and Histological Anatomy in St. Mary's Hospital Medical School. Fcap. 8vo. with 90 Illustrations, price 2s. 6d.

Man's Mechanism.	Digestion.	The Kidneys.
Life.	Respiration.	Nervous System.
Force.	Heat.	Organs of Sense,
Food.	The Skin.	&c. &c.

"Dr. Lawson has succeeded in rendering his manual amusing as well as instructive. All the great facts in human physiology are presented to the reader successively; and either for private reading or for classes, this manual will be found well adapted for initiating the uninformed into the mysteries of the structure and function of their own bodies."—*Athenæum*.

Dr. Lankester on Food.

A Course of Lectures Delivered at the South Kensington Museum. By E. LANKESTER, M.D., F.R.S., F.L.S. New Edition. Price 4s.

Dr. Lankester on the Uses of Animals

In Relation to the Industry of Man. By EDWIN LANKESTER, M.D., F.R.S. A Course of Six Lectures, delivered at the South Kensington Museum. Crown 8vo. pp. 350, cloth, fully illustrated, price 3s.

Silk.	Soap.	Insects.
Wool.	Waste.	Furs.
Leather.	Sponges and Corals.	Feathers, Horns, & Hair.
Bone.	Shell-fish.	Animal Perfumes.

"The information is presented in the most lucid, graceful, and entertaining manner."—*Economist*.

"Every one who peruses them will be grateful to the author. The history of those creatures whose products become through man's skill so useful to him given with such charming feeling that the interest of the reader is attracted, enchained, whether he wills or no."—*Era*.

Practical Physiology: a School Manual of Health.

Being a Practical Guide to the Means of Securing Health and Life. Intended for the use of classes and general reading. By Dr. LANKESTER, F.R.S. Fifth Edition, fcap. 8vo. illustrated, price 2s. 6d.

"It is copiously illustrated. There is not a school of any kind for males or females, rich or poor, in which the book might not be used as a text-book; indeed, it ought to be as common as an English Grammar. Few persons are capable of forming an idea of the increase of human happiness and material prosperity which would follow a more general appreciation of the laws of health."—*Lancet*.

Domestic Medicines: their Uses and Doses.

In the absence of professional assistance, with Tables of Weights and Measures; the preparation of Beverages suitable for the Sick Room. Poisons and their Antidotes. Sixth Thousand, 32mo. cloth, price 1s.

The Changed Cross.

Words by L. P. W. Illuminated by K. K. Dedicated to the Memory of those blessed ones who having, "through much tribulation," finished their course with joy, now rest from their labours; and to those also who are still running with patience the course set before them, "Looking to Jesus." Square 16mo. elegantly printed, with illuminated crosses and border lines, price 6s.

Spiritualism and Animal Magnetism.

By Professor G. G. ZERFFI, Ph.D. Second Edition. Crown 8vo. Price 1s. 6d.

"This is a very clever book. It disposes of Spiritualism pretty conclusively, and it gives expositions of Animal Magnetism in connection therewith that are, as far as they go, as convincing as such efforts well can be."—*Church Herald*.

"A few more such books as this excellent little treatise, and we shall have no more table-turning or spirit-rapping. Perhaps, before long, the force at which we now either ignorantly tremble, or with equal ignorance sneer, may be doing material work."—*Land and Water*.

"As far as Professor Zerffi renders a more rational account of spiritualism than those who are enthusiastically devoted to the study, the little book under notice is worthy attention."—*Bath Journal*.

Degrees and "Degrees;"

Or, the Traffic in Theological, Medical, and other "Diplomas" Exposed. By the Rev. HENRY BELCHER, M.A., Assistant Master in King's College School, Dedicated, by permission, to the Rev. Alfred Barry, D.D., D.C.L., Canon of Worcester, Principal of King's College, London. Demy 8vo. price 1s. 6d.

Notes on Conjectural Amendments of certain Passages in Shakespeare's Plays.

By P. A. DANIEL. Crown 8vo. price 3s. 6d.

Statesman at Home.

A Dramatic Fragment. By G. T. LOWTH. Crown 8vo. limp cloth. price 3s. 6d.

MEDICAL AND SURGICAL WORKS.

The Pocket Guide to the Pharmacopœia.

Being an Explanatory Classification of its Drugs, Preparations, and Compounds. All Essentials being comprised in a form and size adapted to the Practitioners' Note Book. To enable the busy Practitioner to do justice to his vocation, and avail himself of what no memory could otherwise retain, this little book has been compiled by one who has long felt the want of some such aid. Limp cloth for the pocket-book, 1s.

Observations on Sea-Sickness,

And on some of the Means of Preventing it. By Sir JAMES ALDERSON, M.D., D.C.L., F.R.S., Consulting Physician to St. Mary's Hospital. Crown 8vo. cloth, price 2s.

Surgical Appliances and Minor Operative Surgery.

By THOMAS ANNANDALE, F.R.C.S., Edinburgh. Fcap. 8vo. cloth, price 5s.

Abstract of Surgical Principles.

By THOMAS ANNANDALE, F.R.S. and F.R.C.S. (Edin.) price 1s. each.

- I. Inflammation, Suppuration, Abscess, Sinus and Fistula, Mortification, Ulceration, and Ulcers, price 1s.
- II. Tumours, or Morbid Growths, price 1s.
- III. Dislocations, price 1s.
- IV. Fractures, price 1s.

Chemistry.

By Professor BRANDE, D.C.L., F.R.S.L., and Professor ALFRED S. TAYLOR, M.D., F.R.S., F.R.C.P. Lond. Fcap 8vo. cloth, 900 pages, price 12s. 6d.

"For clearness of language, accuracy of description, extent of information, and freedom from the pedantry and mysticism of modern chemistry, no other text-book comes into competition with it. . . . The best guide to the study of chemistry yet given to the world."—*Lancet*.

"Conceived and worked out in the most sturdy common-sense method, this book gives, in the clearest and most summary method possible, all the facts and doctrines of chemistry, with more especial reference to the wants of the medical student."—*Medical Times*.

On the Cure of Clubfoot, without Cutting Tendons.

By RICHARD BARWELL, F.R.C.S., Surgeon and Lecturer on Anatomy, Charing Cross Hospital. With 28 Photographs. Second Edition, greatly enlarged, crown 8vo. 7s. 6d.

By the same Author,

A Treatise on Diseases of the Joints.

Illustrated by Engravings on Wood. Demy 8vo. cloth, price 12s.

Also,

Causes and Treatment of Spinal Curvature.

Crown 8vo. with Woodcuts, price 6s.

A Manual of the Operations of Surgery.

For the Use of Senior Students, House Surgeons, and Junior Practitioners. Illustrated. By JOSEPH BELL, F.R.C.S. Edinburgh, Lecturer on Surgery, Assistant-Surgeon, Clinical Ward, Royal Infirmary. Surgeon to the Eye Infirmary, Edinburgh. Fcp. 8vo. price 6s.

On Inhalation as a Means of Local Treatment of the Organs of Respiration by Atomised Fluids and Gases.

By HERMANN BEIGEL, M.D., L.R.C.P. 8vo. cloth, illustrated, price 6s.

On Surgical Diseases of Women.

By I. BAKER BROWN, F.R.C.S. (by Exam.), Surgeon to the London Surgical Home. Third Edition, revised and enlarged, demy 8vo. price 15s.

By the same Author,

On Ovarian Dropsy:

Its Nature, Diagnosis, and Treatment. The Result of Thirty Years' Experience. Post 8vo. cloth, price 5s.

Also,

On Scarlatina and its Treatment.

Second Edition, fcp. 8vo. cloth, price 2s. 6d.

The Ophthalmoscope :

Its Varieties and its Uses. Translated from the German of ZANDER by R. B. CARTER, F.R.C.S. Eng. (by Exam.). With Notes and Additions. 68 Woodcuts and 3 beautiful Coloured Chromo-lithographs. Royal 8vo., price 9s.

On the Arcus Senilis; or Fatty Degeneration of the Cornea.

By EDWIN CANTON, F.R.C.S., Surgeon to the Charing Cross Hospital. With numerous illustrations. 8vo. cloth, price 10s. 6d.

The Chemical Processes of the British Pharmacopœia,

And the Behaviour with Re-agents of their Products. By HENRY J. CHURCH, F.C.S. Fcp. 8vo. price 3s.

By the same Author,

Carbolic Acid as a Disinfectant.

And as a means of preventing the spread of the Cattle Disease. 8vo. sewed, price 1s.

Disinfection and the Prevention of Disease.

By HENRY BOLLMAN CONDY. 8vo. sewed, price 1s.

Also,

Air and Water, their Impurities and Purification.

8vo. cloth, price 1s.

On Joint Diseases.

Their Pathology, Diagnosis, and Treatment; including the nature and treatment of Deformities and Curvatures of the Spine. By the late HOLMES COOTE, F.R.C.S. 8vo. cloth, price 9s.

Varicocele.

(Clinical Lecture on) Delivered at the Lock Hospital. By WALTER COULSON, F.R.C.S., Surgeon to the Lock Hospital and St. Peter's Hospital. Crown 8vo. price 2s. 6d.

Parturition and its Difficulties.

With Clinical Illustrations and Statistics of 13,783 Deliveries. By J. HALL DAVIS, M.D., F.R.C.P., President of the Obstetrical Society. New Edition, revised and enlarged, crown 8vo. cloth price 7s. 6d.

By the same Author,

Flexions of the Uterus.

With Notes of the Ovarian and other Diseases treated in the Female Ward, Middlesex Hospital, July to July, 1863-64. Price 2s. 6d.

Gleet:

Its Pathology and Treatment. With Memoir on the Treatment of Stricture of the Urethra by Subcutaneous Division. By HENRY DICK, B.A., M.D., Surgeon to the National Orthopædic Hospital. Second Edition, with Woodcuts, price 5s. 6d.

On the Safe Abolition of Pain in Labour and Surgical

Operations by Anæsthesia with Mixed Vapours. By ROBERT ELLIS, M.R.C.S., L.S.A. Crown 8vo. cloth, price 2s. 6d.

On Penetrating Wounds of the Chest.

Founded on Actual Observations in the Camp General Hospital before Sebastopol. By PATRICK FRASER, M.D. Demy 8vo. cloth, price 5s.

Treatment of Diseases of the Skin.

By WILLIAM FRAZER, M.D. Fcp. 8vo. price 3s.

On Diseases of the Kidney, and Dropsy.

By Dr. S. J. GOODFELLOW, M.D., F.R.C.P., Senior Physician to the Middlesex Hospital and Lecturer on Medicine at the Middlesex Hospital Medical School. Crown 8vo. price 7s. 6d. With illustrations from Nature by Tuffen West.

Tooth Extraction.

A Manual on the proper mode of extracting Teeth, compiled from the latest authorities and designed for the use of Students and Junior Practitioners. With a Table showing in parallel columns, the names of all the Teeth, the instruments required for their extraction, and the most approved methods of using them. By JOHN GORHAM, M.R.C.S., Fellow of the Physical Society of Guy's Hospital. Price 1s.

On Epidemic Cholera and Diarrhœa:

Their Prevention and Treatment by Sulphur. Third Edition. By JOHN GROVE, M.D., M.R.C.S., &c. Price 1s.

By the same Author,

Epidemics Examined and Explained.

8vo. cloth, price 5s.

Dr. Grove's System of Medical Book-Keeping.

Only three books required, and no posting necessary. The complete set, suitable to carry a large practice about three years, £4. 14s. 6d. Prospectuses and Examples gratis on application.

On Diseases of the Stomach:

By S. O. HABERSHON, M.D., F.R.C.P., Physician to and Lecturer on *Materia Medica* and Therapeutics at Guy's Hospital, &c. Crown 8vo. price 5s.

An Essay on some Varieties and Effects of Cancerous Disease of Bone.

By WILLIAM HICKMAN, M.B., F.R.C.S., Surgeon to the Western General Dispensary. 8vo. illustrated, 3s. 6d.

On the Use of Artificial Teeth.

In the Prevention of Indigestion, and the Cure of Diseases due to Imperfect Nutrition. By ROBERT T. HULME, M.R.C.S.E. Crown 8vo. cloth, price 2s. 6d.

On the Management of Labour in Contracted Pelvis.

A Thesis which obtained the first prize from the Faculty of Medicine of Paris. By WILLIAM H. JONES, M.D. Paris, M.R.C.S. Eng. Demy 8vo. cloth, price 4s.

Via Medica:

A Treatise on the Laws and Customs of the Medical Profession, in relation especially to Principals and Assistants, with Suggestions and Advice to Students on Preliminary Education. By J. BAXTER LANGLEY, M.R.C.S., F.L.S. Crown 8vo. cloth, price 3s.

Sciatica, Lumbago, and Brachialgia.

Their Nature and Treatment and their immediate relief and rapid Cure by Hypodermic Injection of Morphia. By HENRY LAWSON, M.D., Assistant Physician to St. Mary's Hospital, and Lecturer on Physiology in St. Mary's Hospital Medical School. Crown 8vo. 7s. 6d.

Portrait of Baron Liebig.

From a Negative taken expressly for the Publisher. Mounted on cardboard, price 10s. 6d. ; in elegant gilt frame, price 21s.

Posological Tables :

Being a Tabular Arrangement of all the Medicines contained in the British Pharmacopœia ; with Dose, Action, and Form of Administration, containing also Appendix on Poisons. By ALEXANDER MILNE, M.D. Price 1s. 6d.

Diseases of the Skin :

(The Pathology and Treatment of). By J. L. MILTON, Senior Surgeon to St. John's Hospital for Diseases of the Skin ; corresponding Member, of the New York Dermatological Society. Demy 8vo., cloth, with photographs, price, 10s. 6d.

On Shock after Surgical Operations and Injuries.

With Special Reference to Shock caused by Railway Accidents. By EDWIN MORRIS, M.D., F.R.C.S. Oxon. Crown 8vo. cloth, price 3s. 6d.

Text-Book of Skin Diseases.

By Dr. ISIDOR NEUMANN, Lecturer on Skin Diseases in the Imperial University of Vienna. Translated from the Second German Edition by special permission of the Author. By ALFRED PULLAR, M.D. and C.M. Edin., Fellow of the Royal Medical and Chirurgical Society, Physician to the East London Hospital for Children. Translation revised by the Author. Sixty Woodcuts, illustrating recent microscopic investigations. Royal 8vo. cloth, price 12s. 6d.

Osteology for Students :

A concise Description of the Human Skeleton, adapted for the use of Students in Medicine, accompanied by an Explanatory Atlas of Plates. By ARTHUR TREHERN NORTON, Assistant-Surgeon to St. Mary's Hospital, and Assistant-Lecturer and Demonstrator of Anatomy at the Medical School. 8vo. cloth, price 7s. 6d. 2 vols. complete.

Affections of the Throat and Larynx.

The classification, description, and statistics of 150 consecutive cases occurring in the Throat Department of St. Mary's Hospital. By ARTHUR TREHERN NORTON, F.R.C.S., Assistant Surgeon and Surgeon in charge of the Throat Department of St. Mary's Hospital, Lecturer on Anatomy in the Medical School. Demy 8vo. cloth, price 4s.

The Ward Manual;

Or, Index of Surgical Disease and Injury. For the Use of Students. By T. W. NUNN, F.R.C.S., Surgeon to the Middlesex Hospital. Limp cloth, price 3s.

Hygiene of Air and Water.

Being a Popular Account of the effects of the impurities of Air and Water; their detection and the modes of remedying them. By WILLIAM PROCTER, M.D., F.C.S., Surgeon to the York Dispensary. Crown 8vo. cloth, price 2s. 6d.

Handbook of the Sphygmograph.

Being a Guide to its Use in Clinical Research. By J. BURDON SANDERSON, M.D., F.R.S., Physician to the Hospital for Consumption, Assistant-Physician to Middlesex Hospital, and Joint-Lecturer on Physiology in the Middlesex Hospital Medical College. Crown 8vo. cloth, price 3s. 6d. Illustrated.

On Rupture, Inguinal, Crural, and Umbilical.

The Anatomy, Pathology, Diagnosis, Cause and Prevention; with New Methods of effecting a Radical and Permanent Cure. Embodying the Jacksonian Prize Essay for 1861. By JOHN WOOD, F.R.C.S. Eng. (Exam.) Demonstrator of Anatomy at King's College, London; Assistant Surgeon to King's College Hospital. With numerous Illustrations by Bagg. 8vo. cloth, 12s. 6d.

Lungs and Heart.

A Guide to the Physical Diagnosis of Diseases of the Lungs and Heart; together with an introduction to the Examination of the Urine. By JAMES SAWYER, M.B., Lond., &c. Resident Physician Queen's Hospital, Birmingham. Crown 8vo., price 6s.

Preservation of Sight.

Three Lectures. By DAVID SMITH, M.D., Member of the Royal College of Surgeon of England, Extra Academical Lecturer on the Eye, Glasgow, &c. Crown 8vo. price 3s. 6d.

Clinical Notes on Uterine Surgery:

With Special Reference to the Management of the Sterile Condition. By J. MARION SIMS, M.D., late Surgeon to the Women's Hospital, New York. 8vo. fully illustrated, price £1. 1s.

On Lithotomy.

By WILLIAM F. TEEVAN, B.A., F.R.C.S., Surgeon to the West London Hospital and to St. Peter's Hospital, and Lecturer on Anatomy at Westminster Hospital. Demy 8vo. price 1s.

Trousseau's Clinical Medicine.

Translated and Edited, with Notes and Appendices, by the late P. VICTOR BAZIRE, M.D., London and Paris. Parts 2 and 3, 4s. each; Vol. I. bound in cloth, price 14s.

Vol. II., III., and IV. Translated by JOHN ROSE CORMAC, M.D., Edin., F.R.S.E., Fellow of the Royal College of Physicians, Edinburgh, formerly Lecturer in the Medical School of Edinburgh, &c. Sewed, price 12s.; in cloth, price 14s. each.

A few copies of the 2nd and 3rd parts are still on sale. The work is being now brought out by the New Sydenham Society. All who have bought the early parts as published by Mr. Hardwicke can have the completion from him through their bookseller in the ordinary way.

Spinal Debility:

Its Prevention, Pathology, and Cure, in relation to Curvatures, Paralysis, Epilepsy, and various deformities. By EDWARD W. TUSON, F.R.C.S. Demy 8vo. cloth, price 5s. illustrated.

Plain Directions for dealing with an Insane Patient.

By J. M. WINN, M.D., M.R.C.P., Senior Physician to the St. George's and St. James's Dispensary, late Medical Superintendent of Sussex House Lunatic Asylum, &c. Fcap. 8vo. cloth, price 1s.

PUBLICATIONS of the ROYAL COLLEGE OF SURGEONS.

List on Application.

PUBLICATIONS OF THE RAY SOCIETY.

British Hemiptera-Heteroptera.

By J. W. DOUGLAS and JOHN SCOTT, £1. 10s.

Cirripedia Family.

By C. DARWIN, Esq. 2 vols. £2. 2s.

British Freshwater Polyzoa.

By Professor ALLMAN. £1. 11s. 6d.

Recent Foraminifera.

By Professor WILLIAMSON. £1. 11s. 6d.

Oceanic Hydrozoa.

By Professor HUXLEY. £1. 11s. 6d.

Organization of Trilobites.

From the German of BURMEISTER. 15s.

British Naked-eyed Pulmograde Medusæ.

By Professor E. FORBES. £1. 1s.

The Spiders of Great Britain and Ireland.

By J. BLACKWALL, F.L.S. 2 vols. £3. 13s. 6d.

Foraminifera.

By Dr. CARPENTER. £1. 11s. 6d.

Fructification of Higher Cryptogamia and Coniferæ.

From the German of HOFMEISTER. £1. 5s. 6d.

Reptiles of India.

By Dr. GÜNTHER. £2. 2s.

British Spongiadæ.

By Dr. BOWERBANK. 2 vols. £2. 11s.

Steenstrup on Alternation of Generations.

15s.

British Entomostracous Crustacea.

By Dr. BAIRD. £1. 1s.

British Angiocarpous Lichens.

By Rev. W. A. LEIGHTON. 10s. 6d.

British Nudibranchiate Mollusca.A Monograph of the (with coloured drawings of every species).
By Messrs. ALDER and HANCOCK.

Part	I.	Imp.	4to.	£1	10	0
"	II.	"	"	1	10	0
"	III.	"	"	1	10	0
"	IV.	"	"	1	10	0
"	V.	"	"	1	10	0
"	VI.	"	"	1	1	0
"	VII.	"	"	1	1	0

Meyen's Geography of Plants.Translated from the German by Miss MARGARET JOHNSTON. 8vo.
pp. 422, 10s. 6d.**The Miscellaneous Works of Robert Brown.**

2 vols. demy 8vo. Vol. 1. I. Geographico-Botanical; II. Structural and Physiological Memoirs, price £1. Vol. 2. Systematic Memoirs and Contributions to Systematic Works. £1. Atlas of Plates, imp. 4to. price £1.

Vegetable Teratology.

An Account of the Principal Deviations from the usual construction of Plants. By MAXWELL T. MASTERS, M.D. F.L.S. With numerous illustrations, by E. M. WILLIAMS. Demy 8vo. cloth, £1. 1s.

On Pterylography.

Translated from the German. Edited by PHILIP LUTLEY SCLATER, M.A. Ph.D. F.R.S. Imp. 4to. 10 plates, 16s.

Recent Memoirs on the Cetacea.

By Professors ESCHRICHT, REINHARDT, and LILLJEBORG. Edited by W. H. Flower, F.R.S., &c. Imp. 4to. pp. 312, 6 plates, £1. 4s.

The Shoulder Girdle and the Sternum.

By W. K. PARKER, Esq., F.R.S. Imp. 4to. illustrated with 30 coloured plates, £1. 11s. 6d.

Hydroids.

A Monograph of the Gymnoblatic or Tubularian Hydroids. By GEO. JAMES ALLMAN, M.D., Dublin and Oxon. 2 volumes, with coloured plates. Imp. 4to. £3. 3s. 0d.

POPULAR SCIENCE REVIEW.

EDITED BY HENRY LAWSON, M.D.

THE *Popular Science Review* is the only Journal which takes up the intermediate position between the world of Science and the public generally. There are scientific journals abundant which represent each section of the scientific public properly so called; and there are literary journals which devote a certain portion of their space to scientific matters of general interest. But the only journal which is at once a sort of connecting link between the scientific world on the one side and the people—i.e. those who are beyond the limit of the scientific circle—on the other, is the *Popular Science Review*.

Thus it contains no original theories. Until a theory, in fact, has obtained the sanction of the purely scientific world, it holds no place in the pages of the *Popular Science Review*; but as soon as a fact is recognized as an unquestionable scientific truth, and long before it reaches the general public by any other channel, it takes its place in the pages of the *Popular Science Review*. Here it is laid out in the plainest but most exact language possible, and, if necessary, is illustrated fully,—not in that claptrap style which is adopted by magazines of a particular class, but in a clear, distinct, and yet modest fashion, plainly setting forth the aims and desires of the author.

The plan of the *Popular Science Review* is briefly as follows:—First come a number of articles illustrated by a variety of page-plates, and dealing with those questions in each branch of science which have been most recently the subject of communications to the Scientific Societies. These are invariably done by men who are at once the most eminent and representative in the several branches of science to which they belong: we refer to the names of Contributors as a proof of this.

Leaving the Original Articles, we come next to the Reviews. These are short, but as a rule they are to the point; and they generally contain notices of every scientific book which has been published during the quarter.

Finally, we come to the Scientific Summary. In this we find a very perfect record of the principal work done at home and abroad in the following various branches:—Archæology, Botany, Geology, Palæontology, Chemistry, Mineralogy, Microscopy, Metallurgy, Mining, Physics, Photography, Zoology, Comparative Anatomy, and Medicine. This department is divided amongst a number of men, and generally it may be said to put the general reader *au courant* with the progress of science at home and abroad during the quarter which has elapsed previous to publication.

Thus, altogether, the Journal furnishes an amount of scientific information, in a popular and yet exact form, which cannot be found in any other English periodical.

Vols. I. to XI., Sets in Numbers, £5. 15s. 11 vols. cloth, £8. 14s.
Half-morocco, £8. 8s. Cloth Covers for Binding, price 1s. 6d. each.

Popular Science Review, Vol. 1 contains:—

Corn. By Prof. J. Buckman, F.L.S., F.G.S., F.S.A., &c. Illustrated.
 The Daisy. By Mrs. Lankester. Illustrated.
 The Crown Animalcule. By P. H. Gosse, F.R.S. Illustrated.
 The Lowest Forms of Life. By James Samuelson and Dr. Bruxton Hicks, F.L.S.
 Iron and Steel. By R. Hunt, F.R.S.
 Artificial Light. By Prof. Ansted, F.R.S.
 The Breath of Life. By W. Crookes, F.R.S.
 The West Coast of Equatorial Africa. With a Coloured Map.
 The Great Comet of 1861. By J. Breen. Illustrated.
 Caverns and their Contents. By Prof. Ansted, F.R.S.
 The Lowest Forms of Life. Illustrated.
 The Flower Animalcules. By P. H. Gosse, F.R.S. Illustrated.
 Cotton. By Dr. Lankester, F.R.S. Illustrated.
 Grass. By Prof. Buckman, F.L.S. Illustrated.
 The Reflex Theory. By G. H. Lewes.
 Solar Chemistry. By R. Hunt, F.R.S. With Coloured Diagram.
 Optical Phenomena of the Atmosphere. By G. F. Chambers.
 The Phosphorescence of the Sea. With a Plate.

The Sun and Solar Phenomena. With a Coloured Plate. By James Breen, F.R.A.S.
 Light and Colour. With a Coloured Plate. By Robert Hunt, F.R.S.
 The Application of Science to Electro-plating. By G. Gore, F.R.S.
 Artificial Precious Stones. By W. S. Howgrave.
 The White Clover. By Mrs. Lankester. With two Plates.
 The Human Heart. By Isaac Ashe, B.A., T.C.D.
 The Great Exhibition of 1862.
 The Britannia and Conway Tubular Bridges. With Page Plate. By W. C. Unwin, B.Sc.
 Primitive Astronomy. With two Coloured Illustrations.
 The Physics of a Sunbeam. With Coloured Plate. By R. Hunt, F.R.S.
 The English California. By G. P. Bevan, F.G.S.
 The Contents of Caverns (concluding Part). By D. T. Ansted, F.R.S.
 The Microscope, with Directions for its Use. Illustrated with Woodcuts. By C. Collingwood, M.B., F.L.S.
 The Builder Animalcules. With a Page Plate. By P. H. Gosse, F.R.S.
 The Common Truffle. With a Page Plate. By Jabez Hogg, M.R.C.S., &c.

Popular Science Review, Vol. 2 contains:—

The British Oak. Two Page Illustrations. By Prof. Buckman, F.L.S., F.G.S., &c.
 Tubifex Rivulorum, the Red Worm of our Rivers. With a Coloured Plate. By Edwin Ray Lankester.
 Anesthetics. By Dr. T. L. Phipson, M.B., F.C.S., &c.
 Barmouth and its Scientific Attractions. With a Tinted Illustration.
 The Electro-plating Process (concluding Part). With an Illustration. By George Gore.
 Notes of the Exhibition of 1862.
 The Mistletoe and Parasitic Plants. By Mrs. Lankester. Illustrated.
 The Winter Life of Plants. By Harland Coultas.
 The Vinegar Eel. By Jabez Hogg, F.L.S. With Tinted Plate.
 The Eye of the Ox and its Microscopical Structure. By E. Beckitt Truman. With Tinted Plate.

Mars. By James Breen, F.R.A.S. With Coloured Plate.
 Our Fresh-water Polyzoa. By Rev. W. Houghton, M.A., F.L.S. Two Page Plates.
 The African Lion in its Native Haunts. By Jules Gerard.
 The Human Skin. By Isaac Ashe, M.B. With Page Plate.
 The Furniture Woods of Commerce. By P. L. Simmonds.
 Fossil Birds. By S. J. Mackie, F.G.S. Illustrated.
 History of a Beech-tree. By Harland Coultas.
 The Antiquity of Man. By J. R. Leitch, M.A.
 On the Physical Geography of the Ionian Islands. By Prof. D. T. Ansted, M.A., F.R.S.
 The Telescope. By James Breen, F.R.S.
 Our British Jungfermannias.

Rotifera or Wheel Animalcules. Part IV. The Flexible Creepers (Notommatina). By Philip Henry Gosse, F.R.S. Illustrated.

On the Morphology of *Vinca minor* (small Periwinkle). By James Buckman, F.L.S., F.G.S.

Colour Blindness. By Jabez Hogg, F.L.S. Illustrated.

Popular Science Review, Vol. 3 contains :—

Photographic Printing and Engraving, with a Photograph of a Complete Page of the *Times* Newspaper printed from Stone. By William Crookes, F.R.S.

Fresh Air. By E. Lankester, M.D., F.R.S.

Microscopic Fungi Parasitic on Living Plants. By M. C. Cooke. Coloured Illustrations.

On the Physical Geography of the Ionian Islands. By Prof. D. T. Ansted, M.A., F.R.S.

The Metropolitan Main Drainage Works, with a Map of the Works. By S. J. Mackie, F.G.S.

Diseased Pork, and Microscopic Worms in Man. By John Gamgee. Illustrated.

Bodily Work and Waste. By Francis T. Bond, M.D., B.A. (Lond.), F.C.S.

The Railway Tunnel through the Alps. With Tinted Illustration.

Greek Fire: its Ancient and Modern History. By B. W. Richardson, M.A., M.D.

Notes on Earthquakes. By Rev. W. S. Symonds.

On Printing Telegraphs. With Coloured Fac-simile Illustration. By R. S. Culley.

Herrings and Herring Fishing. By the Editor.

On Proper Clothing. By E. Lankester, M.D., F.R.S.

On the Absorption and Radiation of Heat. With two Page Illustrations. By H. Debus, Ph.D., F.R.S.

Botanical Exercises. By Rev. G. Henslow, M.A., F.L.S.

Pre-historic Dwellings. With Tinted Illustrations. By G. E. Roberts, F.A.S.L.

The Aniline Dyes. With Page of Silk Illustrations. By Dr. T. L. Phipson, F.C.S.

On the Action of Manures. By Baron Liebig.

The Old Red Sandstone Fishes of England. By E. Ray Lankester. Illustrated.

Oysters and Oyster Culture. By the Editor. Illustrated.

The Pneumatic Dispatch. By S. J. Mackie, F.G.S. Illustrated.

Thermometry. By G. F. Chambers.

Popular Science Review, Vol. 4 contains :—

Extraordinary Ships. By S. J. Mackie, F.G.S. Illustrated.

Good Food. By Edwin Lankester, M.D., F.R.S.

On the Origin of Local Floras of Great Britain and Ireland. By Harland Coultas. Illustrated.

Metamorphism. By Prof. D. T. Ansted, M.A., F.R.S., &c.

Morphological Peculiarities of *Linaria spuria*. By E. S. Magrath. Illustrated.

The Metric System. By James Spear. Illustrated.

Recent Investigations into the Natural History of the Red Coral. By the Editor. Illustrated.

What is a Stimulant? By Francis E. Anstie, M.D.

The Source of Heat in the Sun. By Robert Hunt, F.R.S. Illustrated.

Soup for Children. By Baron Liebig.

Vegetables, Fruit, and Water as Sources of Intestinal Worms. By T. Spencer Cobbold, M.D., F.R.S. Illustrated.

The Anatomy and Physiology of the Foraminifera. By W. C. Williamson, F.R.S. Illustrated.

Waves of Heat and Waves of Death. By B. W. Richardson, M.A., M.D., F.R.S.

Origin of our Kitchen Garden Plants. By Harland Coultas.

The Geology of Mineral Springs. By Francis T. Bond, M.D., B.A., F.C.S. Illustrated.

Train-signalling in Theory and Practice. By Charles V. Walker, F.R.S., F.R.A.S.

Extract of Meat. By Charles Boner.

On Sponges. By Robert Patterson, F.R.S.

The Physical Phenomena of other Worlds. By Robert Hunt, F.R.S.

Migration of European Birds. By A. Leith Adams, M.A., M.B., F.G.S., F.L.S.

On the Difficulties in Identifying many of the Lower Kinds of Algae. By J. Braxton Hicks, M.D., F.R.S.

On the Oldest known Fossil (*Bozoon Canadense*) of the Laurentian Rocks of Canada; its Place, Structure, and Significance. By Prof. T. Rupert Jones, F.G.S.

Balloon Ascents and their Scientific Importance. By James Glaisher, F.R.S. Illustrated.

On the Highest Magnifying Powers and their Uses. By Lionel S. Beale, M.B., F.R.S.

Darwin's Observations on the Physiology of the Process of Fertilization in Plants. By M. C. Cooke. Illustrated.

What is a Tonic? By Edward Divers, M.D., F.C.S.

The Eruption of Etna. By S. J. Mackie, F.G.S.

Inside the Eye: the Ophthalmoscope and its Uses. By Ernest Hart, Ophthalmic Surgeon and Lecturer on Ophthalmic Surgery and Medicine at St. Mary's Hospital Medical School. Illustrated.

Is the Whitebait a Distinct Species? By the Editor. Illustrated.

On the Means of Communicating between Guards and Passengers on Railways. By Thomas Symes Pridcaux.

Atlantic Telegraphy. By Robert Hunt, F.R.S. Illustrated.

On Pure Water. By Edwin Lankester, M.D., F.R.S.

On the Nature and Diagnostic Value of Raphides and Other Plant Crystals. By Prof. Gulliver, F.R.S.

On Lake Basins. By Prof. D. T. Ansted, M.A., F.R.S. Illustrated.

Epidemics Past and Present: their Origin and Distribution.

On the Microscopic Anatomy of an Insect Larva (*Corethra plumicornis*). By E. Ray Lankester. Illustrated.

The Moon. By James Breen, F.R.A.S. Illustrated.

Photography, and some of its Applications. By the Editor.

Popular Science Review, Vol. 5 contains:—

Insects Injurious to the Turnip Crops. By Rev. W. Houghton, M.A., F.L.S.

Coffee. By Baron Liebig.

Australia and Europe formerly One Continent. By Berthold Seeman, Ph.D., F.L.S., V.P.A.S.

On Ozone in Relation to Health and Disease. By B. W. Richardson, M.A., F.R.S.

Glaciers and Ice. By W. F. Barrett. Illustrated.

On the Phenomena of Motion, Sensitiveness in Climbing Plants. By Rev. G. Henslow, M.A., F.L.S. With Page Illustration.

On the Construction and Use of the Spectrum Microscope. By H. C. Sorby, F.R.S.

On the Volvox Globator. By J. Braxton Hicks, M.D. (Lond.), F.R.S.

Engraving with a Sunbeam. Woodbury's Relief-printing. By J. Traill Taylor. Illustrated.

Entozoon-like Bodies in the Muscles of Animals destroyed by Cattle Plague. By Lionel S. Beale, M.B., F.R.S.

Our House Spiders. By J. Blackwall, F.L.S.

Raised Beaches, and their Origin. By Edward Hull, B.A., F.G.S.

On Milk, and its Adulteration. By Augustus Voelcher, Ph.D., F.C.S., &c.

The Amœba: its Structure, Development, and Habits. By Prof. W. C. Williamson, F.R.S.

On the Solfatara and Fumaroles in the Neighbourhood of Naples. By Prof. D. T. Ansted, M.A., F.R.S.

The Graphotype Process. By the Editor. Illustrated.

Hydræ or Fresh-water Polypes. By Rev. W. Houghton, M.A., F.L.S.

How to Work with the Telescope. Part I. By R. A. Proctor, B.A., F.R.A.S.

On the Exhaustion of our Coal. By Leonard Lemoran, M.E.

On Hybridization among Plants. By Rev. G. Henslow, M.A., F.L.S.

On the Light-emitting Apparatus of the Glowworm. By Henry Fripp, M.D.

Sun-force and Earth-force. By Dr. Richardson.

The Eruption of Santorin. By Prof. D. T. Ansted, F.R.S.

On the Movements of the Diatomaceæ. By E. Ray Lankester.

Aerolites. By Townshend M. Hall, F.G.S.

On the Electrical Principles of the Atlantic Telegraph. By Prof. G. C. Foster.

The Bone Caverns of Gibraltar, Malta, and Sicily. By A. Leith Adams, A.M., M.B., F.G.S., &c.

Notes on *Lophopus crystallinus*. By J. Josselyn Ranson and T. Graham Ponton.

Genesis or Parthenogenesis? By H. E. Fripp, M.D.

Modern Views of Denudation. By Edward Hull, B.A., F.G.S.

How to Work with the Telescope. Part II. By Richard A. Proctor, B.A., F.R.A.S.

Popular Science Review, Vol. 6 contains :—

- On Growth of some of the Algae. Illustrated by a few of the Common kinds. By J. Braxton Hicks, M.D., F.R.S.
- The Geology of Sinal. By Rev. E. W. Holland, M.A.
- The Planet Mars in January, 1867. By Richard A. Proctor, B.A., F.R.A.S. Illustrated.
- On Water-filters. By Edward Divers, M.D., F.C.S. Illustrated.
- Our Fresh-water Entomostraca, Shell Insects, or Water-Fleas. By W. Baird, M.D., F.L.S. Illustrated.
- How to Photograph Microscopic Objects. By Edward T. Wilson, M.B. Oxon. Illustrated.
- Recent Discoveries in Insect Embryogeny. By Henry Fripp, M.D. Illustrated.
- On the Struggle for Existence among Plants. By J. D. Hooker, M.D., F.R.S.
- How to Study Meteorology. By G. T. Chambers, F.R.A.S. Illustrated.
- On Sensitive Flames. By W. F. Barrett. Illustrated.
- Paraffin Lamps and their Dangers. By John Atfield, Ph.D., F.C.S.
- Venus's Flower-basket (Euplectella). By Dr. J. E. Gray, F.R.S. Illustrated.
- An Attempt to Approximate the Date of the Flint Flakes of Devon and Cornwall. By Spence Bate, F.R.S. Illustrated.
- Jupiter without his Satellites. By Richard A. Proctor, B.A., F.R.A.S. Illustrated.
- Fitzroy Weather Forecasts. By G. F. Chambers, F.R.A.S. Illustrated.
- On Life Insurance and Vital Statistics. By W. Hardwicke, M.D.
- New Electro-Magnetic Machines. By S. J. Mackie. Illustrated.
- The Botany of a Coal-mine. By Wm. Carruthers, F.L.S. Illustrated.
- The Microscope in Geology. By David Forbes, F.R.S. Illustrated.
- Why the Leaves Fall. By Maxwell T. Masters, M.D., F.L.S.
- A Message from the Stars. By Robert Hunt, F.R.S.
- On the Planarie of our Ponds and Streams. By E. Ray Lankester. Illustrated.
- Ventilation and Ventilators. By the Editor. Illustrated.
- Physics of the Brain. By B. W. Richardson, M.A., M.D., F.R.S.

Popular Science Review, Vol. 7 contains :—

- The Common Fresh-water Sponge (*Spongilla fluviatilis*). By Prof. W. C. Williamson, F.R.S. Illustrated.
- The Hurricane, the Typhoon, and the Tornado. By Prof. D. T. Ansted, F.R.S.
- Sensitive Plants. By Maxwell T. Masters, M.D., F.L.S. Illustrated.
- The former range of the Reindeer in Europe. By W. Boyd Dawkins, M.A., F.R.S.
- The Science of a Snow-flake. By Robert Hunt, F.R.S. Illustrated.
- The Food of Plants. By Prof. Church, M.A., F.C.S.
- The Gems and Precious Stones of Great Britain. By J. Morris, F.G.S. Illustrated.
- Is the Fresh-water Sponge (*Spongilla*) an Animal? By John Hogg, M.A., F.R.S.
- How to Dissect a Flower. By M. C. Cooke. Illustrated.
- The Polariscope and how to Work with it. By C. Hockin, M.A. Illustrated.
- Free Nematoids. By H. Carlton Bastian, M.D., F.R.S. Illustrated.
- On the Animals which are most nearly intermediate between Birds and Reptiles. By Prof. Huxley, LL.D., F.R.S. Illustrated.
- The Study of Chemical Geology. By David Forbes, F.R.S., &c.
- Plants known by their Pollen-grains and other Cells. By George Gulliver, F.R.S.
- The Great Eclipse of August 17, 1868. By R. A. Proctor, B.A., F.R.A.S. Illustrated.
- On the Range of the Mammoth. By W. Boyd Dawkins, M.A., F.R.S.
- Caddis-worms and their Metamorphoses. By Rev. W. Houghton, M.A., F.L.S. Illustrated.
- What is Wine? By August Dupré, Ph.D.
- Iron Shields and Iron Forts. By S. J. Mackie, Assoc. Inst. C.E. Illustrated.
- The Air or Swimming Bladder of Fishes. By Rev. W. Houghton, M.A., F.L.S.
- How to make a Geological Section. By Prof. D. T. Ansted, M.A., F.R.S. Illustrated.
- The Life of a Seed. By Maxwell Masters, M.D., F.L.S. Illustrated.

Popular Science Review, Vol. 8 contains :—

- Flying Machines. By Fred. W. Breary, Hon. Sec. to Aëronautical Society.
- The Compound Eye of Insects and Crustacea. By Henryripp, M.D. Illustrated.
- True and False Flint Weapons. By N. Whitley, C.E.
- The Planet Mars in February, 1869. By Richard A. Proctor, B.A., F.R.A.S. Illustrated.
- On the Molecular Origin of Infusoria. By J. Hughes Bennett, M.D., F.R.S.E. Diagrams.
- The Cuttle-fish. By St. George Mivart, F.Z.S. Illustrated.
- The Nature of the Interior of the Earth. By David Forbes, F.R.S.
- On the Use and Choice of Spectacles. By R. Brundell Carter, F.R.C.S. Diagrams.
- The Use of the Spectroscope in Astronomical Observation. By Richard A. Proctor, B.A., F.R.A.S. Illustrated.
- The British Lion. By W. Boyd Dawkins, M.A., F.R.S.
- Passion-flowers. By Maxwell T. Masters, M.D., F.L.S. Illustrated.
- The Natural Development of Bacteria in the Protolamie Parts of Various Plants. By M. A. Bechamp.
- The Planet Saturn in July, 1869. By R. A. Proctor, B.A., F.R.A.S.
- The Sertularian Zoophytes of our Shores. By Rev. T. Hincks, B.A. Illustrated.
- Hydrogenium. By Robert Hunt, F.R.S.
- The Structure and Affinities of the Squirts (Tunicata). By J. C. Galton, M.A., F.L.S. Illustrated.
- The Fertilization of Salvia, and of some other Flowers. By William Ogle, M.D. Illustrated.
- "In Articulo Mortis." By Benjamin W. Richardson, M.D., F.R.S.
- Experimental Illustrations of the Modes of Determining the Composition of the Sun and other Heavenly Bodies by the Spectrum. By Wm. Allen Miller, M.D., D.C.L., V.P.R.S. Illustrated.
- What is Bathybius? By Prof. Williamson, F.R.S.
- Are there any Fixed Stars? By Richard A. Proctor, B.A., F.R.A.S. Illustrated.
- Kent's Hole. By W. Boyd Dawkins, M.A., F.R.S.
- The Lingering Admirers of Phrenology. By Prof. Cleland.
- The Anatomy of a Mushroom. By M. C. Cooke. Illustrated.
- The Chemistry of a Comet. By Edward Divers, M.D., F.C.S.

Popular Science Review, Vol. 9 contains :—

- The Heat of the Moon. By J. Carpenter, F.R.A.S.
- "Under Chloroform." By B. W. Richardson, M.D., F.R.S.
- The Deposits of the Atlantic in Deep Water and their Relation to the White Chalk of Cretaceous Period. By Prof. D. T. Ansted, M.A., F.R.S.
- What is Wine? By A. Dupré, Ph.D.
- The Fertilization of Certain Plants (Didymia). By W. Ogle, M.D., Oxon, F.R.C.P.
- On some Interesting Points in the History of the Polyzoa. By Rev. Thomas Hincks, B.A.
- The Structure of Rock Masses (Stratification Joints, Cleavage). By David Forbes, F.R.S., &c.
- The Planet Jupiter 1869-70. By Rev. T. W. Webb, M.A.
- Microscopic Test Objects under Parallel Light and Corrected Powers. By Rev. J. B. Reade, F.R.S., P.R.M.S.
- The Fertilization of Various Flowers by Insects (Compositæ, Ericaceæ, &c.). By W. Ogle, M.D., F.R.C.P.
- On the Sphærosira Volvox of Ehrenberg. By Prof. Williamson, F.R.S.
- The Turret Ships *Monarch* and *Captain*. By S. J. Mackie, C.E. Illustrated.
- The Usefulness of the Fifth in Music. By Rev. C. Hope Robertson, M.A.
- The Anatomy of the River Mussel. By John C. Garton, M.A., Oxon. F.L.S.
- On a Simple Decimal System for England. By Royston Pigott, M.A., M.D., Cantab., M.R.C.P., F.C.P.S., F.R.A.S.
- What Fills the Star Depths. By R. A. Proctor, B.A., F.R.A.S.
- On the Apparatus employed in Deep Sea Explorations on Board H.M.S. *Porcupine* in the Summer of 1869. By W. Lant Carpenter, B.A., B.Sc.
- The Geology of Mont Cenis Tunnel. By Prof. Ansted, F.R.S. With Map and Section.
- Greenwich Time and its Telegraphic Distribution. By W. Ellis, F.R.A.S.
- The Echinus, or Sea Urchin. By S. G. Mivart, F.R.S. Illustrated.
- The Sun's Corona. By R. A. Proctor, B.A., F.R.A.S.
- Machine Guns or Mitrailleuses. By S. J. Mackie, C.E.

Popular Science Review, Vol. 10 contains :—

Hitting the Mark; or, Cannon-balls and their Striking Velocity. By G. W. Royston Pigott, M.A., M.D. Illustrated.

Natural Selection Insufficient to the Development of Man. By the Rev. George Buckle, M.A.

Polymorphic Fungi. By M. C. Cooke, M.A. Illustrated.

The Eclipse Expeditions. By R. A. Proctor, B.A., F.R.A.S.

Notes on Butterflies. By Rev. C. Hope Robertson, F.R.M.S. Illustrated.

On Sleep. By Dr. Richardson, F.R.S.

The Discophores or Large Medusæ. By Rev. T. Hincks, B.A. Illustrated.

The Issues of the Late Eclipse. By J. Carpenter, F.R.A.S.

Grafting; its Consequences and Effects. By Maxwell T. Masters, M.D., F.R.S. Illustrated.

Coal as a Reservoir of Power. By Robert Hunt, F.R.S.

The Plymouth Breakwater Port. By S. J. Mackie, C.E. Plate.

The Structure of Rock Masses (Foliation and Striation). By David Forbes, F.R.S. Illustrated.

South Africa and its Diamonds. By T. Rupert Jones, F.G.S.

British Bears and Wolves. By W. Boyd Dawkins, M.A., F.R.S.

The "Lotus" of the Ancients. By M. C. Cooke, M.A. Illustrated.

Greenland. By William Pengelly, F.R.S. F.G.S.

Observations on Jupiter in 1870-71. By Rev. T. W. Webb, M.A., F.R.A.S.

The International Exhibition at South Kensington. By S. J. Mackie, C.E. Illustrated by Heliotype Process.

How Fishes Breathe. By J. C. Galton, M.A., M.R.C.S., F.L.S. Illustrated.

Mr. Crooke's New Psychic Force. By J. P. Earwaker.

The Moss World. By R. Braithwaite, M.D., F.L.S. Illustrated.

Theory of a Nervous Ether. By Dr. Richardson, F.R.S.

On Pleistocene Climate and the Relation of the Pleistocene Mammalia to the Glacial Period. By W. Boyd Dawkins, M.A., F.R.S., F.G.S. Illustrated.

Star Streams and Star Sprays. By R. A. Proctor, B.A., F.R.A.S.

Popular Science Review, Vol. 11 contains :—

Mimicry in Plants. By A. W. Bennett, M.A.B., Sc. F.L.S. Illustrated.

Recent Microscopy. By Henry J. Slack, F.G.S., Sec. R.M.S.

Experimental Researches on the Contraction of Rocks. By L. C. Miall. Illustrated.

Psychic Force and Psychic Media. By J. P. Earwaker, Merton Col., Oxford.

Strange News about the Solar Prominences. By Richard A. Proctor, B.A. Illustrated.

Madder Dyes from Coal. By Edward Divers, M.D., F.C.S.

On the Structure of Camerated Shells. By H. Woodward, F.G.S., F.Z.S. Illustrated.

On the Temperature and the Movements of the Deep Sea. By Dr. W. B. Carpenter, F.R.S.

The Eclipse of last December. By R. A. Proctor, B.A., Sec. R.A.S.

The Lithofracteur. By S. J. Mackie, E.C.C. Illustrated.

The Physiological Position of Alcohol. By Dr. Richardson, F.R.S.

The Nature of Sponges. By Henry J. Slack, F.G.S. Illustrated.

On the Probable Existence of Coal-measures in the South-east of England. By Joseph Prestwich, F.R.S., F.G.S. Illustrated.

Bud Variation. By Maxwell T. Masters, M.D., F.R.S.

An Account of a Ganoid Fish from Queensland (*Ceratodus*). By Dr. Günther, F.R.S. Illustrated.

Greenwich Observatory. By James Carpenter, F.R.A.S. Illustrated.

The Recent Fossil Man. By J. Morris, F.G.S., Prof. of Geology in University College, London. Illustrated.

The Hydroid Medusæ. By Rev. Thomas Hincks, B.A., F.R.S. Illustrated.

The First Chapter of the Geological Record. By David Forbes, F.R.S., &c.

Electrical Signalling and the Siphon Recorder. By J. Munroe, Assistant to Sir W. Thomson.

Spontaneous Movements in Plants. By A. W. Bennett, M.A.B., Sec. F.L.S. Plate LXXXIX.

News from the Stars. By Richard A. Proctor, B.A., F.R.A.S.

Life Form of the Past and Present. By Henry Woodward, F.G.S., F.Z.S.

All Microscopists who care to know what is going on at home or
abroad should take in—

THE

MONTHLY MICROSCOPICAL JOURNAL.

*Transactions of the Royal Microscopical Society, and Record of
Histological Research at Home and Abroad.*

THIS Journal is devoted exclusively to the interests of Microscopical Science in the widest and most accurate sense of the term. It contains not only the proceedings of the Royal Microscopical Society, but also embraces communications from the leading Histologists of Great Britain, the Continent, and America, with a comprehensive *résumé* of the latest Foreign Inquiries, Critical Reviews and Short Notices of the more important works, Bibliographical Lists, and Descriptions of all new and improved forms of Microscopes and Microscopic Apparatus; Correspondence on all matters of Histological Controversy; and finally, a Department of "Notes and Queries," in which the student can put such questions as may elicit the special information he desires to obtain.

The Editor has also made arrangements for the publication of the most important Papers read before Local Associations. Contributions requiring illustration are accompanied by most carefully-drawn Plain or Coloured Plates, and the text is printed in clear and legible type, thus affording the Microscopist a readable Monthly Record of all that takes place in the branches of science specially interesting to him. By thus providing a journal at once thoroughly scientific, advanced, and comprehensive, and issued at such short intervals as to meet the requirements of active investigators, the Publisher hopes to receive the support of all workers with the Microscope, and the assistance and co-operation of all who desire to possess a periodical which creditably represents the labours of British and Foreign Histologists.

The Monthly Microscopical Journal,

Volume 1, price 10s. 6d., 392 Pages of Letter-press, 17 whole-page Plates, and numerous Woodcuts, contains :—

- Structure of Papillæ and Termination of Nerves in Muscle of Common Frog's Tongue. By Dr. Maddox. With Plate.
- Relation of Microscopic Fungi to Cholera. By Dr. J. L. W. Thudichum.
- Heliostat for Photo-Micrography. By Dr. Maddox. With plate.
- A Modification of the Binocular Microscope. By M. Nachet. Illustrated.
- Heliostat for Photo-Micrography. By Lieut.-Colonel J. J. Woodward, M.D., U.S. Army Medical Department. With Plate.
- The Vital Functions of the Deep-Sea Protozoa. By Dr. G. C. Wallich.
- The Formation of Blastoderm in Crustacea. By M. van Beneden, Brussels.
- On the Classification and Arrangement of Microscopic Objects. By J. Murie, M.D., F.L.S.
- Immersion Objectives and Test-Objects. By John Mayall, jun., F.R.M.S.
- Notes on Mounting Animal Tissues. By H. Charlton Bastian, M.D., F.R.S.
- Some Undescribed Rhizopods from North Atlantic Deposits. By G. C. Wallich, M.D., F.L.S.
- On the Construction of Object-Glasses. By F. H. Wenham.
- On the Organ of Hearing in Mollusks. By M. Lacaze-Duthiers.
- On a New Infusoria. By J. G. Tatem, F.L.S.
- The Composite Structure of Simple Leaves. By John Gorham, M.R.C.S.
- The Construction of Object-Glasses for the Microscope. By F. H. Wenham.
- On *Triarthra longiseta*. By C. T. Hudson, LL.D.
- On a New Growing Slide. By C. J. Muller.
- Professor Owen's Views on Magnetic and Vital Forces. By Lionel S. Beale, M.B., F.R.S.
- Scale-bearing Poduræ. By S. J. McIntyre, F.R.M.S.
- On the Fibres of the Crystalline Lens of *Petromyzonini*. By George Gulliver, F.R.S.
- Two New Forms of Selenite Stages. By Frederick Blankley, F.R.M.S.
- Researches on the Constitution and Development of the Ovarian Egg of the *Sacculinæ*. By M. J. Gerbe.
- On the Simple Structure of Compound Leaves. By W. R. M'Nab, M.D. Edin.
- On the Microscopical Structure of some Precious Stones. By H. C. Sorby, F.R.S., &c.
- Construction of Object-Glasses. By F. H. Wenham.
- On the Rhizopoda, Primordial Type of Animal Life. By G. C. Wallich, M.D., F.L.S., &c.
- On the Red Blood Corpuscle of Oviparous Vertebrata. By William S. Savory, F.R.S.
- A Small Zoophyte-trough. By W. P. Marshall.
- Preparation of Rock Sections. By David Forbes, F.R.S., &c.
- Markings on the *Pleurosigma angulatum* and *Lepisma saccharina*. By J. B. Dancer, F.R.A.S.
- Notes on Zoosperms of Crustacea. By Alfred Sanders, M.R.C.S., F.R.M.S.
- Protoplasm and Living Matter. By Dr. Lionel S. Beale, F.R.S.
- On some New Infusoria from the Victoria Docks. By Wm. S. Kent, F.R.M.S.
- Parkeria* and *Loftusia*. By Dr. Carpenter, V.P.R.S., and H. B. Brady, F.L.S.
- The Microscope in Silkworm Cultivation. By M. Cornalia.
- On the Proboscis of the Blow-fly. By W. T. Suffolk, F.R.M.S.
- A New Universal Mounting and Dissecting Microscope. By W. P. Marshall, President of the Birmingham Natural History and Microscopical Society.
- On Crystals enclosed in Blowpipe Beads. By H. C. Sorby, F.R.S.
- On Free-swimming *Amœbæ*. By J. G. Tatem, Esq.
- Action of Anæsthetics on the Blood Corpuscles. By J. H. McQuillen, M.D., D.D.S.
- Note on the Blood-vessel System of the Retina of the Hedgehog. By J. W. Hulke, F.R.S.
- A New Process of Preparing Specimens of Filamentous Algae for the Microscope. By A. M. Edwards.

The Monthly Microscopical Journal,

Volume 2, with 344 Pages of Letterpress, 19 whole-page Plates, price
10s. 6d., contains:—

- On the Rectal Papillæ of the Fly. By B. T. Lowne, M.R.C.S. With Plate.
- On the Diatom Prism, and the True Form of Diatom Markings. By the Rev. J. B. Reade, M.A., F.R.S.
- Observations on the Supposed Cholera Fungus. By the Rev. M. J. Berkeley, M.A., F.L.S.
- On the Correlation of Microscopic Physiology and Microscopic Physics. By John Browning, F.R.A.S.
- Notes on Hydatina lenta. By C. T. Hudson, LL.D. With plate.
- Some Remarks on the Structure of Diatoms and Podura Scales. By F. H. Wenham.
- Structure of the Adult Human Vitreous Humour. By David Smith, M.D., M.R.C.S.
- On the Use of the Chloride of Gold in Microscopy. By Thomas Dwight, jun., M.D.
- On a Simple Form of Micro-Spectroscope. By John Browning, F.R.A.S.
- On the Structure and Affinities of some Exogenous Stems from the Coal-Measures. By W. C. Williamson, F.R.S. With plate.
- On the Battledore Scales of Butterflies. By John Watson, Esq. With Three Plates.
- On Methods of Microscopical Research. By Herr S. Stricker.
- On the Construction of Object-Glasses for the Microscope. By F. H. Wenham.
- Jottings from the Note-book of a Student of Heterogeny. By Metcalfe Johnson, M.R.C.S.
- A Supposed Mammalian Tooth from the Coal-measures. By T. P. Barkas, F.G.S.
- On Holtenia, a Genus of Vitreous Sponges. By Wyville Thompson, LL.D., F.R.S.
- Icospectroscopy. Results of Spectrum Analysis. By Jabez Hogg, F.L.S.
- Memorandum of Spectroscopic Researches on the Chlorophyll of Various Plants. By the late William Bird Herapath, M.D., F.R.S.
- Observations on Mucor Mucedo. By R. L. Maddox, M.D. With plate.
- Floscularia coronetta, a New Species, with Observations on some Points in the Economy of the Genus. By Charles Cubitt, Assoc. Inst. C.E., F.R.M.S. With two plates.
- On the Detection by the Microscope of Red and White Corpuscles in Blood-stains. By Joseph G. Richardson, M.D.
- On the Staining of Microscopical Preparations. By Dr. W. R. M'Nab.
- Some further Remarks on an Illumination for verifying the Structure of Diatoms, and other Minute Objects. By F. H. Wenham.
- On the Rhizopodal Fauna of the Deep Sea. By William B. Carpenter, M.D., V.P.R.S.
- On the Structure of the Stems of the Arborescent Lycopodiaceæ of the Coal-measures. By W. Carruthers, F.L.S. Illustrated.
- On the Development of the Ovum of the Pike. By E. B. Truman. Illustrated.
- On the Presence of Foraminifera in Mineral Veins. By Charles Moore.
- On the Relations of the Ciliary Muscle to the Eye of Birds. By Henry Lawson, M.D. Illustrated.
- Experiments on Spontaneous Generation. By Edward Parfitt, Curator of the Devon and Exeter Institution.
- The Histology of the Eye. By John Whitaker Hulke, F.R.S., F.R.C.S., Assistant-Surgeon to the Middlesex Hospital, and Surgeon to the Royal London Ophthalmic Hospital.
- On Collecting and Mounting Entomotrachea. By J. G. Tatem.
- Further Remarks on the Nineteen-Band Test-plate of Nobert, and on Immersion Lenses. By Col. Woodward.
- On High Power Definition, with Illustrative Examples. By G. W. Royston Pigott, M.A., M.D.
- My Experience in the Use of Microscopes. By Dr. H. Hagen.
- Further Remarks on the Plumules of Battledore Scales of some of the Lepidoptera. By John Watson.
- The Development of Organisms in Organic Infusions. By C. T. Staniland Wake, F.A.A.S.

The Monthly Microscopical Journal,

Volume 3, 334 Pages of Letter-press, 20 whole-page Plates, price 10s. 6d., contains:—

- Structure of the Scales of Certain Insects of the Order Thysanura. By S. J. McIntire, F.R.M.S.
- Organisms in Mineral Infusions. By C. Staniland Wake, F.A.L.S.
- Markings on the Podura Scale. By G. Royston Pigott, M.D.
- Cultivation, &c., of Microscopic Fungi. By R. L. Maddox, M.D.
- Jottings by a Student of Heterogeny. By Metcalfe Johnson, M.R.C.S. No. 2.
- The Mode of Examining the Microscopic Structure of Plants. By W. R. McNab, M.D. Edin.
- On the Microscopical Examination of Milk under certain Conditions. By J. B. Dancer, F.R.A.S.
- On the Stylet-Region of the Ommatoplean Proboscis. By W. C. McIntosh, M.D., F.R.S.E., F.L.S.
- On a Method of Measuring the Position of Absorption-bands with a Micro-Spectroscope. By J. Browning, F.R.A.S., F.R.M.S.
- On an Undescribed Stage of Development of *Tetrarhynchus corollatus*. By A. Sanders, M.R.C.S., F.R.M.S.
- On a New Instrument for Cutting Thin Sections of Wood. By M. Mouchet, Hon. F.R.M.S.
- On the Calcareous Spicula of the Gorgonaceæ their Modification of Form, and the Importance of their Characters as a Basis for Generic and Specific Diagnosis. By W. S. Kent, F.L.S.
- On Pollen; as an Aid in the Differentiation of Species. By C. Bailey, Esq.
- On Professor Listing's Recent Optical Improvements in the Microscope. By Dr. H. Hagen.
- On the Structure of the Stems of the Arborescent Lycopodiaceæ of the Coal-Measures. By W. Carruthers, F.L.S., F.G.C., Botanical Dept., Brit. Museum.
- The Mode of Examining the Microscopic Structure of Plants. By W. R. McNab, M.D. Edin.
- Description of some Peculiar Fish's Ova. By W. B. Carpenter, M.D.,
- On the Shell Structure of *Fusulina*. By W. B. Carpenter, M.D., F.R.S.
- On the Comparative Steadiness of the Ross and the Jackson Microscope-stands. By W. B. Carpenter, M.D., F.R.S.
- A New Method of Using Darker's Films. By Edward Richards.
- A New Tube-dwelling Stentor. By C. A. Barrett, M.R.C.S.
- A Contribution to the Teratology of Infusoria. By J. G. Tatem.
- The Polymorphic Character of the Products of Development of *Monas* Lens. By Metcalfe Johnson, M.R.C.S.E.
- On the Reparation of the Spines of Echinida. By W. B. Carpenter, M.D., V.P.R.S.
- On the Colouring Matters derived from the Decomposition of some Minute Organisms. By H. C. Sorby, F.R.S., &c.
- Cercarie, Parasitic on *Lymnæa stagnalis*. By Jabez Hogg, Hon. Sec., R.M.S.
- Experimental Researches on the Anatomical and Functional Regeneration of the Spinal Cord. By M. M. Masius and Van Lair.
- Observations on some Points in the Economy of *Stephanoceros*. By C. Cubitt, Assoc. Inst. C.E., F.R.M.S.
- Notes on Diatomaceæ. By Prof. Arthur Mead Edwards.
- The New Binocular Microscope. By Samuel Holmes.
- Reminiscences of the Early Times of the Achromatic Microscope. By J. S. Bowerbank, LL.D., F.R.S.
- On an Apparatus for Collecting Atmospheric Particles. By R. L. Maddox, M.D.
- The Magnesium and Electric Light as applied to Photo-Micrography. By Brevet Lieut.-Colonel J. J. Woodward, U. S. Army.
- Remarks on High Power Definition. By F. H. Wenham, R.M.S.
- On a New Critical Standard Measure of the Perfection of High Power Definition, as afforded by Diatoms and Nobert's Lines. By Dr. ~~W. B. Carpenter~~ Pigott.

The Monthly Microscopical Journal,

Volume 4, 342 Pages of Letter-press, 14 whole-page Cuts, price 10s. 6d., contains:—

- On Fungi and Fermentation. By J. Bell, F.C.S.
- On the Origin of the Colouring Matter in Mr. Sheppard's Dichroic Fluid. By E. Ray Lankester, B.A., F.R.M.S.
- Object-Glasses and their Definition. By F. H. Wenham.
- On the Optical Advantages of Immersion Lenses and the Use of Deviation Tables for Optical Research. By Royston Pigott, M.A., M.D.
- On *Synchaeta mordax*. By C. T. Hudson, LL.D.
- Notes on Diatomaceæ. By Prof. Arthur Mead Edwards.
- On an Erecting Binocular Microscope. By J. W. Stephenson, F.R.A.S., F.R.M.S.
- Further Remarks on the Oxycalcium Light, as applied to Photo-Micrography. By Brevet Lieut.-Col. J. J. Woodward.
- Cursory Remarks on the Podura Scale, *Lepidocyrtus curvicolis*, and *Degeeria domestica*, or the Speckled Variety. By R. L. Maddox, M.D.
- Description of a Simple Air Sieve. By Metcalfe Johnson, S.C.E.
- The Microscopic Structure of the Human Liver. By Dr. H. D. Schmidt, of New Orleans.
- Microscopic Examination of the Atmosphere. By Geo. Sigerson, M.D.
- Microscopical Examination of Rocks and Minerals. By S. Allport, F.G.S.
- On the Structure of the *Pleurosigma angulatum* and *Pleurosigma quadratum*. By John Anthony, M.D., Cantab.
- Object-Glasses and their Definition. By F. H. Wenham.
- The Ciliary Muscle and Crystalline Lens in Man. By J. W. Hulke, F.R.S.
- On the Preparation of Specimens of Soundings for the Microscope. By Prof. A. M. Edwards.
- Circulation of the Latex in the Laticiferous Vessels. By H. C. Perkins, M.D.
- On a New Species of Parasite from the Tiger. By T. Graham Penton, F.Z.S.
- On the Application of the Microscope to the Study of Rocks. By H. C. Sorby, F.R.S., &c.
- On the Focal Length of Microscopic Objectives. By C. R. Cross.
- The Patterns of Artificial Diatoms. By H. J. Slack, Sec. R.M.S.
- Ancient Water-Fleas of the Ostracodous and Phyllopodous Tribes Bivalved Entomostraca. By Professor T. Rupert Jones, F.G.S.
- On the Real Nature of Disease-Germs. By Lionel S. Beale, F.R.S.
- On the Histology of Minute Blood-vessels. By Brevet Lieut.-Col. Woodward.
- On the Formation of Microscopic in closed Cells. By A. W. Wills.
- The Ciliary Muscle and Crystalline Lens in Man. By J. W. Hulke.
- On the "Hexactinellidæ" or Hexradiate Spiculed Silicious Sponges taken in the "Norma" Expedition. By W. Saville Kent, F.Z.S.
- On a Mode of Ascertaining the Structure of Scales of *Thysanuradeæ*. By Joseph Beck, F.R.M.S.
- On the Advancing Aplanatic Power of the Microscope and New Double Star and Image Tests. By G. W. Royston Pigott, M.A., M.D.
- American Microscopes and their Merits. By C. Stodder.
- On a New Anchoring Sponge, *Dorvillia agariciformis*. By W. Saville Kent, F.Z.S.
- On Aplanatic Definition and Illumination, with Optical Illustrations. By G. Royston Pigott, M.A., M.D., &c.
- On Selecting and Mounting Diatoms. By Cap. F. H. Lang.
- On Certain Cattle-plague Organisms. By Boyd Moss, F.R.C.S.
- Notes on New Infusoria. By J. G. Tatem.

The Monthly Microscopical Journal,

Volume 5, 286 Pages of Letterpress, 20 whole-page Plates, contains:—

- Notes on Fluorescence v. Pseudodichroism. By the Late Rev. J. B. Reade, F.R.S., P.R.M.S.
- Notes on the Minute Structure of the Scales of Certain Insects. By S. J. McIntire, F.R.M.S.
- On an Optical Illusion Slide; Cracks in Silica Films. By Henry J. Slack, F.R.M.S.
- Object-Glasses and their Definition. By F. H. Wenham.
- On the Mounting of the Diatom Prism. By F. W. Griffin, Ph.D.
- On *Pterodina valvata*: A New Species. By C. T. Hudson, LL.D.
- Observations on the Use of the Aëroconoscope, or Air-Dust-collecting Apparatus. By R. L. Maddox, M.D.
- On the Employment of Colloid Silica in the Preparation of Crystals for the Polariscope. By Henry J. Slack, F.G.S., Sec. R.M.S.
- The Development of Phycocyan. By T. C. White, M.R.C.S.
- The Anatomy of the Round Worm (*Ascaris lumbricoides*, Linn.). By B. T. Lowne, M.R.C.S., Eng.
- On the History, Refractions, Definition, and Powers of Immersion Lenses and New Refractometers. By Royston Pigott, M.A., M.D.
- On Microscopical Appliances. By Dr. Royston Pigott.
- Recent Investigations into Minute Organisms. By H. J. Slack, Sec. R.M.S.
- On a New Form of Binocular Eye-piece and Binocular Microscope for High Powers. By C. D. Ahrens.
- On Crystalline Forms modified by Colloid Silica. By Henry James Slack, F.R.M.S.
- Object-Glasses and their Definition. By F. H. Wenham, R.M.S.
- Robert's Nineteenth Brand and its Observers. By Charles Stodder.
- On the Structure of the Podura Scale, and Certain other Test Objects by Photo-Micrography. By Lieut.-Col. D. M. Woodward.
- Microscopical Examination of Water for Domestic Use. By James Bell, F.C.S.
- On the Winter Habits of the Rotatoria. By C. Cubitt, F.R.M.S.
- The Magnifying Power of the Microscope. By Count Castracane.
- A Few Experiments bearing on Spontaneous Generation. By Metcalfe Johnson, M.R.C.S.E.
- On the Mode of Working out the Morphology of the Skull. By W. Kitchen Parker, F.R.S., President R.M.S.
- Linear Projection considered in its Application to the Delineation of Objects under Microscopic Observation. By C. Cubitt, F.R.M.S.
- Optical Appearances of Cut Lines in Glass. By Henry J. Slack.
- Transmutation of Form in Certain Protozoa. By Metcalfe Johnson, M.R.C.S.E.
- Microscopical Examination of Two Minerals. By Professor A. M. Edwards.
- Additional Observations concerning the Podura Scale. By Lieut. J. J. Woodward.
- Remarks on the Scales of some of the Lepidoptera, as "Test Scale" of *Lepidocyrtus curvicolis*. By R. L. Maddox, M.D.
- On the so-called Suckers of *Dytiscus* and the Pupvilli of Insects. By B. T. Lowne, M.R.C.S.

The Monthly Microscopical Journal,

Volume 6, 302 Pages of Letterpress and 18 whole-page Plates, price 10s. 6d., contains:—

- On Bog Mosses. By R. Braithwaite, M.D., F.L.S.
 Structure of Podura Scales. By F. H. Wenham, Vice-president R.M.S.
 On some New Parasites. By T. Graham Ponton, F.Z.S.
 On some Improvements in the Spectrum Method of Detecting Blood. By H. C. Sorby, F.R.S., &c.
 On the Cellular Structure of the Red Blood Corpuscle. By Joseph G. Richardson, M.D.
 On the Use of Nobert's Plate. By Assistant-Surgeon J. J. Woodward, U. S. Army.
 On the Employment of Dammin in Microscopy. By Professor Arthur Mead Edwards, New York.
 Experiments on Angular Aperture. R. B. Tolles.
 Mycetoma: the Madura or Fungus-foot of India. By Jabez Hogg, Hon. Sec. R.M.S.
 Diatomaceous Earth from the Lake of Valencia, Caracas. By A. Ernst, Esq., and H. J. Slack, F.G.S.
 The Silicious Deposit in Pinnulariæ. By Henry J. Slack, F.G.S., Sec. R.M.S.
 Observations and Experiments with the Microscope on the Chemical Effects of Chloral Hydrate and other Agents, on the Blood. By Thomas Shearman Ralph, M.R.C.S., England.
 Floscularia Cyclops; a New Species. By Charles Cubitt, F.R.M.S.
 Mr. Tolles's Experiments on Angular Aperture." By F. H. Wenham.
 On the Microscopical Structure and Composition of a Phonolite from the "Wolf Rock." By S. Allport, F.G.S., with Chemical Analysis by J. A. Phillips.
 On Spore-cases in Coals. By J. W. Dawson, LL.D., F.R.S.
 On a New Rotifer. By C. T. Hudson.
 On the Examination of Mixed Colouring Matters with the Spectrum Microscope. By H. C. Sorby, F.R.S.
 On Spectra formed by the passage of Polarized Light through Double-refracting Crystals seen with the Microscope. By Francis Deas, M.A.
 Remarks on some Parasites found on the Head of a Bat. By R. L. Maddox, M.D.
 Notes on the Resolution of *Amphipleura pellucida* by a Tolles Immersion $\frac{1}{4}$ th. By Assistant-Surgeon J. J. Woodward, U. S. Army.
 Are Melicerian. By Chas. Cubitt.
 Micro-ruling on Glass and Steel. By John F. Stanistreet, F.R.A.S., with Illustrative Remarks by Henry J. Slack.
 The Fungoid Origin of Disease and Spontaneous Generation. By Jabez Hogg, Hon. Sec. R.M.S.
 On an Improved Method of Photographing Histological Preparations by Sunlight. By J. J. Woodward.
 Hematozoa in Blood of Ceylon Deer. By Boyd Moss, M.D.
 Microscopical Fissures in the Masticating Surface of Molars and Bicuspids. By J. H. McQuillen, M.D.
 Transmutation of Form in Certain Protozoa. By Metcalfe Johnson.
 On Gnats' Scales. By Jabez Hogg, Esq., Hon. Sec. R.M.S.
 The Examination of Nobert's Nineteenth Band. By F. A. P. Barnard.
 An Incident in the Life of a Chelifer. By S. J. McIntire, F.R.M.S.
 On the Form and Use of the Facial Arches. By W. K. Parker, F.R.S., President R.M.S.
 On the Angular Aperture of Immersion Objectives. By Robt. B. Tolles, of Boston, U. S. A.
 Notes on Pedalion Mira. By C. T. Hudson, LL.D.
 Another Hint on Selecting and Mounting Diatoms. By Capt. F. H. Lang.
 The Monads' Place in Nature. By Metcalfe Johnson, M.R.S.C.E.
 Mapping with the Micro-Spectroscope, with the Bright-line Micrometer. By H. G. Bridge.
 Some Remarks on a "Note on the Resolution of *Amphipleura pellucida* by a Tolles immersion $\frac{1}{4}$ th. By Assistant-Surgeon J. J. Woodward, U. S. Army.
 Infusorial Circuit of Generations. By Theod. C. Hilgard.
 Notes on Professor James Clark's Flagellate Infusoria, with Description of New Species. By W. Saville Kent, F.Z.S., F.R.M.S.
 Instrument for Micro-ruling on Glass and Steel. By J. F. Stanistreet.
 On the Conjugation of *Amœba*. By J. G. Taten, Esq.
 Crystallization of Metals by Electricity under the Microscope. By Philip Braham, Esq.
 Infusorial Circuit of Generations. By Theod. C. Hilgard.
 On the Connection of Nerves and Chromoblasts. By M. Geo. Punnett.

The Monthly Microscopical Journal,

Volume 7, 294 Pages of Letterpress, 20 whole-page Plates, price 10s. 6d., contains :—

- Markings on Battledore Scales of some Lepidoptera. By John Anthony, M.D. Cantab., F.R.M.S.
 The Nerves of Capillary Vessels and their Probable Action in Health and Disease. By Dr. Lionel S. Beale, F.R.S., F.R.C.P.
 New Erecting Arrangement for Binocular Microscopes. By R. H. Ward, M.A., M.D.
 On a New Micrometric Goniometer Eye-piece for the Microscope. By J. P. Southworth.
 Action of Hydrofluoric Acid on Glass, viewed Microscopically. By H. F. Smith.
 On the Relation of Nerves to Pigment and Other Cells or Elementary Parts. By Dr. Lionel S. Beale, F.R.C.P.
 On the Structure of the Stems of the Arborescent Lycopodiaceæ of the Coal-Measures. By W. Carruthers, F.R.S.
 On Bog Mosses. By R. Braithwaite, M.D., F.L.S.
 The Advancing Powers of Microscopic Definition. By Dr. Royston Pigott, M.A., Cantab.
 Microscopic Object-Glasses and their Power. By Edwin Bicknell.
 Remarks on a Tolles Immersion $\frac{1}{4}$ th. By Edwin Bicknell.
 Maltwood's Finder supplemented. By W. K. Bridgman.
 On a New Micro-Telescope. By Prof. R. H. Ward.
 Mycetoma, the Fungus Disease of India. By Jabez Hogg, F.L.S.
 The American Spongilla, a craspedote, flagellate Infusorian. By H. James Clarke, A.B., B.S.
 Refractive Powers of Peculiar Objectives. By R. B. Tolles (U.S.).
 On the Development of Vegetable Organisms within the Thorax of Living Birds. By Dr. James Murie, F.L.S., F.G.S., &c.
 Remarks on the Finer Nerves of the Cornea. By Dr. E. Klein.
 Note on the Resolution of *Amphipleura pellucida* by certain Objectives made by R. and J. Beck and by William Wales. By Dr. J. J. Woodward, U. S. Army.
 Stephenson's Erecting Binocular. By J. W. Stephenson, F.R.A.S.
 On a Presumed Phase of Actinophryan Life. By J. G. Tatem.
 On the Various Phenomena exhibited by the *Podura*. Test under Microscopic Resolving Powers. By G. W. Royston Pigott, M.A., M.D.
 Researches on the First Stages of the Development of the Common Trout (*Salmo Fario*). By Dr. E. Klein.
 On the Classification and Arrangement of Microscopic Objects. By Dr. James Murie, F.L.S., F.G.S., &c.
 On Bichromatic Vision. By J. W. Stephenson, F.R.A.S., Treas. R.M.S.
 The Supposed Fungus on *Coleus Leaves*, and Notes on *Podisoma fuscum* and *P. juniperi*. By Henry J. Slack, F.G.S.
 Optical Curiosities of Literature. By the Rev. S. Leslie Brakey, M.A.
 On an Improved Reflex Illuminator for the Highest Powers of the Microscope. By F. H. Wenham.
 On a Silvered Prism for the Successive Polarization of Light. By J. W. Stephenson, F.R.A.S., Treas. R.M.S.
 Structure of Battledore Scales. By J. Anthony, M.D., F.R.M.S.
 Beale's Nerve Researches. Dr. Beale. in Reply to Dr. Klein.
 Crystallization of Metals by Electricity. By Philip Braham.
 On the Means of Distinguishing the Fibres of New Zealand Flax from those of Manila or Sisal by the Microscope. By Captain Hutton.

CHEAP NATURAL HISTORY PERIODICAL.

Hardwicke's Science-Gossip:

An Illustrated Medium of Interchange and Gossip for Students and Lovers of Nature,

About Animals, Aquaria, Bees, Beetles, Birds, Butterflies, Ferns, Fish, Fossils, Lichens, Microscopes, Mosses, Reptiles, Rocks, Seaweeds, Wildflowers.

Edited by J. E. TAYLOR, F.G.S., Author of "Half-hours at the Seaside," "Geological Stories," &c.

"This is a very pleasant journal, that costs only fourpence a month, and from which the reader who is no naturalist ought to be able to pick up a good fourpenny-worth of pleasant information. It is conducted and contributed to by expert naturalists who are cheerful companions, as all good naturalists are; technical enough to make the general reader feel that they are in earnest, and are not insulting him by writing down to his comprehension, but natural enough and direct enough in their records of facts, their questioning and answering each other concerning curiosities of nature. The reader who buys for himself their monthly budget of notes and discussions upon pleasant points in natural history and science, will probably find his curiosity excited and his interest in the world about him taking the form of a little study of some branch of this sort of knowledge that has won his readiest attention. For when the study itself is so delightful, and the enthusiasm it excites so genuine and well-directed, these enthusiasms are contagious. The fault is not with itself, but with the public if this little magazine be not in favour with a very large circle of readers."—*Examiner*.

Hardwicke's Science-Gossip, Vol. 1. Selection from the Index.*

Amceba, the, 45.
 Ants, 113, 116, 143, 179, 185, 234, 235, 239, 262, 263.
 Aphides, Swarms of, 287.
 Aquarian Difficulties, 154, 188, 213, 259.
 Badger, the, 87.
 Balance of Power, 193.
 Bees, 34, 41, 93, 137, 143, 166, 167, 185, 214, 257, 263, 286, 287.
 Bitten by a Viper, 131.
 Black Cradle, 270.
 Breeze-fly, the, 194.
 Bromley, and What I found there, 246.
 Cabbage Butterfly and its Metamorphoses, 30, 74.
 Cat-fleas, 278.
 Caterpillars, Brood of, 126, 168, 288.
 Chapter from the Life of a Volvox, 244.
 Circle of Life, 145.
 Cleaning Diatomaceæ, 52.
 Colour of Birds' Eggs, 39, 47, 87, 142, 231.
 Common Things Unknown, 88.
 Coral Reefs, 112, 220, 285.

Crickets, 42, 66, 84, 113, 128, 166.
 Cui Bono? 25.
 Dead Fly on the Window, 10.
 Diatoms, 27, 52, 85, 95, 114, 140, 148, 163, 167, 237, 250.
 Diet of Worms, 180, 214.
 Duckweeds, 5, 258, 286.
 Ferns, 20, 34, 37, 44, 66, 67, 93, 169, 114, 117, 187, 188, 190, 214, 291, 262, 284.
 Fly in Pike-fishing, 280.
 Fly Parasites, 93, 227.
 Foot of a Fly, 253.
 Four Years' Acquaintance with a Toad, 12.
 Gathering Seaweeds, 173.
 Gossamer Spiders, 151, 191, 213.
 Gossip about Mansuckers, 49.
 Green Drake-fly, 231.
 Hair worm, more Notes on the, 197.
 Hairs, Chapter on, 29.
 Hermit Rooks, 226.
 House Ants, 170, 239.
 House Fly, the, 82.
 Humming-bird Hawk-moth, 208.

* In making a selection from the Index to each volume, space precludes giving more than about one-tenth of each volume.

y-developed Flowers, 103.
 ace, 241.
 riam," 265.
 olds, 133.
 e of Starling, 13
 alcule, 58.
 s, 248.
 impet and Parasite, 122.
 ings, 52.
 ocket, 149.
 abæus, 98.
 ic Illumination, 130.
 114, 273, 283.
 Objects, 46, 65, 93, 94, 116,
 he Hawthorn, 198.
 222.
 ow to Grow, 124, 162.
 l, the, 79.
 ells, 106.
 ills, 272.
 he, 232.
 nals, 177.
 ettles, 275.
 Light, 224.
 s, Popular History, 100.
 rth thinking about, 127.
 stler, the, 242.

Sea-Anemones, 40, 155, 158, 167, 183
 190, 196, 213, 239, 260, 285, 286, 287.
 Sea-wrack, 204.
 Short Commons, 49.
 Silurus, European, 56.
 Six-spot Burnet, 119, 151.
 Snake-stones, 37, 61, 94.
 Spiders, 24, 36, 39, 86, 143, 151, 206,
 213, 215, 239, 256, 282.
 Spiracles of Insects, 199, 254.
 Splitters and Lumpers, 73.
 Strange Remedies, 85.
 The Deep, deep Sea, 169.
 Tit in Moustaches, 26.
 Toads, 12, 62, 87, 111, 114, 233, 256.
 Tom Tidler's Ground, 126.
 Variations in British Plants, 32, 228.
 Vipers, 2, 95, 108, 131, 143, 160, 191.
 What do Crickets eat? 113, 128, 166.
 What Katy did, 146.
 What to seek and what to avoid in the
 Choice of a Microscope, 267.
 What's your Hobby? 1.
 Why objects appear larger through the
 Microscope, 8, 45.
 Window Gardens, 92, 117, 141, 284.
 Wren, the Blue, 199.

icke's Science-Gossip, Vol. 2. Selection from the Index.

he, 223.
 f Form, 266.
 oads and Frogs, 47, 69, 94,
 , a Chapter of, 221.
 159, 213, 238, 272.
 14, 21, 22, 46, 66, 69, 74, 95,
 , 166, 191, 215, 239, 260, 261,
 ss, 81.
 thing of, 73.
 22, 47, 70, 71, 115, 119.
 1, 71, 88, 89, 183, 190, 279.
 rs, 219.
 ighfisher, 26.
 a, Star of, 115, 136, 163, 186.
 ie Little, 200, 277.
 f Grasses, 53.
 rvæ, 95, 109.
 alsam, 175.
 wl, 4.
 88, 255, 260, 276.
 rs, 133, 161, 176, 182, 213.
 f Anecdotes, 221.
 and Acalephs, 124.
 153.
 imalcule, 253.
 tch, 34, 75, 254, 278.
 eae, 101, 147.
 British, 62, 87, 108, 112, 133,
 281.
 shes, 268.

Fairy-ring Champignon, 225.
 Ferns, 46, 61, 95, 104, 173, 269.
 Fossil Plants, 37.
 Fossil Wood, 250.
 Fossil Wood in Flint, 15.
 French Marygold, 163.
 Galls and Gall Insects, 165, 215, 228.
 Gill Fans of Sabellæ, 29.
 Glow-worm, 15, 238, 243.
 Grey Mullet, the, 145.
 Hairs, Star-shaped, 248.
 Illuminators for High Powers, 32, 65,
 66.
 Imperfectly-developed Plants, 8.
 Insect Fungi, 127, 176.
 Insect Vovarium, 80, 118, 207.
 Insects, 41, 55, 91, 93, 112, 117, 147, 161,
 184, 185, 204, 213, 249, 282.
 Jute: what is it? 84.
 Ladybirds, 169.
 Lizard, the Common, 79.
 Mistletoe of the Oak, 152, 186, 212.
 Moa, the, 14.
 Mounting, 19, 20, 23, 33, 47, 93, 94, 114,
 125, 209, 245, 260, 263, 282, 283.
 Odd Fishes, 50, 99, 171.
 Our Club, 193.
 Periodic Phenomena, 49.
 Pests, and their Checks, 54.
 Pin-centres and Rose-centres, 196.
 Puff Balls, 270.
 Rural Natural History, 83, 163.

Salamander, Japanese, 130.
 Saw-fly, the Great, 181, 273.
 Scales of Insects, 55, 91, 112.
 SCIENCE-GOSSIP, 1, 135, 158.
 Serpents at Meals, 244.
 Shooting Stars, 274.
 Snails and their Houses, 195, 230.
 Spectrum Microscope, 52.
 Speedwells, 121.
 Spider-crabs and their Parasites, 178, 211.
 Spiders, 7, 22, 40, 63, 119, 138, 141, 169, 189, 201, 209, 213, 255, 277, 279.
 Spring, 97.
 Starch, 34.

Sticklebacks, 5, 45, 153, 165.
 Teachings of Natural Science, 251.
 Track of the Pygmies, 154, 203, 239, 251.
 Turtle, the Edible, 247.
 Under the Snow, 25.
 Vegetable Caterpillars, 176.
 Vegetable Fibres, 10.
 Venus, Observations on, 110.
 Vivarium for Insects, 80, 118, 207.
 Viviparous Fish, 241.
 Wasps, 22, 88, 116, 276.
 Water-Beetles, 183.
 Water-Fleas, 156.
 Wings of British Butterflies, 27.
 Woodpecker, 6, 41, 95, 119.

Hardwicke's Science-Gossip, Vol. 3. Selection from the Index.

- A Century ago, 127.
 Age of Niagara, 139.
 Amidst the Ruins, 31.
 Analogy of Smell, 59.
 Aquaria, dust on, 69, 117, 118, 141, 142.
Atropos, 51.
 Bitten by a Viper, 175, 199, 213.
 Blindworm, the, 179, 260.
 Blood Beetle, the, 27, 62, 71, 94.
 Bouquet from Helvellyn, 242.
 Bugs, 269, 276, 282.
 Century ago, 127.
 Chignon-fungus, the, 107.
 Cholera-fungus, the, 206.
 Cockroaches, to Kill, 166, 212, 280.
 Cornish Colloquies, 182.
 Crickets, how to get rid of, 263, 279, 281.
 Crocodile in England, 7, 41.
 Death's-head Moth, 190, 213, 214, 262.
 Death-watch, the, 29.
 Dermestes? what is, 28, 206.
 Diatoms, 9, 35, 81, 91, 103, 115, 153, 156, 180, 188, 228, 269.
 Disguises of Insects, 193, 233, 234, 261, 279.
 Dodo, the White, 5, 52.
 Dragon-fly, the, 225.
 Echoes from the Club, 231.
 Edible Bird's Nest, 39.
 Fangs of Spiders, 237, 270, 276.
 Feast of Roses, 145.
 Foraminifera, 36, 129, 215, 236, 263.
 Freshwater Sponge, 247.
 Freshwater Sticklebacks in Seawater, 38.
 Germination of the Toad-rush, 150.
 Hairs of Dermestes Larva, 28, 206.
 Hardy Foreign Ferns, 83.
 Helps to Distribution, 244.
 Hints to Object-mounters, 91, 139.
 Huxley on the Study of Natural History, 73.
 Is Lichen-growth detrimental to Trees? 241.
Leaf-mining Larvæ, 169, 212.
Left by the Tide, 217.
 Lichen Dyes, 266.
 Maple, Aphids of the, 204.
 May Mushrooms, 112, 136.
 Melicerta, N. E. Green on, 33.
Mermis nigrescens, 221.
 Monmouth Deposit, 133, 156, 180.
 Mosquitoes, 78.
 Mounting in Balsam and Chloroform, 8, 21, 23.
 My little Green Monkey, 179.
 Nest of Wood Wasp, 247.
 Organization of Mosses, 249.
 Oxlip, the, 137, 163, 165, 197, 235.
 Perils of a Naturalist, 15.
 Poduræ, 45, 53.
 Preservation of Fossils, 22, 23, 44.
 Primeval Britain, 198.
 Primroses, 42, 114, 136, 141, 167, 235.
 Ramble in South Africa, 154.
 Rare Birds, Shooting, 69, 93, 94, 160.
 Rhythm of Flames, 49, 95.
 Rural "Folk-lore," 177.
 Rural Natural History, 86, 117, 118.
 Sandstone, Markings in, 20.
 Skeleton Leaves, 22, 117, 141, 246.
 Skeleton of Purple Urchin, 82.
 Snakes, 273.
 Sociable Mites, 124.
 Something to do, 99.
 Spiders' Nests, 37, 40.
Spirogyra, 60.
 Sponge-washings, 228.
 Sticklebacks in Seawater, 38, 87.
 Swallows, 50, 101.
 Temperature of Lakes, 272.
 Toad-flaxes, 201.
 Travellers' Tales, 25.
 Unity of Mankind, 110, 152, 173, 243, 269.
 Viper, the, 22, 175, 199, 280.
 Wasp, Sting and Poison-gland of, 60.
 What's in the Honey? 30, 68.
 Wheat Mildew, 16.
 White Dodo, the, 5, 52.
 Winter Work, 1.

ke's Science-Gossip, Vol. 4. Selection from the Index.

" 248.
 Ant of Texas, 1.
 t never Die, 16, 40, 62, 106.
 88, 117, 118, 139, 143, 159,
 19, 234, 261, 263, 282.
zfor, 194.
 41, 64, 88, 95, 160, 185, 191,
 k," 232.
 , 46, 214.
 Plumules of, 44, 137, 186,
 69.
 ns and their Cases, 152, 189.
 , a, 183.
 s, 15, 22, 215, 239.
 ottles, 111.
 i Hedge-sparrow, 113, 143,
 14, 261.
 s, 158.
 legs, 256.
 e Heart of, 227, 279.
 , what is, 241.
 a, the, 87, 113.
 s, 117, 226.
 the Pupa of a, 245.
 s, Phenomena of, 217.
 , 221.
 61, 162, 183, 187, 213, 231,
 140, 261, 263, 281.
 10ts, 97.
 of Fern Seeds, 183.
 i, the, 53.
 Actinia, a, 247.
 9, 94, 206, 213.
 , 49, 114, 160, 209, 271.
 es of the, 55.
 i, 58, 124, 143.
 the Study of British, 197,
 r, the Large Green, 196, 236.
 lian Bats, 26.
 he, 109, 160.
 Variation in the, 267.
 23, 69, 81, 100.
 229.
 the, 107.

How Birds and Insects fly, 9,
 Infusoria, 44, 57, 125, 155, 164.
 Irritability and Sensation, 25.
 Kingfisher, the Common, 204, 234
 King of the Rats, 135.
 Kite, the, 251.
 Maine Deposits, 85.
 Maple Blight, 136, 188.
 Merlin, the, 156.
 Metamorphoses of Insects, 35.
 Microscopic Seeds, 253.
 Mole Mite, 232.
 Mosquitoes, 207, 211, 212, 215, 236.
 Oxlip, the, 35.
 Palates of Mollusca, 20, 200.
 Pebble-finding, 134.
 Perley's Meadow Deposit, 131.
 Phantom Larvæ, 78.
 Polyzoön from Victoria Docks, 255.
 Primroses, Pink, 43, 66, 147, 187.
 Reptiles, from the Coal-Measures, 104,
 142, 167, 214.
 Reptiles in Confinement, 272.
 Sand Wasp, the, 205.
 Scleriform Tissue, 276.
 Sensorial Vision, 145.
 Silver-mining in Eastern Nevada, 193.
 Slug Parasite, 274.
 Smew, the, 55.
 Spicules of Echinoderms, 175.
 Spiders, 8, 11, 21, 22, 23, 24, 41, 47, 51,
 58, 63, 82, 124, 128, 143, 161, 168, 167,
 195, 213, 238, 261, 262, 263, 281, 283.
 Splits, 169.
 Spring Phenomena of Plant Life, 121.
 Stag-Beetle, the, 109.
 Stings and Poison-glands of Bees, &c.,
 148, 205.
 Storm-glass, 24, 93, 117, 143, 167.
 Trees, Age of, 202, 231, 259.
 Unity of Mankind, 6, 34.
 Vegetable Hairs, 11, 101.
 Vipers, 23, 46, 70, 95, 165, 180, 212.
 Waxwing, the, 181.
 Why? 265.
 Wood-sorrel, the, 52, 280.

Hardwicke's Science-Gossip, Vol. 5. Selection from the Index.

- Acherontia atropos*, 220, 257, 278.
 Admiral, the Red, 257, 262, 273.
 Age of Fish, 141, 214.
 Animal from the Salt Lake, 78, 130, 234.
 Aphid Lion and Lacewing Fly, 11.
 Ash-Trees, Violets under, 91, 116, 117, 166, 188, 189.
 Badger and Otter, the, 90, 118, 137, 258, 262, 277.
 Botanical Allusion in Tennyson, 91, 116, 117, 166, 189, 189.
 British Birds, 39, 85, 107, 113, 156, 179, 227, 253, 274.
 Buds as Objects for Winter Study, 34.
 Butterflies, 58, 116, 140, 164, 212, 273, 274.
 Cats and Starfishes, 214, 234, 239, 263, 282.
 Cats before a Storm, 117, 141, 164, 167.
 Celandine, 52.
 Cells for Microscopic Objects, 139, 236, 260, 281.
 Centipede, a Luminous, 46, 47, 69, 71.
Cheyleti, 5.
 Christmas and the Microscope, 44.
 Christmas Berries, 13, 94, 138.
 Cilia, about, 53.
 Coal, the Story of a Piece of, 1, 46, 71, 96.
Cochlearia officinalis, 43, 66, 91, 114, 143.
Comatula rosacea, 209.
 Cuckoo, the, 16, 64, 65, 185, 262.
 Death's-Head Moth, 220, 257, 278.
 Dendritic Spots on Paper, 22, 46, 71, 80.
 Diatomaceæ, 22, 61, 67, 72, 92, 109, 139, 158, 163, 183, 187, 220.
 Drawing from the Microscope, 87, 139, 165.
 Early Birds, 113, 136, 137.
 Enchanter's Nightshade, 62.
 English Plant Names, 25.
Epistylis, 83.
 Fish Scales, 12, 41, 67, 163, 187, 260, 281.
 Floral Giants, 9.
 Food for Bullfinch, 215, 237, 238.
Fragillaria Crotoneensis, 109, 158, 183.
 Frog, 63, 76, 161.
 Fumart, the, 22, 45, 68.
 Garden Decoration, Wild Flowers for, 169, 265.
Geophilus electricus, &c., 49, 65, 71.
Geranium Robertianum, 153, 191, 288.
 Geraniums, Carpels of, 211, 235, 238, 261.
 Gnats, 16.
 Hawk-Moths, 16, 119, 220, 234, 257, 273.
 Hawthorn, 22, 23, 43, 70, 93, 116, 118.
 Holly, the, 213, 235, 238, 259, 280, 283.
 Humble Bee, Winter Home of the, 41, 134, 164, 166.
 Hybernation of Bees, 41, 90, 93, 134, 164, 166.
Janthina, 31, 64.
 Influence of Light on Insects, 57, 77, 137, 165, 188, 273.
 Insects of the Season, 233, 234.
 Kestrel, the, 179, 257.
 Ladybird, 232, 239, 267, 283.
 Lampreys and Lamperns, 145.
 Lapwing, the, 107, 167.
 Laurel Berries, &c., 47, 70, 114.
 Laurel-Leaves, 20, 45, 68, 71.
 Leeches, 76, 93, 143, 161, 165.
Lepisma saccharina, 94, 118, 142, 163.
 Light Attracting Insects, 57, 77, 137, 165, 188.
 Ligurian Bees, 213, 237, 256, 263, 283.
 Luminous Centipede, 46, 47, 69, 71.
 Monsters of the Deep, 55.
 Mosquitoes, 17, 54.
 Moth, the Fish, 94, 118, 142.
 Myriapods, 49.
 Norfolk, Rare Birds in, 160, 161, 184, 192.
 Otter and Badger, the, 90, 137, 258.
 Otters, 90, 118, 137, 161, 165, 184.
Phronima, 73.
 Poppy Seeds, 11.
 Rudd, Scales of the, 12.
 Scales of Butterflies, 212, 214.
 Scurvy Grass, 43, 66, 67, 91, 114, 143.
 Sea Anemones, 56, 90, 198, 210.
 Sea Birds, Association for the Protection of English, 10, 42.
 Sections of Fossil Wood, 18.
 Siekin, the, 39.
 Starfish, Cats and, 214, 234, 239, 282, 282.
 Surirella, a New, 61.
 Vine Diseases, new, 59.
 Violets under Ash-Trees, 91, 116, 117, 166, 188, 189.

Hardwicke's Science-Gossip, Vol. 6. Selection from the Index.

- Abstinence of Insects, 64.
 Animated Oats, 199, 211, 212, 237.
 Antique, Studies from the, 14, 45, 71.
 Ants, 87, 185, 241, 263.
 Apterous Insects, 49, 105, 126, 164.
 Aquarium Difficulty, 40, 121, 142, 165, 179, 191, 215, 216.
 Bees, 21, 24, 34, 42, 47, 65, 70, 117, 119, 141, 142, 143, 161, 164, 165, 166, 167, 188, 190, 212, 213, 215, 236, 237, 257, 269.
 Beetles, 112, 113, 143, 233, 263.
 Birds, 13, 35, 54, 58, 65, 86, 108, 113, 132, 157, 159, 179, 184, 246, 276.
 Borago, 165, 189, 212, 213, 214, 239.
 Butterflies, 22, 143, 251, 283.
 Caterpillar, 47, 70, 149, 239.
 Cats, 16, 17, 23.
Charadrius plumalis, 58, 95, 114, 188.
Chenopodium Bonus Henricus, 189, 214, 234, 238.
Chrysopa perla, 231, 237, 239.
 Cleaning Shells, 190, 214.
 Cowper and the Nightingale, 142, 175, 209.
 Cowslips, 141, 164, 166, 190.
 Cuckoo, the, 108, 138, 281.
 Derivation of Foxglove, 43, 67, 69, 91, 93, 115, 118, 135, 166.
 Diatoms, 22, 37, 55, 61, 140, 178, 235.
 Dodder, the, 117, 118, 120.
 D'Orbigny's Foraminifera, 81, 106, 155.
 Dragon-fly in Town, 237, 262.
 Dun Cow, Ribs of the, 23, 38, 63, 69, 94, 96.
 Eggs, 92, 114, 132, 185, 191, 212, 251, 259, 283.
 Embryos, Tricotyledonous, 115, 139, 145, 226.
 English Plant-Names, 127, 210, 211, 227, 234.
 Female Moths, Attractive Power of, 153, 174, 213.
 Fish, Scales of, 92, 140, 187, 235, 279, 280.
 Flies, 65, 249, 273.
 Foraminifera, 9, 68, 81, 106, 155, 167, 209.
 Foxglove, 43, 67, 69, 91, 93, 115, 118, 135, 166, 260.
 Frog, the, 71, 119, 191, 233.
 Good King Harry, 189, 214, 234, 238.
 Hairs, 111, 116, 212, 235.
 Hawk-moths, 114, 161, 209, 229, 232, 233, 283.
 "Hoddy-doddy," 21, 70, 93, 94, 212.
 Holly Berries in July, &c., 186, 210, 213, 234.
 Insects, Brain of, 262.
 Insects, 8, 41, 49, 64, 76, 92, 105, 116, 126, 133, 137, 164, 224, 236, 247, 255, 262.
 July, Holly Berries in, 186, 210, 213, 234.
 Lotus, its History and Traditions, 124, 208, 231, 272.
 Manipulation, 20, 187.
Mergus albellus, 95, 110, 137.
 Mosquitoes, 167, 168.
 Mosses, 83, 139, 275.
 Otters, 21, 54, 90, 113, 114, 137, 138, 209.
Phacelia tenacetifolia, 167, 189, 211.
 Plant Names, 127, 210, 211, 227, 234, 262.
 Plover, the Golden, 58, 95, 114, 188.
 Podder, 91, 117, 118, 120.
Polyxenus lagurus, 187, 209.
 Pygidium of Lacewing Fly, 231, 237, 239.
 Queen of Spain Fritillary, the, 233, 258.
 Ribs of "the Dun Cow," 23, 38, 63, 69, 94, 96.
 Rue and Rosemary, 39, 40, 67, 118.
 "Sembling" of Moths, 153, 174, 213.
 Shells, New British, 130, 138, 161, 277.
 Smew, the, 95, 110, 137.
 Snow Bunting, the, 71, 90, 114.
Sphinx atropos, 114, 161.
 Studies from the Antique, 14, 45, 71.
 Study of Natural History, 112.
 Sandew, Hairs of the, 111, 212.
 Titmouse, the Bearded, 65, 95.
 Toad-flax, the Ivy-leaved, 43, 71.
 Toads devouring Bees, 117, 141, 142, 161, 164, 166.
 Tricotyledonous Embryos, 115, 139, 145, 226.
Veronica Busbaumii, 43, 91, 186.
 Wasps and Bees—Do Toads eat them? 117, 141, 142, 161, 164, 166.
 Wiltshire, Spur-winged Goose in, 51, 95.
 Wingless Insects, 49, 105, 126, 164.
 Woodpecker, Greater Spotted, 184.
 Woodruff, the Sweet, 47, 69, 70, 93.
 Zinc Troughs for Bees, 142, 167.
Zonites glaber, 161, 277.

Hardwicke's Science-Gossip, Vol. 8—continued.

Singing Mice, 47, 65, 94.
 Snake, the Smooth, 266, 232, 239, 258.
 Spicules of Sponges, &c., 20, 95, 186.
 Spiders, 140, 182, 213, 261.
 Sponges, 17, 20, 25, 56, 95, 106, 160, 185,
 281.
 Stag Beetle, 42, 92, 95, 139, 190.
 Starlings, 93, 117, 118, 140, 213, 237.
 Swallows, 57, 69, 115, 143, 212, 235, 237,
 239.
 Tissues, on Staining, 111, 120, 136.

Test-scales of Podura, 16, 41, 47, 100,
 266.
 Tufted Duck, the, 70, 94, 143, 164.
 Unicorn, the, 46, 68, 95.
Vitruva pellucida, 44, 69, 116.
 Wasps, 70, 163, 214, 239, 282.
 Web-weaving Caterpillars, 58, 94, 11
 142.
 Whitebait, 44, 263, 281, 282.
 Young Caterpillars in Confinement, 161,
 188, 237, 262.

HARDWICKE'S SCIENCE-GOSSIP,

*Published Monthly, price 4d. Each Volume can be had separately
 Price 5s., in purple cloth. All the numbers are kept in print.*


Volume 9 commenced January, 1873.

MR. HARDWICKE begs to inform Authors of Works on Natural History, Travel, General Science, and Miscellaneous Literature, that he has at his command the requisite means for bringing all Works published by him prominently before the Public, both at home and abroad. Being practically acquainted with Printing, and having been many years engaged in business requiring an intimate knowledge of the best modes of Illustration, he is enabled to offer great facilities to Gentlemen who entrust their Works to him.

Estimates of Cost, Terms of Publishing, and other particulars on application.

LONDON:

192, Piccadilly, W.



22

— 5 —

1







